



EUROfusion

Topic 2 I: AUG experiment analysis meeting

N. Vianello and V. Naulin for the Topic 2 I Scientific Team

15 September 2017



This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.



✓ L-Mode experiment, CW I7.


1. Performed similar density ramps in an I_p scan at constant q_{95}
2. Performed similar density ramps in an I_p scan at constant B_t



✓ L-Mode experiment, CW 17.


✓ H-Mode experiment, CW 21.

1. Compare divertor/midplane fueling effect on filamentary transport and profiles without cryo-pumps
2. Compare profiles with the same fueling with/without cryopumps
3. Determine an H-Mode with the cryopumps matching similar divertor pressure and SOL profiles




- ✓ We matched correctly the shape and the $L_{||}$ here shown from outer divertor plate up to X-point

L-Mode analysis: I_p scan at constant q_{95}



- ✓ The scan was performed with similar puffing rate (0.8-1 MA) whereas we reduced it at lower current to avoid early disruption
- ✓ We have comparable edge density and divertor neutral pressure

L-Mode analysis: I_p scan at constant q_{95}



- ✓ At comparable edge density
Upstream profiles are different
with the tendency to develop
shoulder easier at lower current