



## Topic 21: AUG experiment Week 17 Summary

---

Topic 21 Scientific Team

08 May 2017



This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.



bla bla bla

# Objective of Week 17 campaign



- ✓ Check shoulder formation and filamentary transport behavior at different current keeping  $q_{95}$  constant in L-Mode

# Objective of Week 17 campaign




- ✓ Check shoulder formation and filamentary transport behavior at different current keeping  $q_{95}$  constant in L-Mode
- ✓ Check shoulder formation and filamentary transport behavior at different current keeping  $B_t$  constant in L-Mode

# Objective of Week 17 campaign




- ✓ Check shoulder formation and filamentary transport behavior at different current keeping  $q_{95}$  constant in L-Mode
- ✓ Check shoulder formation and filamentary transport behavior at different current keeping  $B_t$  constant in L-Mode
- ✓ Find suitable H-Mode scenario at *high power* (compatible with probe operation) for shoulder behavior and filamentary transport studies in H-Mode

# L-Mode: Current scan at constant $q_{95}$




- ✓ Reference shot with **EOC** Shape, 0.8MA,  $B_\phi = -2.5$  T, 0.5MW NBI heating, fueling in order to keep heating approximately constant during the current scan

# L-Mode: Current scan at constant $q_{95}$




- ✓ Reference shot with **EOC** Shape,  $0.8\text{MA}$ ,  $B_\phi = -2.5\text{T}$ ,  $0.5\text{MW}$  NBI heating, fueling in order to keep heating approximately constant during the current scan
- ✓ The fueling has been reduced at lower current for not encountering disruption too early

# L-Mode: Current scan at constant $q_{95}$




- ✓ Reference shot with **EOC** Shape,  $0.8\text{MA}$ ,  $B_\phi = -2.5\text{T}$ ,  $0.5\text{MW}$  NBI heating, fueling in order to keep heating approximately constant during the current scan
- ✓ The fueling has been reduced at lower current for not encountering disruption too early
- ✓ Up to 3 seconds 0.6 and 0.8 discharge exhibits close edge and core density as well as neutral pressure from gauges under the dome

# L-Mode: Current scan at constant $q_{95}$




- ✓ Reference shot with **EOC** Shape, 0.8MA,  $B_\phi = -2.5$  T, 0.5MW NBI heating, fueling in order to keep heating approximately constant during the current scan
- ✓ The fueling has been reduced at lower current for not encountering disruption too early
- ✓ Up to 3 seconds 0.6 and 0.8 discharge exhibits close edge and core density as well as neutral pressure from gauges under the dome
- ✓ In all the cases the profiles evolve and tends to become flatter. Preliminary indication suggest this occurs at similar  $\Lambda_{div}$

# L-Mode: Current scan at constant $B_t$




- ✓ The fueling has been reduced at lower current for not encountering disruption too early

# L-Mode: Current scan at constant $B_t$




- ✓ The fueling has been reduced at lower current for not encountering disruption too early
- ✓ Up to 3 seconds 0.6 and 0.8 MA discharges exhibits close edge and core density as well as neutral pressure from gauges under the dome

# L-Mode: Current scan at constant $B_t$




- ✓ The fueling has been reduced at lower current for not encountering disruption too early
- ✓ Up to 3 seconds 0.6 and 0.8 MA discharges exhibits close edge and core density as well as neutral pressure from gauges under the dome
- ✓ In all the cases the profiles evolve and tends to become flatter. Work in progress to evaluate development as a function of divertor collisionality  $\Lambda_{\text{div}}$


# Fluctuations



# H-Mode




# H-Mode




- ✓ **EOC** shape, initially with 2MW NBI + 1.4MW ECRH density ramp without N seeding
- ✓ We increase the NBI power up to 4MW with the same ECRH adding N in feedforward

# H-Mode




- ✓ **EOC** shape, initially with 2MW NBI + 1.4MW ECRH density ramp without N seeding
- ✓ We increase the NBI power up to 4MW with the same ECRH adding N in feedforward
- ✓ We end up in a degraded H-Mode with all the 4 phases. Fluctuations from MEM available in all the phases

# H-Mode




- ✓ **EOC** shape, initially with 2MW NBI + 1.4MW ECRH density ramp without N seeding
- ✓ We increase the NBI power up to 4MW with the same ECRH adding N in feedforward
- ✓ We end up in a degraded H-Mode with all the 4 phases. Fluctuations from MEM available in all the phases
- ✓ We reduce the N dosing finding a proper scenario **but without cryopumps**




- ✓ **EOC** shape, initially with 2MW NBI + 1.4MW ECRH density ramp without N seeding
- ✓ We increase the NBI power up to 4MW with the same ECRH adding N in feedforward
- ✓ We end up in a degraded H-Mode with all the 4 phases. Fluctuations from MEM available in all the phases
- ✓ We reduce the N dosing finding a proper scenario **but without cryopumps**
- ✓ In this cases we still encounters a degraded H-Mode but without all the phases

# H-Mode



- ✓ **EOC** shape, initially with 2MW NBI + 1.4MW ECRH density ramp without N seeding
- ✓ We increase the NBI power up to 4MW with the same ECRH adding N in feedforward
- ✓ We end up in a degraded H-Mode with all the 4 phases. Fluctuations from MEM available in all the phases
- ✓ We reduce the N dosing finding a proper scenario **but without cryopumps**
- ✓ In this cases we still encounters a degraded H-Mode but without all the phases
- ✓ With respect to the reference we achieve higher power and higher neutral pressure in the divertor. We will need to adjust once the cryopump will operate in the next session

# Fast particle acceleration



- ✓ L-mode shots: no evidence of electron or ion acceleration during flat-top, but TAE excitation (top) & neutron time traces during current ramp-up in three shots suggest bulk ion acceleration
- ✓ H-mode shots: many examples found of SXR spikes from plasma edge at start of ELMs → electron acceleration), although enhancements are less dramatic than in earlier H-mode pulses with lower pedestal  $\nu^*$  (lower plot)
- ✓ ELMs also linked to enhanced fast ion losses, as in previous H-mode pulses – not yet clear whether beam ions are accelerated during ELMs, but fast ion loss detector settings will be optimised in remaining pulses of this experiment

K. McClements & J. Galdon-Quiroga