

Filamentary transport in high-power H-mode conditions and in no/small-ELM regimes to predict heat and particle loads on PFCs for future devices

J. Madsen and N. Vianello
15 November 2016







 Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics



- Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics
- 2. The balance is observed to change at high density in L-Mode and this is at the basis of the *shoulder formation*



- Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics
- The balance is observed to change at high density in L-Mode and this is at the basis of the shoulder formation
- 3. This is observed in a variety of devices including AUG (Carralero et al. 2014), TCV (Garcia et al. 2007) and MAST (Militello et al. 2016)



- Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics
- The balance is observed to change at high density in L-Mode and this is at the basis of the shoulder formation
- This is observed in a variety of devices including AUG (Carralero et al. 2014), TCV (Garcia et al. 2007) and MAST (Militello et al. 2016)
- 4. The observations exhibit a dependence on  $I_p$  (increasing the current at constant  $B_\phi$  shoulder tends to disappear)



- Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics
- The balance is observed to change at high density in L-Mode and this is at the basis of the shoulder formation
- This is observed in a variety of devices including AUG (Carralero et al. 2014), TCV (Garcia et al. 2007) and MAST (Militello et al. 2016)
- The observations exhibit a dependence on I<sub>p</sub> (increasing the current at constant B<sub>φ</sub> shoulder tends to disappear)
- 5. The H-Mode behavior is still a work in progress and exhibit a dependence on the level of fueling




- Recent experimental and theoretical investigation agrees on suggesting SOL density profiles is determined by a balance between radial convective filamentary transport and parallel dynamics
- The balance is observed to change at high density in L-Mode and this is at the basis of the shoulder formation
- This is observed in a variety of devices including AUG (Carralero et al. 2014), TCV (Garcia et al. 2007) and MAST (Militello et al. 2016)
- The observations exhibit a dependence on I<sub>p</sub> (increasing the current at constant B<sub>φ</sub> shoulder tends to disappear)
- 5. The H-Mode behavior is still a work in progress and exhibit a dependence or the level of fueling
- 6. L-Mode and H-Mode behavior are not yet reconciled and different devices show somehow contradictory behavior

## L-Mode studies




✓ AUG and JET (Carralero et al. 2015) suggest that \(\Lambda\_{div}\) dominates this process and a transition from sheath-limited to inertial regime



### L-Mode studies



- ✓ AUG and JET (Carralero et al. 2015) suggest that \(\Lambda\_{div}\) dominates this process and a transition from sheath-limited to inertial regime
- √ Tested by changing n<sub>e</sub> and T<sub>e</sub> through fueling/seeding/heating



### L-Mode studies



- ✓ AUG and JET (Carralero et al. 2015) suggest that  $\Lambda_{div}$  dominates this process and a transition from sheath-limited to inertial regime
- √ Tested by changing n<sub>e</sub> and T<sub>e</sub> through fueling/seeding/heating
- √ This determines a change of the velocity-size scaling

