

Observational evidence of star formation stochasticity in the CALIFA dataset

Nicolás Romero Díaz¹

Advisor: Jaime Forero-Romero¹

¹Departamento de Física
Universidad de los Andes

Undergraduate thesis presentation
December 1 2016



Table of Contents

1 Motivation

- Stochasticity in star formation
- Measuring stochasticity

2 The CALIFA survey

- CALIFA datacubes
- The Pipe3D pipeline

3 Results

4 Conclusions



Table of Contents

1 Motivation

- Stochasticity in star formation
- Measuring stochasticity

2 The CALIFA survey

- CALIFA datacubes
- The Pipe3D pipeline

3 Results

4 Conclusions



Stochasticity in star formation

Star formation

- Star forming regions
- Young stars ionize gas clouds
- Spectral quantities: equivalent width

$$\text{EW} \equiv \frac{L_{Ly\alpha}}{L_{\lambda,UV}} \quad (1)$$

- Low SFR: EW and $\frac{\text{H}\alpha}{\text{FUV}}$ ratios fluctuate [1], [2]



Stochasticity in star formation

Where does stochasticity arise?

- Star formation
- Initial Mass Function (IMF)
 - $m_{min} - m_{max}$
 - Relative abundance
- Cluster Mass Function (CMF)



Stochasticity in star formation

Where does stochasticity arise?

- Sampling of mass functions – SFR
- Low SFR:
 - Lowers probability of massive stars — finite sampling in mass [2]
 - Stellar mass and luminosity — highly nonlinear [3]
 - Total mass and luminosity — no longer deterministic [4]
 - Finite sampling in time — stellar phases [2]
 - $m_{max} < M_{ecl}$ [2]
 - Bursts of SFR [2]



Measuring stochasticity

How does stochasticity translate into observable quantities?

- SLUG: Stochastically Light Up Galaxies [5]
- EW, $\frac{H_\alpha}{FUV}$ fluctuates, [1][2]
- “We find that stochasticity alone induces a broad distribution in L_α and EW at a fixed SFR, and that the widths of these distributions decrease with increasing SFR” [1]



Measuring stochasticity

Double power law

$$P(\mathcal{M}|\text{SFR}) = P_0 \left[\left(\frac{\mathcal{M}}{\mathcal{M}_0} \right)^{-\alpha} + \left(\frac{\mathcal{M}}{\mathcal{M}_0} \right)^\gamma \right]^{-1} \quad (2)$$




Figure: SFR distributions, $\mathcal{M} = \frac{\text{EW}}{\text{EW}_0}$

Measuring stochasticity




Figure: Mas-Ribas et.al 2016 [6]



Motivation

General objective

“Look for observational evidence of stochasticity in star formation processes in the data published by CALIFA”.

Specific objectives

- Develop a simple theoretical model to measure the effects of stochasticity in the EW of the H_{α} and O_{II} emission lines
- Analyze data from the CALIFA survey collaboration
- Compare results between the observed data and the theoretical model
- Conclude if there is enough evidence to claim that stochastic effects have been detected in CALIFA data



Measuring stochasticity

- Balmer decrement
 - Interstellar dust
 - $\frac{H_\alpha}{H_\beta} = 2.85 \rightarrow \frac{H_\alpha}{H_\beta} \geq 2.85^1$
 - Interstellar reddening




Figure: Domínguez et. al. 2012 [7]

¹Osterbrock, Astrophysics of Planetary Nebulae and Active Galactic Nuclei, University Science Books, 1989



Table of Contents

1 Motivation

- Stochasticity in star formation
- Measuring stochasticity

2 The CALIFA survey

- CALIFA datacubes
- The Pipe3D pipeline

3 Results

4 Conclusions



The CALIFA survey

- ~ 600 galaxies
- “Largest and most comprehensive wide-field IFU survey of galaxies carried out to date”[8]




Figure: http://califaserv.caha.es/CALIFA/DATA/Figs/CALIFA_HexDR2.png



The CALIFA survey

- Calar Alto observatory
- Almería, Spain. 2010
- Many spectra per galaxy
- Evolution in mass, brightness of galaxies



The CALIFA datacubes

- 2 setups²
- V500
 - 3745 – 7500 Å [9]
 - $\lambda/\Delta\lambda \sim 850$
- V1200
 - 3700 – 4800 Å
 - $\lambda/\Delta\lambda \sim 1650$ [9]



Data analysis

- Fit spectral lines
 - Stellar population
 - Strong emission lines




Figure: Spectral fit [7]



Data analysis

- Pipe3D: analysis pipeline [10]
- *"The final product of the data reduction from both surveys is a regular grid datacube, with x and y coordinates that indicate the right ascension and declination of the target, and the z coordinate a common step in wavelength..."*



Table of Contents

1 Motivation

- Stochasticity in star formation
- Measuring stochasticity

2 The CALIFA survey

- CALIFA datacubes
- The Pipe3D pipeline

3 Results


4 Conclusions



Results





NGC0001
<http://www.galaxyzooforum.org/index.php?topic=280028.0>



NGC0036 <http://cseigman.com/text/atlas/ngc36.jpg>



Results



<http://califaserv.caha.es/CALIFA/DATA/V1200/v2.2/web/NGC0001> <http://califaserv.caha.es/CALIFA/DATA/V500/v2.2/web/NGC0036>



Results




Figure: NGC0001

Results




Figure: NGC0036



Results




Figure: NGC0001




Figure: NGC0036



Results

- H_{α} luminosity is a direct measure of SFR [11]
- Higher H_{α} luminosity → higher SFR [12]

$$\text{SFR}(\text{M}_{\odot}\text{yr}^{-1}) = 7.9 \times 10^{-42} L(H_{\alpha})(\text{ergs s}^{-1}) \quad (3)$$



Results




Figure: NGC0001




Figure: NGC0036



Results




Figure: NGC0001




Figure: NGC0036



Results




Figure: NGC0001




Figure: NGC0036

Results

- OII EW distribution




Figure: NGC0001




Figure: NGC0036



Table of Contents

1 Motivation

- Stochasticity in star formation
- Measuring stochasticity

2 The CALIFA survey

- CALIFA datacubes
- The Pipe3D pipeline

3 Results

4 Conclusions



Conclusions

- Fluctuation of $\frac{H_\alpha}{H_\beta}$, O_{II} EW was detected.
- Distribution of values follows the double power law (eq. 2)
- We propose stochasticity as a candidate that explains fluctuation of spectral parameters.
- **Further work:** Theoretical work on what is expected of distributions
- Quantify the impact of interstellar dust effects on our results
- Determine if stochasticity is observed
- Expand data sample (MaNGA [13])



References I

-  J. E. Forero-Romero and M. Dijkstra. "Effects of star-formation stochasticity on the Ly α and Lyman continuum emission from dwarf galaxies during reionization". In: 428 (Jan. 2013), pp. 2163–2170. DOI: 10.1093/mnras/sts177. arXiv: 1206.0726.
-  M. Fumagalli, R. L. da Silva, and M. R. Krumholz. "Stochastic Star Formation and a (Nearly) Uniform Stellar Initial Mass Function". In: 741, L26 (Nov. 2011), p. L26. DOI: 10.1088/2041-8205/741/2/L26. arXiv: 1105.6101.

References II



[JE Andrews et al.](#) "AN INITIAL MASS FUNCTION STUDY OF THE DWARF STARBURST GALAXY NGC 4214Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with program GO-11360." In: *The Astrophysical Journal* 767.1 (2013), p. 51.



[R. L. da Silva, M. Fumagalli, and M. R. Krumholz.](#) "SLUG - Stochastically Lighting Up Galaxies - II. Quantifying the effects of stochasticity on star formation rate indicators". In: 444 (Nov. 2014), pp. 3275–3287. DOI: [10.1093/mnras/stu1688](https://doi.org/10.1093/mnras/stu1688). arXiv: 1403.4605.



References III

-  Robert L. da Silva, Michele Fumagalli, and Mark Krumholz. "SLUG—Stochastically Lighting Up Galaxies. I. Methods and Validating Tests". In: *The Astrophysical Journal* 745.2 (2012), p. 145. URL:
<http://stacks.iop.org/0004-637X/745/i=2/a=145>.
-  L. Mas-Ribas, M. Dijkstra, and J. E. Forero-Romero. "Boosting Ly α and H β 1640Å Line Fluxes from Pop III Galaxies: Stochastic IMF Sampling and Departures from Case-B". In: *ArXiv e-prints* (Sept. 2016). arXiv: 1609.02150.
-  A. Domínguez et al. "Dust Extinction from Balmer Decrement of Star-forming Galaxies at $0.75 < z < 1.5$ with Hubble Space Telescope/Wide-Field-Camera 3 Spectroscopy from the WFC3 Infrared Spectroscopic Parallel Survey". In: *The Astrophysical Journal* 763.2 (2013), p. 145. URL:
<http://stacks.iop.org/0004-637X/763/i=2/a=145>.

References IV

-  S. F. Sánchez et al. “CALIFA, the Calar Alto Legacy Integral Field Area survey. I. Survey presentation”. In: 538, A8 (Feb. 2012), A8. DOI: 10.1051/0004-6361/201117353. arXiv: 1111.0962.
-  J. Sánchez Almeida et al. “Qualitative Interpretation of Galaxy Spectra”. In: 756, 163 (Sept. 2012), p. 163. DOI: 10.1088/0004-637X/756/2/163. arXiv: 1207.3928.
-  S. F. Sánchez et al. “Pipe3D, a pipeline to analyze Integral Field Spectroscopy Data: II. Analysis sequence and CALIFA dataproduccts”. In: 52 (Apr. 2016), pp. 171–220. arXiv: 1602.01830 [astro-ph.IM].
-  L. J. Kewley et al. “The H α and Infrared Star Formation Rates for the Nearby Field Galaxy Survey”. In: 124 (Dec. 2002), pp. 3135–3143. DOI: 10.1086/344487. eprint: astro-ph/0208508.



References V

-  R. C. Kennicutt Jr. "Star Formation in Galaxies Along the Hubble Sequence". In: 36 (1998), pp. 189–232. DOI: [10.1146/annurev.astro.36.1.189](https://doi.org/10.1146/annurev.astro.36.1.189). eprint: [astro-ph/9807187](https://arxiv.org/abs/astro-ph/9807187).
-  K. Bundy et al. "Overview of the SDSS-IV MaNGA Survey: Mapping nearby Galaxies at Apache Point Observatory". In: 798, 7 (Jan. 2015), p. 7. DOI: [10.1088/0004-637X/798/1/7](https://doi.org/10.1088/0004-637X/798/1/7). arXiv: [1412.1482](https://arxiv.org/abs/1412.1482).

