

Homework 4

Nikole Vannest, Emiliano Espinoza, Elena Ortiz

5/28/2021

1. Create a quadratic damage function relating the \$\$ value of damages to the change in global mean temperature and plot with underlying data.

```
#damages %>% ggplot(aes(x = warming, y = damages)) + geom_point()
```

Equation: $Y = 1.94x - 1.035x^2$

```
#y = ax^2 + bx + c
```

```
damages_quadratic <- cbind("warming_sq" = (damages$warming)^2, damages)
```

```
damages_lm <- lm(damages ~ warming_sq + warming + 0, data = damages_quadratic) #linear model
```

```
a <- damages_lm$coefficients[["warming_sq"]] #coefficient a
```

```
b <- damages_lm$coefficients[["warming"]] #coefficient b
```

```
damages_lm2 <- a*damages_quadratic$warming_sq + b*damages_quadratic$warming #Calculating quadratic equation
```

```
damages_quadratic %>% ggplot(aes(x = warming, y = damages)) +
```

```
  geom_point() +
```

```
  geom_line(aes(y = damages_lm2)) +
```


```
  ylab("Damage ($)") +
```

```
  xlab("Warming scenario (degrees C)") +
```

```
  theme_minimal()+
```

```
  ggtitle("Estimating Damages Under Warming Scenarios")
```

Estimating Damages Under Warming Scenarios




2. Use warming data to predict damages under baseline and pulse scenarios.

```
damages_baseline <- a*((warming$warming_baseline)^2) + b*warming$warming_baseline


warming %>% ggplot(aes(x = year, y = damages_baseline)) +
  geom_point() +
  geom_line(aes(y = damages_baseline)) +
  ylab("Damages ($)") +
  xlab("Year") +
  theme_minimal()+
  ggtitle("Estimating Damages Under Warming Baseline Scenario")
```

Estimating Damages Under Warming Baseline Scenario



```
damages_pulse <- a*((warming$warming_pulse)^2) + b*warming$warming_pulse


warming %>% ggplot(aes(x = year, y = damages_pulse)) +
  geom_point() +
  geom_line(aes(y = damages_pulse)) +
  ylab("Damages ($)") +
  xlab("Year") +
  theme_minimal()+
  ggtitle("Estimating Damages Under Pulse Warming Scenario")
```




```
difference_scenario <- damages_pulse - damages_baseline

warming %>% ggplot(aes(x = year, y = difference_scenario)) +
  geom_point() +
  geom_line(aes(y = difference_scenario)) +
  ylab("Damages ($)") +
  xlab("Year") +
  theme_minimal()+
  ggtitle("Estimating Differences in Damages Between Baseline and Pulse")
```

Estimating Differences in Damages Between Baseline and Pulse



```
per_ton <- difference_scenario / 35000000000 #to get per ton CO2  
  
warming %>% ggplot(aes(x = year, y = per_ton)) +  
  geom_point() +  
  geom_line(aes(y = per_ton)) +  
  ylab("Damages ($/ton CO^2)") +  
  xlab("Year") +  
  theme_minimal()+  
  ggtitle("Differences in Damages Between Baseline and Pulse in Per ton CO^2 Emitted")
```



3. Calculate the SCC against the discount rate for a reasonable range of discount rates.

- Obama: 3%
- NY State: 2%

Here we have calculated the SCC at discount rates between 1-5%


```
#cost/(1+rate)^year

SCC <- c()
rates <- c(0.01, 0.02, 0.03, 0.04, 0.05) #Discount rates chosen
for(i in 1:length(rates)) { #Calculating PV's in the loop
  temp <- (per_ton)/(1+rates[i])^(warming$X)
  SCC[i] <- sum(temp)
}

SCC_table <- round(data.frame("rates" = rates, #put into a data frame to run in ggplot
                             "SCC" = SCC), 2)

SCC_table %>% ggplot(aes(x = rates)) + #create a graph of SCC's under discount rates
  geom_point(aes(y = SCC)) +
  geom_line(aes(y = SCC)) +
  theme_minimal() +
  xlab("Discount Rate") +
  ylab("Social Cost of Carbon (SCC) in $/ton of CO^2") +
  ggtitle("SCC under Discount Rates 1-5%")
```

SCC under Discount Rates 1–5%



- SCC at 1%: \$113.64/ton CO²
- SCC at 2%: \$73.59/ton CO²
- SCC at 3%: \$50.02/ton CO²
- SCC at 4%: \$35.61/ton CO²
- SCC at 5%: \$26.45/ton CO²

4. What is the SCC using the Ramsey Rule?

```
#r = p + ng


p <- 0.001
n <- 2
g <- 0.01

ramsey <- p + n*g #Ramsey Rule equation

SCC_ramsey <- c()
rates_ramsey <- c(0.01, 0.02, 0.03, 0.04, 0.05, ramsey) #Adding Ramsey to the loop
for(i in 1:length(rates_ramsey)) {
  temp <- (per_ton)/(1+rates_ramsey[i])^(warming$X)
  SCC_ramsey[i] <- sum(temp)
}

ramsey_table <- data.frame("rates" = rates_ramsey, #Moving to a data frame for ggplot
                           "SCC" = SCC_ramsey)
```

```
ramsey_table %>% ggplot(aes(x = rates)) + #Graphing the SCC's under several discount rates
  geom_point(aes(y = SCC)) +
  geom_line(aes(y = SCC)) +
  theme_minimal() +
  ggtitle("SCC Discounted with 1-5% and the Ramsey Rule")
```



The point calculated using the Ramsey Rule is located at location (x = 0.021,y = 70.65).

5. What are the expected damages up to year 2100 under Policy A and Policy B?

Facts

- **Policy A:** Either baseline (probability 0.5) or warming each year will be 1.5 times that of the baseline (probability 0.5)
- **Policy B:** Continue until 2050 at baseline, then stay at 1.29 degrees warming forever
- Society is risk neutral
- Discount rate = 2%

```
#(Scenario 1 * probability) + (Scenario 2 * probability)
warming_1.5 <- warming$warming_baseline * 1.5 #create warming scenario
damages_1.5 <- a*((warming_1.5)^2) + b*warming_1.5 #plug into quadratic equation

policy_a <- (damages_baseline*0.5) + (damages_1.5*0.5) #calculate overall expected damages

policy_a_discount <- (policy_a)/(1+0.02)^(warming$X) #calculate present value for each year
policy_a_pv <- sum(policy_a_discount) #Sum for NPV
```



```

b_1.29 <- rep(1.29, 50) #Create warming sequence for scenario b
policy_b_warming <- c(warming$warming_baseline[1:30], b_1.29) #combine with baseline
damages_b <- a*((policy_b_warming)^2) + b*policy_b_warming #put into quadratic equation
policy_b <- (damages_b)/(1+0.02)^(warming$X) #calculate present value
policy_b_pv <- sum(policy_b) #find NPV

difference_ab <- policy_a_pv-policy_b_pv #Value of X to meet damages in policy A

baseline_0.02 <- (damages_baseline)/(1+0.02)^(warming$X)
baseline_sum <- sum(baseline_0.02) #Calculate sum of PV of baseline scenario alone

warm_1.5_0.02 <- (damages_1.5)/(1+0.02)^(warming$X) #Calculate sum of PV of 1.5 warming scenario alone
warm1.5_sum <- sum(warm_1.5_0.02)

```

- Expected damages under Policy A: \$3.0340e+15
- Expected damages under Policy B: \$7.7825e+14
- Initial undertaking of Policy B would have to cost less than \$2.2558e+15 to justify choosing Policy A over Policy B
- If society was risk averse and the expected values for both policies were the same, Policy B would still be the preferred outcome because the costs are guaranteed. On the other hand, Policy A is a riskier option because the expected value is not guaranteed (there are two scenarios).