

Initial Analysis

Kyle Ligon

Read the data in and library calls

```
library(tidyverse)
library(zoo)
library(lubridate)
library(gridExtra)

crime <- read_csv("BPD_Part_1_Victim_Based_Crime_Data.csv", progress = FALSE)
```

Looking at the data

```
head(crime)

## # A tibble: 6 x 15
##   CrimeDate CrimeTime CrimeCode Location      Description `Inside/Outside`
##   <chr>      <time>      <chr>      <chr>      <chr>          <chr>
## 1 9/2/2017  23:30      3JK       4200 AUDRE~ ROBBERY - RE~ I
## 2 9/2/2017  23:00      7A        800 NEWING~ AUTO THEFT  0
## 3 9/2/2017  22:53      9S        600 RADNOR~ SHOOTING    Outside
## 4 9/2/2017  22:50      4C        1800 RAMSA~ AGG. ASSAULT I
## 5 9/2/2017  22:31      4E        100 LIGHT ~ COMMON ASSAU~ 0
## 6 9/2/2017  22:00      5A        CHERRYCRE~ BURGLARY    I
## # ... with 9 more variables: Weapon <chr>, Post <int>, District <chr>,
## #   Neighborhood <chr>, Longitude <dbl>, Latitude <dbl>, `Location
## #   1` <chr>, Premise <chr>, `Total Incidents` <int>

names(crime)

## [1] "CrimeDate"      "CrimeTime"      "CrimeCode"
## [4] "Location"       "Description"     "Inside/Outside"
## [7] "Weapon"         "Post"           "District"
## [10] "Neighborhood"   "Longitude"       "Latitude"
## [13] "Location 1"     "Premise"         "Total Incidents"
```

Looks like we have information about the crime, where it happened, when it happened, what happened in the form of Description, and the responding Post.

Counting up the Number of Crimes

```
descCounts <- crime %>%
  group_by(Description) %>%
  tally() %>%
  arrange(desc(n))

descCounts

## # A tibble: 15 x 2
##   Description      n
##   <chr>          <int>
```

```
## 1 LARCENY 60528
## 2 COMMON ASSAULT 45518
## 3 BURGLARY 42538
## 4 LARCENY FROM AUTO 36295
## 5 AGG. ASSAULT 27513
## 6 AUTO THEFT 26838
## 7 ROBBERY - STREET 17691
## 8 ROBBERY - COMMERCIAL 4141
## 9 ASSAULT BY THREAT 3503
## 10 SHOOTING 2910
## 11 ROBBERY - RESIDENCE 2866
## 12 RAPE 1637
## 13 HOMICIDE 1559
## 14 ROBBERY - CARJACKING 1528
## 15 ARSON 1464
```

Counting up Where the Crimes Occurred

```
hoodCounts <- crime %>%
  group_by(Neighborhood) %>%
  tally() %>%
  arrange(desc(n))
head(hoodCounts)
```

```
## # A tibble: 6 x 2
##   Neighborhood      n
##   <chr>          <int>
## 1 Downtown      9048
## 2 Frankford     6642
## 3 Belair-Edison 5977
## 4 Brooklyn     4516
## 5 Cherry Hill   4086
## 6 Sandtown-Winchester 4026
```

Let's focus on Arsons. Particularly, let's see if the number of Arsons committed in one month are more varied in the winter months than in the other nine months of the year. For this I will:

- Summarize the number of arsons by month
- Run an F test on number of arsons between the two groups
- Write a conclusion for the test

1) Are the distributions of arsons in the winter months less varied than other 9 months?

```
arsons <- crime %>%
  filter(Description == "ARSON")
head(arsons)
```


```
## # A tibble: 6 x 15
##   CrimeDate CrimeTime CrimeCode Location Description `Inside/Outside`
##   <chr>      <time>    <chr>    <chr>      <chr>      <chr>
## 1 9/1/2017  22:00      8AO      300 N FREMON~ ARSON      I
## 2 8/30/2017 22:00      8H      2600 FLORA ST ARSON      <NA>
```

```
## 3 8/30/2017 19:30      8H      3700 CLIFTMO~ ARSON      0
## 4 8/30/2017 15:26      8AV     4600 PARK HE~ ARSON      I
## 5 8/29/2017 03:30      8H      800 RAPPOLLA~ ARSON     <NA>
## 6 8/28/2017 06:50      8H      3300 TIVOLY ~ ARSON      0
## # ... with 9 more variables: Weapon <chr>, Post <int>, District <chr>,
## #   Neighborhood <chr>, Longitude <dbl>, Latitude <dbl>, `Location
## #   1` <chr>, Premise <chr>, `Total Incidents` <int>
```

Checking the dimensions of the new frame

```
dim(arsons)
```

```
## [1] 1464   15
```



With “normal” distributions, we will proceed with the hypotheses.

Hypotheses: $H_0: \text{var_Winter} = \text{var_SprSumFall}$ $H_1: \text{var_Winter} < \text{var_SprSumFall}$

Variance of the Winter Months

```
#degrees of freedom for W
arsons_grouped %>% filter(WinterBin == "Winter") %>% tally() - 1
```

```
##      n
## 1 16
```

```
var_W
```

```
## # A tibble: 1 x 1
##   Variance
##     <dbl>
```

```
## 1      27.3
```

Variance of the Other Months

```
arsons_grouped %>% filter(WinterBin != "Winter") %>% tally() - 1
```

```
##      n
```

```
## 1 51
```

```
var_SSF
```

```
## # A tibble: 1 x 1
```

```
##   Variance
```

```
##   <dbl>
```

```
## 1      45.9
```

Test Statistic for F Test

```
f <- as.numeric(round(var_W/var_SSF, 4))
```

```
f
```

```
## [1] 0.5945
```

Rejection Region

```
qf(0.975, 16, 51)
```

```
## [1] 2.075301
```

```
f < qf(0.025, 16, 51)
```

```
## [1] FALSE
```

```
f > qf(0.975, 16, 51)
```

```
## [1] FALSE
```

Since our F Test Statistic is not larger and not smaller than the F Stat for alpha, we do not have enough evidence to reject the null hypothesis that the variances are equal. It does not appear that the the Winter months experience a less varied number of arsons than the other 9 months.

2) Are there more Auto Thefts on Weekends over Weekdays?

```
auto_thefts <- crime %>%  
  filter(Description %in% c("ROBBERY - CARJACKING", "AUTO THEFT"))  
head(auto_thefts)
```

```
## # A tibble: 6 x 15
```

```
##   CrimeDate CrimeTime CrimeCode Location      Description `Inside/Outside`
```

```
##   <chr>      <time>    <chr>    <chr>      <chr>      <chr>
```

```
## 1 9/2/2017  23:00      7A      800 NEWINGTO~ AUTO THEFT  0
```

```
## 2 9/2/2017  08:00      7A      4700 HOMESDA~ AUTO THEFT  I
```

```
## 3 9/2/2017  02:00      7C      1500 RUSSELL~ AUTO THEFT  0
```

```
## 4 9/1/2017  22:30      7A      300 E LORRAI~ AUTO THEFT  0
```

```
## 5 9/1/2017  21:30      7A      3500 CHESTER~ AUTO THEFT  0
```

```
## 6 9/1/2017  20:45      7A      OSTEND ST & ~ AUTO THEFT  0
```

```
## # ... with 9 more variables: Weapon <chr>, Post <int>, District <chr>,
```

```
## #   Neighborhood <chr>, Longitude <dbl>, Latitude <dbl>, `Location
```

```
## #   1` <chr>, Premise <chr>, `Total Incidents` <int>
```

```


at_form <- auto_thefts %>%
  mutate(CrimeDate= as.Date(CrimeDate, format = "%m/%d/%Y"),
         DateName = wday(CrimeDate, label = TRUE))
weekend <- at_form %>%
  filter(DateName %in% c("Sat", "Sun")) %>%
  group_by(CrimeDate, DateName) %>%
  summarize(Count = sum(`Total Incidents`))

weekday <- at_form %>%
  filter(!(DateName %in% c("Sat", "Sun"))) %>%
  group_by(CrimeDate, DateName) %>%
  summarize(Count = sum(`Total Incidents`))

ggplot(data = weekday, aes(x = Count)) + geom_histogram() + ggtitle("Weekday Distribution")

```

Weekday Distribution




```

ggplot(data = weekend, aes(x = Count)) + geom_histogram() + ggtitle("Weekend Distribution")

```

Weekend Distribution



With our data cleaned, we can now go about testing to see if the mean of the weekend set is larger than the mean of the weekday set. But first... equal variance check. Which, is just an F test:

H_0: Var_Weekday = var_Weekend H_1: var_Weekday < var_weekend

```
#Weekday Variance
var(weekday$Count)
```

```
## [1] 28.16756
```

```
#df of Weekday
nrow(weekday)-1
```

```
## [1] 1479
```

```
#Weekend Variance
var(weekend$Count)
```

```
## [1] 30.29062
```

```
#df of Weekends
nrow(weekend)-1
```

```
## [1] 591
```

Test Statistic

```
f_w <- var(weekday$Count)/var(weekend$Count)
f_w
```

```
## [1] 0.9299104
```

Rejection Region:

```
up <- qf(0.975, 1479, 591)
up
```

```
## [1] 1.146795
```

```
dwn <- qf(0.025, 1479, 591)
dwn
```

```
## [1] 0.8754564
```

```
f_w > up
```

```
## [1] FALSE
```

```
f_w < dwn
```

```
## [1] FALSE
```

Conclusion: Since our F Stat was less than the upper rejection region and more than the lower rejection region, we do not have enough evidence to reject the hypothesis that the variances are equal. It does not appear that the variances are different. Now we can test the means.

Hypotheses: H_0 : mean_weekday = mean_weekend H_1 : mean_weekday < mean_weekend

Test Statistic:

```
t.test(x = weekday$Count, y = weekend$Count)
```

```
##
```

```
## Welch Two Sample t-test
```

```
##
```

```
## data: weekday$Count and weekend$Count
```

```
## t = -1.5301, df = 1054.2, p-value = 0.1263
```

```
## alternative hypothesis: true difference in means is not equal to 0
```

```
## 95 percent confidence interval:
```

```
## -0.9252956 0.1144848
```

```
## sample estimates:
```

```
## mean of x mean of y
```

```
## 13.57432 13.97973
```

Rejection Region

```
lower <- qt(0.05, 2070)
lower
```

```
## [1] -1.64559
```

```
-1.5301 < lower
```

```
## [1] FALSE
```

Conclusion: Since -1.5301 is greater than our rejection region, we do not have enough proof to reject our null hypothesis that the means are the same. It does not appear that there are more Auto thefts on weekends in comparison to weekdays.

3) On a given night are there more shootings inside a residence than outside?

```
shootings <- crime %>%
# mutate(CrimeTime = strptime(CrimeTime, format = "%H:%M")) %>%
```

```

filter(Description == "SHOOTING", CrimeTime > "17:00") %>%
group_by(CrimeDate, `Inside/Outside`) %>%
summarize(Count = sum(`Total Incidents`))
head(shootings)

```

```


## # A tibble: 6 x 3
## # Groups:   CrimeDate [5]
##   CrimeDate `Inside/Outside` Count
##   <chr>      <chr>          <int>
## 1 1/1/2012 Outside            1
## 2 1/1/2013 Inside              1
## 3 1/1/2013 Outside            1
## 4 1/1/2014 Outside            2
## 5 1/1/2015 Outside            1
## 6 1/1/2016 Outside            3

```

```
dim(shootings)
```

```
## [1] 1493    3
```

```
ggplot(shootings, aes(x = Count)) + geom_histogram() + facet_grid(~ `Inside/Outside`)
```



With non-normal distributions, our route that we should run down is to check to see if the medians are different between the two groups. Using the Wilcoxon Rank Sum test, we'll see what if the outside median is larger than the inside median.

Hypothesis: $H_0: M_{\text{Inside}} = M_{\text{Outside}}$ $H_1: M_{\text{Inside}} > M_{\text{Outside}}$


```
w <- wilcox.test(Count ~ `Inside/Outside`, data = shootings, alternative = "greater")
```

Test Statistic

```
w$statistic
```

```
##      W
```

```
## 98797
```

Rejection Region

```
## [1] FALSE
```

Conclusion/Interpretation: Since our p-value is greater than 0.05, we do not have enough evidence to reject the null hypothesis that the Medians are the same. It does not appear there are more shootings inside a residence than outside.

4) On a given day if you are assaulted, is it more likely to be a common assault?


Hypotheses: H_0 : Mean_aggAssault = Mean_CommonAssault H_1 : Mean_aggAssault < Mean_CommonAssault

```
assaults <- crime %>%
  filter(Description %in% c("AGG. ASSAULT", "COMMON ASSAULT")) %>%
  select(CrimeDate, Description, `Total Incidents`) %>%
  group_by(CrimeDate, Description) %>%
  summarize(Total = sum(`Total Incidents`)) %>%
  ungroup()

day_merge <- assaults %>%
  group_by(CrimeDate) %>%
  summarize(Count = sum(Total))

assaults <- assaults %>%
  group_by(CrimeDate) %>%
  left_join(day_merge) %>%
  mutate(percent = Total/Count) %>%
  ungroup() %>%
  mutate(CrimeDate = as.Date(CrimeDate, format = "%m/%d/%Y")) %>%
  arrange(desc(CrimeDate))

ggplot(data = assaults, aes(x = percent)) + geom_histogram() + facet_grid(~Description)
```



```
assaults_split <- t.test(formula = percent ~ Description, data = assaults, alternative = "less")
```

Test Statistic/p-value

```
assaults_split$statistic
```

```
##          t
## -80.19273
```

```
assaults_split$p.value
```

```
## [1] 0
```

Conclusion: Since our p-value is less than 0.05, we have sufficient evidence to reject the null hypothesis that the means are the same. It appears that the Mean percent of the Aggravated Assault group is less than the Mean percent of the Common Assault.

5) Is there a District with a distinctly higher number of Burglary's in the Summer Months?
If so, which are significantly different?

H₀: means are all the same H₁: at least one mean is different


```
summer_burglaries <- crime %>%
  filter(substr(CrimeDate, 1,1) %in% c('6','7','8'), Description == "BURGLARY") %>%
  mutate(CrimeDate <- as.Date(CrimeDate, format = "%m/%d/%Y")) %>%
  mutate(YearMon = as.yearmon(CrimeDate, "%m/%d/%Y")) %>%
  group_by(District, YearMon) %>%
  summarize(Count = sum(`Total Incidents`)) %>%
```

```

    filter(District != "NA") %>%
    ungroup()

ggplot(data= summer_burglaries, aes(x = Count)) + geom_histogram() + facet_wrap(~ District)

```



Test Choice: Since there is a mess of distributions that appear to fit an ANOVA model, I'll use the ANOVA test to see if there's one mean that's different.

```

summer_burglaries <- summer_burglaries %>%
  mutate(District = as.factor(District))

sum_burg_anova_check <- aov(Count ~ District, data = summer_burglaries)

y <- quantile(sum_burg_anova_check$residuals[!is.na(sum_burg_anova_check$residuals)], c(0.25, 0.75))
x <- qnorm(c(0.25, 0.75))
slope = diff(y)/diff(x)
int <- y[1L] - slope*x[1L]


norm_res <- ggplot(data = sum_burg_anova_check, aes(x = sum_burg_anova_check$residuals)) + geom_histogram()

resid_jitter <- ggplot(data = sum_burg_anova_check, aes(x = sum_burg_anova_check$fitted.values, y = sum

```

```
qq <- ggplot(data = sum_burg_anova_check) + stat_qq(aes(sample = sum_burg_anova_check$residuals), alpha = 0.5) +
  geom_abline(slope = slope, intercept = int, color = "blue") + ggtitle("Q-Q Plot")

res_sum <- as.table(summary(sum_burg_anova_check$residuals), nrow = 2, ncol = 6)
grid.arrange(resid_jitter, norm_res, qq, nrow = 2, ncol = 2)
```



Checking ANOVA/Test Statistic/p-value

```
summary(sum_burg_anova_check)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## District      8  88078   11010   32.54 <2e-16 ***
## Residuals    153  51772     338
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Since our p-value is considerably less than 0.05, we can conclude that it appears that one mean is different than the other. We will proceed with Tukey's HSD to show which district is different from another.

```
thsd_sum_burg <- TukeyHSD(sum_burg_anova_check)
thsd_sum_burg
```

```
##      Tukey multiple comparisons of means
##      95% family-wise confidence level
##
## Fit: aov(formula = Count ~ District, data = summer_burglaries)
##
```

## \$District		diff	lwr	upr	p adj
## EASTERN-CENTRAL	-3.055556	-22.3503538	16.239243	0.9998975	
## NORTHEASTERN-CENTRAL	68.222222	48.9274240	87.517020	0.0000000	
## NORTHERN-CENTRAL	43.000000	23.7052018	62.294798	0.0000000	
## NORTHWESTERN-CENTRAL	18.722222	-0.5725760	38.017020	0.0647509	
## SOUTHEASTERN-CENTRAL	33.944444	14.6496462	53.239243	0.0000046	
## SOUTHERN-CENTRAL	37.611111	18.3163129	56.905909	0.0000003	
## SOUTHWESTERN-CENTRAL	17.055556	-2.2392427	36.350354	0.1295154	
## WESTERN-CENTRAL	-6.222222	-25.5170205	13.072576	0.9839710	
## NORTHEASTERN-EASTERN	71.277778	51.9829795	90.572576	0.0000000	
## NORTHERN-EASTERN	46.055556	26.7607573	65.350354	0.0000000	
## NORTHWESTERN-EASTERN	21.777778	2.4829795	41.072576	0.0145920	
## SOUTHEASTERN-EASTERN	37.000000	17.7052018	56.294798	0.0000004	
## SOUTHERN-EASTERN	40.666667	21.3718684	59.961465	0.0000000	
## SOUTHWESTERN-EASTERN	20.111111	0.8163129	39.405909	0.0340026	
## WESTERN-EASTERN	-3.166667	-22.4614649	16.128132	0.9998657	
## NORTHERN-NORTHEASTERN	-25.222222	-44.5170205	-5.927424	0.0020286	
## NORTHWESTERN-NORTHEASTERN	-49.500000	-68.7947982	-30.205202	0.0000000	
## SOUTHEASTERN-NORTHEASTERN	-34.277778	-53.5725760	-14.982980	0.0000036	
## SOUTHERN-NORTHEASTERN	-30.611111	-49.9059094	-11.316313	0.0000556	
## SOUTHWESTERN-NORTHEASTERN	-51.166667	-70.4614649	-31.871868	0.0000000	
## WESTERN-NORTHEASTERN	-74.444444	-93.7392427	-55.149646	0.0000000	
## NORTHWESTERN-NORTHERN	-24.277778	-43.5725760	-4.982980	0.0035837	
## SOUTHEASTERN-NORTHERN	-9.055556	-28.3503538	10.239243	0.8645865	
## SOUTHERN-NORTHERN	-5.388889	-24.6836871	13.905909	0.9937698	
## SOUTHWESTERN-NORTHERN	-25.944444	-45.2392427	-6.649646	0.0012955	
## WESTERN-NORTHERN	-49.222222	-68.5170205	-29.927424	0.0000000	
## SOUTHEASTERN-NORTHWESTERN	15.222222	-4.0725760	34.517020	0.2488070	
## SOUTHERN-NORTHWESTERN	18.888889	-0.4059094	38.183687	0.0601195	
## SOUTHWESTERN-NORTHWESTERN	-1.666667	-20.9614649	17.628132	0.9999991	
## WESTERN-NORTHWESTERN	-24.944444	-44.2392427	-5.649646	0.0024032	
## SOUTHERN-SOUTHEASTERN	3.666667	-15.6281316	22.961465	0.9995979	
## SOUTHWESTERN-SOUTHEASTERN	-16.888889	-36.1836871	2.405909	0.1381106	
## WESTERN-SOUTHEASTERN	-40.166667	-59.4614649	-20.871868	0.0000000	
## SOUTHWESTERN-SOUTHERN	-20.555556	-39.8503538	-1.260757	0.0273407	
## WESTERN-SOUTHERN	-43.833333	-63.1281316	-24.538535	0.0000000	
## WESTERN-SOUTHWESTERN	-23.277778	-42.5725760	-3.982980	0.0064007	

- The Northeastern District has distinctly larger number of burglaries than all other groups.
- The Northwestern District has a distinctly larger number of burglaries than the Western, Northern, and Eastern.
- The Southern District has a distinctly larger number of burglaries in the Summer months over the Central, Eastern, and the Southwest.
- The Southwest District has a distinctly larger number of burglaries in the Summer months over the Eastern, Northern, and the Western districts.
- The Western District has a distinctly larger number of burglaries in the Summer months over the Northern and southern Districts.
- The Northern District has a distinctly larger number of burglaries in the Summer months over the Central and Eastern Districts.
- The Southeastern District has a distinctly larger number of burglaries in the Summer months over the

Central and Eastern Districts.