

# Hyperbolic PDEs and GENERIC

Nikolas Siccha

August 7, 2020

## Abstract

# Acknowledgements

# Contents

<b>Abstract</b>	<b>i</b>
<b>Acknowledgements</b>	<b>i</b>
<b>Table of contents</b>	<b>i</b>
<b>List of Figures</b>	<b>v</b>
<b>List of Tables</b>	<b>vi</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Outline . . . . .	2
<b>2 Finite and infinite dimensional geometry</b>	<b>3</b>
2.1 Differential geometry . . . . .	4
2.1.1 Manifolds . . . . .	4
2.1.2 Tensor fields . . . . .	4
2.1.3 The Lie derivative . . . . .	4
2.1.4 Metrics . . . . .	4
2.1.5 Connections . . . . .	4
2.1.6 Covariant derivatives and geodesics . . . . .	4
2.1.7 Curvature . . . . .	4
2.1.8 Torsion . . . . .	4
2.1.9 The Levi-Civita connection . . . . .	4
2.2 Poisson structures . . . . .	7
2.2.1 Poisson manifolds . . . . .	7
2.2.2 The Poisson bivector . . . . .	7
2.2.3 The Schouten bracket . . . . .	7
2.2.4 Constant Poisson structures and symplectic structures . .	7
2.2.5 Symplectic foliations . . . . .	7
2.2.6 Poisson structure reductions . . . . .	7
2.3 Lie groups and Lie algebras . . . . .	8
2.3.1 Lie-Poisson structures . . . . .	8
2.3.2 Extensions of Lie algebras . . . . .	8
2.3.3 Groups of diffeomorphisms . . . . .	8

2.3.4	Semidirect products . . . . .	8
<b>3</b>	<b>The geometry of hyperbolic PDEs of metriplectic type</b>	<b>9</b>
3.1	GENERIC / metriplectic dynamics . . . . .	10
3.1.1	Gradient flows . . . . .	10
3.1.2	Symplectic flows . . . . .	10
3.1.3	Reversibility . . . . .	10
3.1.4	Irreversibility . . . . .	10
3.1.5	Rational extended thermodynamics . . . . .	10
3.2	Hyperbolic partial differential equations . . . . .	11
3.2.1	Symmetric hyperbolic thermodynamically compatible systems . . . . .	15
3.2.2	Skew-selfadjoint hyperbolic systems of balance laws . . . . .	15
3.3	Poisson brackets of hydrodynamic type . . . . .	16
3.3.1	with nondegenerate metric . . . . .	18
3.3.1.1	in one spatial dimension . . . . .	18
3.3.1.2	in multiple spatial dimensions . . . . .	18
3.3.2	with degenerate metric . . . . .	20
3.3.2.1	in one spatial dimension . . . . .	20
3.3.2.2	in multiple spatial dimensions . . . . .	22
3.3.3	of Lie-Poisson type . . . . .	22
3.3.3.1	in one spatial dimension . . . . .	22
3.3.3.2	in multiple spatial dimensions . . . . .	22
3.4	Hyperbolic Poisson brackets . . . . .	24
3.4.1	of constant type . . . . .	24
3.4.2	with non-degenerate metric . . . . .	27
3.4.2.1	in one spatial dimension . . . . .	27
3.4.2.2	in multiple spatial dimensions . . . . .	33
3.4.3	with degenerate metric . . . . .	33
3.4.3.1	in one spatial dimension . . . . .	33
3.4.3.2	in multiple spatial dimensions . . . . .	33
3.4.4	of Lie-Poisson type . . . . .	33
3.4.4.1	in one spatial dimension . . . . .	33
3.4.4.2	in multiple spatial dimensions . . . . .	33
3.5	Hyperbolic gradient flows . . . . .	34
<b>4</b>	<b>Structure preserving integrators</b>	<b>35</b>
4.1	Symplectic integrators . . . . .	36
4.2	Energy preserving integrators . . . . .	37
4.3	Poisson integrators . . . . .	38
4.4	Multi-symplectic integrators . . . . .	39
4.5	Dissipation preserving integrators . . . . .	40
4.6	Conformal symplectic integrators . . . . .	41
4.7	Metriplectic integrators . . . . .	42
4.8	Finite volume methods . . . . .	43
4.9	Symmetric integrators . . . . .	44

4.9.1	The discontinuous Galerkin time stepping method . . . .	44
4.9.2	The adjoint of the discontinuous Galerkin time stepping method . . . . .	44
4.9.3	The symmetric discontinuous Galerkin time stepping method	44
4.9.4	The symmetric-dissipative discontinuous Galerkin time step- ping method . . . . .	44
<b>5</b>	<b>Conclusion</b>	<b>45</b>

# List of Figures

3.1	Constructions for PROPOSITION . . . . .	23
3.2	"Excess hyperbolicity" for a counterexample to [PPK20], Theo- rem 2.10 . . . . .	29

# List of Tables



# Chapter 1

## Introduction

## 1.1 Outline

## Chapter 2

# Finite and infinite dimensional geometry

## 2.1 Differential geometry

### 2.1.1 Manifolds

### 2.1.2 Tensor fields

### 2.1.3 The Lie derivative

### 2.1.4 Metrics

### 2.1.5 Connections

### 2.1.6 Covariant derivatives and geodesics

### 2.1.7 Curvature

### 2.1.8 Torsion

### 2.1.9 The Levi-Civita connection

For now, we will use this as a dump for formulae and definitions:

*Christoffel symbols of the second kind*

$$\Gamma_{i,j}^k = \frac{1}{2}g^{k,l}(\partial_i g_{j,l} + \partial_j g_{i,l} - \partial_l g_{i,j}) \quad (2.1)$$

*Riemannian curvature endomorphism*

$$R_{i,j,k}^l = [\partial_j \Gamma_{i,k}^l + \Gamma_{i,k}^m \Gamma_{j,m}^l]_{[i,j]} \quad (2.2)$$

*Riemannian curvature tensor*

$$Rm_{i,j,k,l} = R_{i,j,k}^n g_{l,n} \quad (2.3)$$

*flatness criterion*

$$Rm_{i,j,k,l} = 0 \quad (2.4)$$

*transformed metric*

$$g_{i,\hat{j}} = u_i^i g_{i,j} u_j^j \quad (2.5)$$

*transformed hessian*

$$h_{i,\hat{j}} = u_i^i h_{i,j} u_j^j + h_i u_{i,\hat{j}}^i \quad (2.6)$$

Following definitions just a test for back referencing.

**Definition 2.1.1.** A metric  $g(u)$  is called Riemannian if it is everywhere positive definite.

**Definition 2.1.2.** A metric  $g(u)$  is called pseudo-Riemannian if it is everywhere non-degenerate.

All following definitions collected from [Lee12].

**Definition 2.1.3.** Consider a continuous map  $\pi : \mathcal{M}^d \rightarrow N$ . A section of  $\pi$  is a continuous right inverse for  $\pi$ , i.e. a continuous map  $\sigma : N \rightarrow \mathcal{M}^d$  such that  $\pi \circ \sigma = \text{Id}_N$ . A local section need only be defined on some open subset  $U \subseteq \mathcal{M}^d$ .

**Definition 2.1.4.** Let  $\mathcal{M}^d$  be a topological space. A (real) vector bundle of rank  $k$  over  $\mathcal{M}^d$  is a topological space  $E$  together with a surjective continuous map  $\pi : E \rightarrow \mathcal{M}^d$  satisfying the following conditions:

- For each point  $p \in \mathcal{M}^d$ , the fiber  $E_p = \pi^{-1}(p)$  over  $p$  is endowed with the structure of a  $k$ -dimensional real vector space.
- For each point  $p \in \mathcal{M}^d$ , there exist a neighborhood  $U$  of  $p$  in  $\mathcal{M}^d$  and a homeomorphism  $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$  satisfying
  - $\pi_U \circ \Phi = \pi$  and
  - for each  $q \in U$ , the restriction of  $\Phi$  to  $E_q$  is a vector space isomorphism from  $E_q$  to  $\{q\} \times \mathbb{R}^k$ .

**Definition 2.1.5.** Consider a vector bundle  $\pi : E \rightarrow \mathcal{M}^d$ . A section of  $E$  is a section of the map  $\pi$ . A local section need only be defined on some open subset  $U \subseteq \mathcal{M}^d$ , while a global section is defined on all of  $\mathcal{M}^d$ .

**Definition 2.1.6.** The Lie bracket of two smooth vector fields  $X$  and  $Y$  is defined by

$$[X, Y](f) = X(Y(f)) - Y(X(f)) \quad (2.7)$$

for every smooth function  $f \in C^\infty(\mathcal{M}^d)$ .

**Definition 2.1.7.** A smooth distribution  $D$  on a smooth manifold  $\mathcal{M}^d$  of rank  $k$  is a rank- $k$  smooth subbundle of the tangent bundle  $T\mathcal{M}^d$ .

**Definition 2.1.8.** A smooth distribution  $D$  on  $\mathcal{M}^d$  is said to be involutive if given any pair of smooth local sections of  $D$ , their Lie bracket is also a local section of  $D$ .

**Definition 2.1.9.** A nonempty immersed submanifold  $N \subseteq \mathcal{M}^d$  is called an integral manifold of  $D$  if  $T_p N = D_p$  at each point  $p \in N$ .

**Definition 2.1.10.** A smooth distribution  $D$  on  $\mathcal{M}^d$  is said to be integrable if each point of  $\mathcal{M}^d$  is contained in an integral manifold of  $D$ .

**Definition 2.1.11.** Consider a smooth manifold  $\mathcal{M}^d$  and let  $\mathcal{F}$  be any collection of  $k$ -dimensional submanifolds of  $M$ . A smooth chart  $(U, \phi)$  is said to be flat for  $\mathcal{F}$  if  $\phi(U)$  is a cube in  $\mathbb{R}^d$  and each submanifold in  $\mathcal{F}$  intersects  $U$  in either the empty set or a countable union of  $k$ -dimensional slices of the form  $x^i = c^i$  for  $i = k+1, \dots, n$ .

**Definition 2.1.12.** A smooth distribution  $D$  is said to be completely integrable if there exists a flat chart for  $D$  in a neighborhood of each point  $p \in \mathcal{M}^d$ .

**Proposition 2.1.13.** *Every involutive distribution is completely integrable and vice versa.*

*Proof.* See e.g. [Lee12], theorem 19.12. □

**Definition 2.1.14.** *A collection  $\mathcal{F}$  of disjoint, connected, nonempty, immersed  $k$ -dimensional submanifolds of  $\mathcal{M}^d$ , whose union is  $\mathcal{M}^d$ , such that in a neighborhood of each point  $p \in \mathcal{M}^d$  there exists a flat chart for  $\mathcal{F}$  is called foliation of dimension  $k$  on  $\mathcal{M}^d$ .*

## 2.2 Poisson structures

### 2.2.1 Poisson manifolds

### 2.2.2 The Poisson bivector

### 2.2.3 The Schouten bracket

### 2.2.4 Constant Poisson structures and symplectic structures

### 2.2.5 Symplectic foliations

### 2.2.6 Poisson structure reductions

## 2.3 Lie groups and Lie algebras

### 2.3.1 Lie-Poisson structures

### 2.3.2 Extensions of Lie algebras

### 2.3.3 Groups of diffeomorphisms

### 2.3.4 Semidirect products



## Chapter 3

# The geometry of hyperbolic PDEs of metriplectic type

## 3.1 GENERIC / metriplectic dynamics

The main references are

- [GO97], [OG97] “First” papers
- [DV80] “First” first mention
- [Arn66] French hydrodynamic Lie groups
- [GMR96] SHTC
- [MR98] RET Book
- [Ott05] “Standard” reference
- [PKG14] “Sensible” reversibility
- [PKG18] Better “standard” reference
- [PPRG18] GENERIC  $\leftrightarrow$  SHTC
- [PPK20] Hamiltonian continuum mechanics
- [OST20] XGENERIC

### 3.1.1 Gradient flows

Give motivation, examples and properties for (physical) gradient flows.

### 3.1.2 Symplectic flows

Give motivation, examples and properties for (physical) symplectic flows.

### 3.1.3 Reversibility

Discuss parity, time-reversal transformations etc. See e.g. [PKG14].

### 3.1.4 Irreversibility

Discuss two options:

- Phase-space area shrinkage
- (Free) energy decrease

### 3.1.5 Rational extended thermodynamics

Give short discussion of RET.

## 3.2 Hyperbolic partial differential equations

The main references are

- [BGS06] Standard hyp. PDE Book
- [God62] Ex. solutions
- [Rug] Ex. solutions
- [FL71] Convex entropy extension
- [Tad84] Skew-selfadjoint hyp. PDEs
- [PPRG18] GENERIC  $\leftrightarrow$  SHTC
- [PPK20] Hamiltonian continuum mechanics

Throughout, unless otherwise stated, we will use a  $d$ -dimensional spatial manifold  $\mathcal{M}^d$ , a simply connected  $n$ -dimensional target manifold  $\mathcal{U}^n$  and assume  $u(t) \in C^\infty(\mathcal{M}^d; \mathcal{U}^n)$  for all  $t$  considered. Furthermore, indices in products which appear once above and once below will be summed. Greek indices like  $\mu, \nu$  run from 1 to  $d$  and latin indices like  $i, j, k$  run from 1 to  $n$ , unless otherwise noted.

**Definition 3.2.1.** *We consider systems of quasilinear first-order partial differential equations*

$$A_j^{0,i} u_t^j(x) + A_j^{\mu,i}(u(x)) u_\mu^j(x) = \sigma^i(u(x)). \quad (3.1)$$

We are interested in the following subclasses:

**Definition 3.2.2.** *A system of quasilinear first-order PDEs is called hyperbolic if the eigenvalue problem*

$$[A(\xi) + -1\lambda A^0]v = 0 \quad (3.2)$$

*has real eigenvalues and  $n$  linearly independent eigenvectors for every  $\xi \in \mathcal{S}^{[d-1]}$ , where*

$$A : \mathcal{S}^{[d-1]} \rightarrow T_1^1 \mathcal{U}^n, A(\xi) = A^\mu \xi_\mu. \quad (3.3)$$

*If  $A^0 = \text{Id}$  this translates to  $A(\xi)$  being diagonalizable over the real numbers for every direction  $\xi \in \mathcal{S}^{[d-1]}$ .*

**Note:** Unless otherwise noted, we will assume  $A^0 = \text{Id}$ .

**Definition 3.2.3.** *A system of quasilinear first-order PDEs is called a system of balance laws if there exist smooth flux functions  $F^\mu \in \Gamma(T\mathcal{U}^n)$  such that*

$$\frac{\partial F^{\mu,i}}{\partial u^j} = A_j^{\mu,i} \quad (3.4)$$

*for  $\mu = 0, \dots, d$ . If  $\sigma = 0$  it is called a system of conservation laws.*

**Definition 3.2.4.** A system of quasilinear first-order PDEs is called symmetric hyperbolic if  $A^0$  is symmetric positive definite and all  $A^\mu$  are symmetric. A system that can be brought into this form is called symmetrizable.

**Definition 3.2.5.** A pair  $(\eta, q)$  is called an entropy-entropy flux pair to the system of quasilinear first-order PDEs if the additional balance law

$$\frac{\partial \eta}{\partial u^i} \left( A_j^{0,i} u_t^j(x) + A_j^{\mu,i}(u(x)) u_\mu^j(x) \right) = \eta_t(u(x)) + q_\mu^\mu(u(x)) \quad (3.5)$$

is implied by equation (3.1).

**Lemma 3.2.6.** A system of quasilinear first-order PDEs admits an entropy-entropy flux pair  $(\eta, q)$  if the compatibility condition

$$\frac{\partial q^\mu}{\partial u^j} = \frac{\partial \eta}{\partial u^i} A_j^{\mu,i} \quad (3.6)$$

is satisfied.

**Lemma 3.2.7.** Given an entropy density  $\eta$ , a system of quasilinear first-order PDEs admits an entropy-entropy flux pair  $(\eta, q)$  if and only if the integrability condition

$$\left[ \frac{\partial}{\partial u^k} \left( \frac{\partial \eta}{\partial u^i} A_j^{\mu,i} \right) \right]_{[j,k]} = 0 \quad (3.7)$$

is satisfied.

*Proof.* Follows from lemma 3.2.6 combined with the requirement

$$\left[ \frac{\partial^2 q_\mu}{\partial u^j \partial u^k} \right]_{[j,k]} = \left[ \frac{\partial}{\partial u^k} \left( \frac{\partial \eta}{\partial u^i} A_j^{\mu,i} \right) \right]_{[j,k]} = 0 \quad (3.8)$$

and from the assertion that the  $n$ -dimensional target manifold  $\mathcal{U}^n$  is simply connected (see e.g. [Zor16], section 14.3).  $\square$

**Lemma 3.2.8.** A system of balance laws admitting an entropy-entropy flux pair  $(\eta, q)$  with strictly convex entropy density  $\eta$  is symmetrizable.

*Proof.* See e.g. [FL71].  $\square$

**Lemma 3.2.9.** A system of quasilinear first-order PDEs can be written as a system of balance laws in original coordinates  $u$  if

$$\left[ \frac{\partial A_j^{\mu,i}}{\partial u^k} \right]_{[j,k]} = 0 \quad (3.9)$$

holds. WILL ADD CONTENT FROM [Bog11].

*Proof.* Follows from the requirement

$$\left[ \frac{\partial^2 F^{\mu,i}}{\partial u^j \partial u^k} \right]_{[j,k]} = \left[ \frac{\partial A_j^{\mu,i}}{\partial u^k} \right]_{[j,k]} = 0 \quad (3.10)$$

and from the assertion that the  $n$ -dimensional target manifold  $\mathcal{U}^n$  is simply connected.  $\square$

It has been shown that if the entropy and the fluxes are *homogeneous*, symmetrizable, admitting an entropy-entropy flux pair and having a so called *skew-selfadjoint form* are all equivalent (see [Tad84], theorem 2.1 for details).  
WILL ADD DEFINITION, LEMMA AND DISCUSSION.

A weaker requirement than homogeneity and having a skew-selfadjoint form is the following:

**Lemma 3.2.10.** *Consider an entropy functional  $H = \int \eta(u(x)) dx$  and a skew-symmetric bilinear form*

$$L(v, w) = \left\langle v_i, \left( g^{\mu,i,j} \frac{\partial}{\partial x_\mu} + b_k^{\mu,i,j} u_\mu^k \right) w_j \right\rangle. \quad (3.11)$$

*Then the evolution equation*

$$u_t^i = \left[ L \frac{\partial \eta}{\partial u} \right]^i = \left( g^{\mu,i,j} \eta_{j,k} + b_k^{\mu,i,j} \eta_j \right) u_\mu^k \quad (3.12)$$

*admits an entropy-entropy flux pair  $(\eta, q)$ . If*

$$\left[ \frac{\partial}{\partial u^k} \left( b_l^{\mu,j,i} \eta_j \right) \right]_{[k,l]} = 0, \quad (3.13)$$

*it is a system of balance laws. If the entropy density  $\eta$  is strictly convex, the system is symmetrizable.*

*Proof.* First we note that from the skew-symmetry of  $L$  it follows that

$$2L(v, w) = 2 \left\langle v_i, b_k^{\mu,i,j} u_\mu^k w_j + g^{\mu,i,j} w_{j,\mu} \right\rangle \quad (3.14)$$

$$= L(v, w) - L(w, v) \quad (3.15)$$

$$= 2 \left\langle v_i, b_k^{\mu,i,j} u_\mu^k w_j \right\rangle_{[v,w]} + \frac{1}{2} (\langle v_i, g^{\mu,i,j} w_{j,\mu} \rangle - \langle w_i, g^{\mu,i,j} v_{j,\mu} \rangle) \quad (3.16)$$

$$= 2 \left\langle v_i, \left[ b_k^{\mu,i,j} \right]_{[i,j]} u_\mu^k w_j \right\rangle + \left\langle v_j, \frac{\partial}{\partial x_\mu} (g^{\mu,i,j} w_i) \right\rangle + \langle v_i, g^{\mu,i,j} w_{j,\mu} \rangle \quad (3.17)$$

$$= 2 \left\langle v_i, \left[ b_k^{\mu,i,j} \right]_{[i,j]} u_\mu^k w_j \right\rangle + \left\langle v_j, \frac{\partial g^{\mu,i,j}}{\partial u^k} u_\mu^k w_i \right\rangle + 2 \left\langle v_i, \left[ g^{\mu,i,j} \right]_{(i,j)} w_{j,\mu} \right\rangle \quad (3.18)$$

$$= \left\langle v_i, \underbrace{\left( 2 \left[ b_k^{\mu,i,j} \right]_{[i,j]} + \frac{\partial g^{\mu,i,j}}{\partial u^k} \right)}_{2b_k^{\mu,i,j}} u_\mu^k w_j + \underbrace{2 \left[ g^{\mu,i,j} \right]_{(i,j)}}_{2g^{\mu,i,j}} w_{j,\mu} \right\rangle. \quad (3.19)$$

Hence, we get

$$2 \left[ b_k^{\mu,i,j} \right]_{(i,j)} = \frac{\partial g^{\mu,i,j}}{\partial u^k} \text{ and } \left[ g^{\mu,i,j} \right]_{[i,j]} = 0. \quad (3.20)$$

**To check the existence of an entropy-flux function:** From equation (3.12) we get

$$\eta_t(u(x)) = \frac{\partial \eta}{\partial u^i} \left( g^{\mu,i,j} \eta_{j,k} + b_k^{\mu,i,j} \eta_j \right) u_\mu^k \quad (3.21)$$

and thus identify the *gradient of the entropy-flux function*

$$\frac{\partial q^\mu}{\partial u^k} = \frac{\partial \eta}{\partial u^i} \left( b_k^{\mu,i,j} \frac{\partial \eta}{\partial u^j} + g^{\mu,i,j} \frac{\partial^2 \eta}{\partial u^j \partial u^k} \right) \quad (3.22)$$

and can check the conditions of lemma 3.2.7:

$$\left[ \frac{\partial^2 q^\mu}{\partial u^k \partial u^l} \right]_{[k,l]} = \left[ \frac{\partial}{\partial u^l} \left( \frac{\partial \eta}{\partial u^i} \left( b_k^{\mu,i,j} \frac{\partial \eta}{\partial u^j} + g^{\mu,i,j} \frac{\partial^2 \eta}{\partial u^j \partial u^k} \right) \right) \right]_{[k,l]} \quad (3.23)$$

$$= \left[ \underbrace{\eta_{i,l} b_k^{\mu,i,j} \eta_j}_{\eta_i b_k^{\mu,j,i} \eta_{j,l}} + \underbrace{\eta_{i,l} g^{\mu,i,j} \eta_{j,k}}_{[() ]_{[k,l]} = 0} + \dots \right]_{[k,l]} \quad (3.24)$$

$$= \left[ \dots + \eta_i \left( \frac{\partial}{\partial u^l} \left( b_k^{\mu,i,j} \eta_j \right) + \underbrace{\frac{\partial}{\partial u^l} (g^{\mu,i,j} \eta_{j,k})}_{[() ]_{[k,l]} = [g_l^{\mu,i,j} \eta_{j,k}]_{[k,l]}} \right) \right]_{[k,l]} \quad (3.25)$$

$$= \left[ \eta_i \left( b_k^{\mu,j,i} \eta_{j,l} + \frac{\partial}{\partial u^l} \left( b_k^{\mu,i,j} \eta_j \right) + g_l^{\mu,i,j} \eta_{j,k} \right) \right]_{[k,l]} \quad (3.26)$$

$$= \left[ \eta_i \left( \underbrace{2 \left[ b_k^{\mu,j,i} \right]_{(i,j)}}_{g_k^{\mu,i,j}} \eta_{j,l} + \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \eta_j + g_l^{\mu,i,j} \eta_{j,k} \right) \right]_{[k,l]} \quad (3.27)$$

$$= \left[ \underbrace{\eta_i \left( g_k^{\mu,i,j} \eta_{j,l} + g_l^{\mu,i,j} \eta_{j,k} \right)}_{[O]_{[k,l]}=0} + \underbrace{\eta_i \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \eta_j}_{=\eta_i \left[ \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \right]_{(i,j)} \eta_j = \frac{1}{2} \eta_i \frac{\partial g_k^{\mu,i,j}}{\partial u^l} \eta_j} \right]_{[k,l]} \quad (3.28)$$

$$= \left[ \frac{1}{2} \eta_i \frac{\partial^2 g^{\mu,i,j}}{\partial u^k \partial u^l} \eta_j \right]_{[k,l]} \quad (3.29)$$

$$= 0. \quad (3.30)$$

**To check the existence of flux functions:** From the evolution equation we identify  $A_k^{\mu,i} = g^{\mu,i,j} \eta_{j,k} + b_k^{\mu,i,j} \eta_j$ . We check

$$\left[ \frac{\partial A_k^{\mu,i}}{\partial u^l} \right]_{[k,l]} = \left[ \frac{\partial}{\partial u^l} \left( g^{\mu,i,j} \eta_{j,k} + b_k^{\mu,i,j} \eta_j \right) \right]_{[k,l]} \quad (3.31)$$

$$= \left[ \frac{\partial}{\partial u^l} \left( g^{\mu,i,j} \eta_{j,k} \right) + \frac{\partial}{\partial u^l} \left( b_k^{\mu,i,j} \eta_j \right) \right]_{[k,l]} \quad (3.32)$$

$$= \left[ \frac{\partial g^{\mu,i,j}}{\partial u^l} \eta_{j,k} + g^{\mu,i,j} \frac{\partial \eta_{j,k}}{\partial u^l} + \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \eta_j + b_k^{\mu,i,j} \frac{\partial \eta_j}{\partial u^l} \right]_{[k,l]} \quad (3.33)$$

$$= \left[ b_l^{\mu,j,i} \eta_{j,k} + \underbrace{b_l^{\mu,i,j} \eta_{j,k} + b_k^{\mu,i,j} \eta_{j,l}}_{[O]_{[k,l]}=0} + \eta_j \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \right]_{[k,l]} \quad (3.34)$$

$$= \left[ \frac{\partial}{\partial u^k} \left( b_l^{\mu,j,i} \eta_j \right) + -1 \frac{\partial b_l^{\mu,j,i}}{\partial u^k} \eta_j + \eta_j \frac{\partial b_k^{\mu,i,j}}{\partial u^l} \right]_{[k,l]} \quad (3.35)$$

$$= \left[ \frac{\partial}{\partial u^k} \left( b_l^{\mu,j,i} \eta_j \right) + \eta_j \frac{\partial^2 g^{\mu,i,j}}{\partial u^k \partial u^l} \right]_{[k,l]} \quad (3.36)$$

$$= \left[ \frac{\partial}{\partial u^k} \left( b_l^{\mu,j,i} \eta_j \right) \right]_{[k,l]} \quad (3.37)$$

which completes the proof.  $\square$

### 3.2.1 Symmetric hyperbolic thermodynamically compatible systems

### 3.2.2 Skew-selfadjoint hyperbolic systems of balance laws

### 3.3 Poisson brackets of hydrodynamic type

The main references are

- [LGPV13] Standard Poisson structures reference
- [Nov82], [DN83a], [DN83b] (Slightly erroneous) initial papers
- [GD80b], [GD80a], [GD82] preceeding papers
- [Gri85] (Slightly erroneous) treatment of degenerate brackets of **constant rank**
- [Mok89] Correction of [DN83a]
- [Mok92] Correction of [Gri85]
- [Bog07b], [Bog07c] Extension of [Gri85]
- [Sav16b] Extension of [Gri85]
- [Mor80] (Slightly erroneous) treatment of Maxwell-Vlasov
- [MG80] Hamiltonian MHD
- [WM81] Correction of [Mor80]
- [KHO01] Automatic Jacobi-check
- [KH10] Improvement of [KHO01]
- [KVV17] Inaccessible, more general improvement of [KH10], [KHO01]
- [Vit19] Like [KVV17]
- [Rey10] Hydrodynamic Poisson structures
- [Sav16a] Hydrodynamic Poisson structures
- [MF90], [Mok92] Nonlocal generalization

**Definition 3.3.1.** *A bilinear bracket*

$$\{\cdot, \cdot\} : V \times V \rightarrow \mathbb{R} \quad (3.38)$$

*is called Poisson if  $\{\cdot, \cdot\}$*

- *is antisymmetric:*  $\{F, G\} = -\{G, F\}$ ,
- *is a derivation:*  $\{FG, H\} = F\{G, H\} + G\{F, H\}$  and
- *satisfies the Jacobi-identity:*  $\{F, \{G, H\}\} \circ_{F, G, H} = 0$

*JUST ADDED THIS TO BE ABLE TO REFERENCE IT, BUT THE BRACKET IS REALLY A BIVECTOR-field, I.E. IT MAY DEPEND ARBITRARILY ON U. WILL MOVE TO POISSON SECTION AND UPDATE.*

In [DN83a] and [DN83b] Dubrovin and Novikov introduced Poisson brackets and Hamiltonians of *hydrodynamic type*:

**Definition 3.3.2.** *A Poisson bracket  $\{\cdot, \cdot\}$  is called hydrodynamic if it satisfies*

$$\{u^i(x), u^j(y)\} = g^{\mu, i, j}(u(x))[\delta']_{\mu}(x-y) + u_{\mu}^k b_k^{\mu, i, j}(u(x))\delta(x-y). \quad (3.39)$$

**Definition 3.3.3.** *A functional (or Hamiltonian)*

$$H : C^{\infty}(\mathcal{M}^d; \mathcal{U}^n) \rightarrow \mathbb{R} \quad (3.40)$$



is called hydrodynamic if it is of the form

$$H[u] = \int h(u(x)) dx \quad (3.41)$$

for some energy density  $h \in C^\infty(\mathcal{U}^n)$ .

By requiring that brackets and Hamiltonians do not depend themselves on spatial derivatives of the unknown field variables  $u \in C^\infty(\mathcal{M}^d; \mathcal{U}^n)$ , we ensure that the resulting evolution equation will be a system of quasilinear first-order PDEs.

For any two Functionals  $F(u) = \int f(u(x)) dx$  and  $H(u) = \int h(u(x)) dx$  we get

$$\{F, H\} = \int \frac{\partial f}{\partial u^i} L^{i,j} \frac{\partial h}{\partial u^j} dx \quad (3.42)$$

with differential operator  $L^{i,j} = g^{\mu,i,j} \frac{\partial}{\partial x_\mu} + b_k^{\mu,i,j} u_\mu^k$ . Finally, given a Poisson bracket and a hydrodynamic Hamiltonian functional  $H$  we get the *evolution equation*

$$u_t^i = L^{i,j} h_j = \left( g^{\mu,i,j} h_{j,k} + b_k^{\mu,i,j} h_j \right) u_\mu^k, \quad (3.43)$$

where we can identify the  $(1,1)$ -tensors

$$A_k^{\mu,i} = g^{\mu,i,j} h_{j,k} + b_k^{\mu,i,j} h_j. \quad (3.44)$$

For the above bracket to be Poisson, it has to satisfy the *Jacobi identity*

$$\{F, \{G, H\}\}_{\circlearrowleft F, G, H} = 0 \quad (3.45)$$

for all hydrodynamic Functionals  $F, G, H$  which, while innocent looking, translates to (see [Mok98], Lemma 2.1):

$$0 = [g^{\mu,i,j}]_{[i,j]} \quad (3.46)$$

$$0 = \frac{\partial g^{\mu,i,j}}{\partial u^k} + -2 [b_k^{\mu,i,j}]_{(i,j)} \quad (3.47)$$

$$0 = \left[ [g^{\mu,i,l} b_l^{\nu,j,k}]_{[i,j]} \right]_{(\mu,\nu)} \quad (3.48)$$

$$0 = \left[ [g^{\mu,i,l} b_l^{\nu,j,k}]_{[(i,\mu),(j,\nu)]} \right]_{\circlearrowleft i,j,k} \quad (3.49)$$

$$0 = \left[ g^{\mu,i,l} \left[ \frac{\partial b_l^{\nu,j,k}}{\partial u^m} \right]_{[m,l]} + [b_l^{\mu,i,j} b_m^{\nu,l,k}]_{[j,k]} \right]_{(\mu,\nu)} \quad (3.50)$$

$$0 = \left[ g^{\mu,i,l} \frac{\partial b_m^{\nu,j,k}}{\partial u^l} + -1 b_l^{\nu,i,j} b_m^{\mu,l,k} + -1 b_l^{\nu,i,k} b_m^{\mu,j,l} \right]_{[(i,\mu),(j,\nu)]} \quad (3.51)$$

$$0 = \left[ \frac{\partial}{\partial u^n} \left( g^{\mu,i,l} \left[ \frac{\partial b_l^{\nu,j,k}}{\partial u^m} \right]_{[m,l]} + \left[ b_l^{\mu,i,j} b_m^{\nu,l,k} \right]_{[j,k]} \right) - \left[ b_m^{\mu,l,i} \left[ \frac{\partial b_l^{\nu,j,k}}{\partial u^m} \right]_{[m,l]} \right]_{\odot i,j,k} \right]_{((\mu,m),(\nu,n))}. \quad (3.52)$$

**Definition 3.3.4.** A collection of metrics  $g^\mu$  is called Poisson, if there exists a collection  $b^\mu$  such that the pair  $(g, b)$  induces a hydrodynamic Poisson bracket.

### 3.3.1 with nondegenerate metric

**Definition 3.3.5.** A metric  $g(u)$  is called non-degenerate if  $\det(g(u)) \neq 0$  everywhere.

**Definition 3.3.6.** A Poisson bracket (3.39) is called non-degenerate if all metrics  $g^\mu(u)$  are non-degenerate.

#### 3.3.1.1 in one spatial dimension

The main result, due to Dubrovin and Novikov ([DN83b], theorem 1) and most succinctly stated in [Mok00], is the following

**Theorem 3.3.7.** Let  $\det(g(u)) \neq 0$  and  $d = 1$ . The bracket (3.39) is a Poisson bracket if and only if  $g(u)$  is an arbitrary flat pseudo-Riemannian metric and  $b_k^{i,j}(u) = -1 g^{i,l}(u) \Gamma_{l,k}^j(u)$ , where  $\Gamma_{l,k}^j(u)$  is the Riemannian connection generated by the metric  $g(u)$ .

*Proof.* Stated without explicit proof in [DN83b]. Can be proved by ‘direct verification’.  $\square$

Due to flatness of the metric, we immediately get existence of (local) coordinates  $u$  where  $g^{i,j}(u) = e_i \delta_j^i$  with  $e_i = \pm 1$  and  $b(u) = 0$ . Hence all non-degenerate one-dimensional Poisson brackets are classified by the signature of their metric.

#### 3.3.1.2 in multiple spatial dimensions

A generalization of theorem 3.3.7, summarized in [Mok06], uses heavily the notion of *compatible metrics* developed in [Mok00]:

**Definition 3.3.8.** Two pseudo-Riemannian contravariant metrics  $g^{1,i,j}(u)$  and  $g^{2,i,j}(u)$  of constant Riemannian curvature  $K_1$  and  $K_2$ , respectively, are said to be compatible if an arbitrary linear combination

$$g^{i,j}(u) = \lambda_1 g^{1,i,j}(u) + \lambda_2 g^{2,i,j}(u) \quad (3.53)$$

of these metrics, where  $\lambda_1$  and  $\lambda_2$  are arbitrary constants such that  $\det(g^{i,j}(u)) \neq 0$ , is a metric of constant Riemannian curvature  $\lambda_1 K_1 + \lambda_2 K_2$  and the coefficients of the corresponding Levi-Civita connections are related by the same linear formula

$$\Gamma_k^{i,j}(u) = \lambda_1 \Gamma_{1,k}^{i,j}(u) + \lambda_2 \Gamma_{2,k}^{i,j}(u). \quad (3.54)$$

This definition naturally mirrors the definition of *compatible Poisson brackets*, of which linear combinations are again required to be Poisson brackets (see e.g. [Fer01] or [Mok02c]). Indeed, the following result is immediate:

**Theorem 3.3.9.** *The metrics  $g^{\mu,i,j}$  defining a multidimensional Poisson bracket of the form (3.39) are compatible.*

*Proof.* See [Mok98], chapter II.  $\square$

However, this is far from enough. Indeed, again from [Mok98], theorem 2.1:

**Theorem 3.3.10.** *Flat non-degenerate metrics  $g^{\mu,i,j}(u)$  and the Riemannian connections  $\Gamma_{j,k}^{\mu,i}$  defined by these metrics (the Levi-Civita connections) generate a Poisson structure (3.39) if and only if the tensors  $T_{j,k}^{\mu,\nu,i} = \Gamma_{j,k}^{\nu,i} - \Gamma_{j,k}^{\mu,i}$  satisfy the following relations (no summation over  $\mu$ ):*

$$[T^{\mu,\nu,i,j,k}]_{[i,k]} = 0 \quad (3.55)$$

$$[T_{j,k}^{\mu,\nu,i}]_{\odot i,j,k} = 0 \quad (3.56)$$

$$[T^{\mu,\nu,i,j,m} T_{m,l}^{\mu,\nu,k}]_{[k,l]} = 0 \quad (3.57)$$

$$\nabla_l^\mu T^{\mu,\nu,i,j,k} = 0 \quad (3.58)$$

where  $T^{\mu,\nu,i,j,k} = -1g^{\nu,k,m} T_m^{\mu,\nu,i,j} = -1g^{\nu,k,m} g^{\mu,i,n} T_{m,n}^{\mu,\nu,j}$  and  $\nabla_l^\mu$  is the covariant derivative associated with  $g^\mu$ .

Similarly to the case of one spatial dimension, there are multi-dimensional hydrodynamic Poisson brackets that can be reduced to *constant* form. However, contrary to the one-dimensional case, non-degeneracy of the associated metric(s) is not a sufficient condition. Instead, we need additionally that the *obstruction tensors*  $T_{j,k}^{\mu,\nu,i}$  are identically zero. We will need the concept of *non-singular* pairs of metrics:

**Definition 3.3.11.** *A pair of metrics  $g^{1,i,j}(u)$  and  $g^{2,i,j}(u)$  is called non-singular if the eigenvalues of this pair of metrics, i.e. the roots of the equation  $\det((g^{1,i,j}(u) + -1\lambda g^{2,i,j}(u))) = 0$ , are distinct.*

Finally we can state

**Theorem 3.3.12.** *If for a non-degenerate multidimensional Poisson bracket (3.39) one of the metrics  $g^{\mu,i,j}(u)$  forms non-singular pairs with all the remaining metrics of the bracket, then this Poisson bracket can be reduced to constant form by a local change of coordinates.*

And for all other (non-degenerate) Poisson brackets we have

**Theorem 3.3.13.** *Consider a Poisson bracket (3.39). If the metric  $g^{1,i,j}$  is non-degenerate, then in flat coordinates  $u$  where  $g^{1,i,j}(u)$  is constant and thus  $b_k^{1,i,j} = 0$ , every other metric  $g^{\mu,i,j}$  is linear.*

*Proof.* See [DN83a], theorem 1 and [Rey10], proposition 2.15.  $\square$

**Remark.** *Non-degeneracy of all metrics is not required.*

### 3.3.2 with degenerate metric

#### 3.3.2.1 in one spatial dimension

Following the classification of non-degenerate one-dimensional Poisson brackets (3.39) in [DN83a], Grinberg provided answers to the question of classification of *degenerate* one-dimensional Poisson brackets with metric  $g(u)$  of constant rank in [Gri85]. However, proofs were lacking in the literature until the publication of [Bog07b]. We will summarize the main statements.

For now, we do not require the  $(2,0)$ -tensor field  $g$  to be associated with a Poisson bracket. First, let us partition the  $n$ -dimensional target manifold  $\mathcal{U}^n$  by defining the subsets of rank  $k$

$$\mathcal{O}_k = \{u \in \mathcal{U}^n | \text{rank}(g(u)) = k\} \subset \mathcal{U}^n. \quad (3.59)$$

Next, to depart from the non-degenerate case, let us require that at least one  $\mathcal{O}_m$  is nonempty and open for some  $m < n$ . Now, in slight disagreement with [Bog07b], we have:

**Proposition 3.3.14.** *Consider a  $(2,0)$ -tensor field  $g$  with nonempty, open set  $\mathcal{O}_m$  as in 3.59. Then*

- *$g$  being smooth and Poisson does not imply  $\overline{\cup_{k=0}^{n-1} \mathcal{O}_k} = \mathcal{U}^n$ ,*
- *if  $g$  is analytic, then  $\mathcal{O}_m$  is dense in  $\mathcal{U}^n$ , i.e.  $\overline{\mathcal{O}_m} = \mathcal{U}^n$ .*

*Proof.* For the first point we will construct a counterexample. All functions used in this part will be smooth. Consider a diagonal metric

$$g(u) = \begin{bmatrix} g^1(u) & 0 \\ 0 & g^2(u) \end{bmatrix} \quad (3.60)$$

on the target manifold  $\mathcal{U}^n = \mathbb{R}^2$ . Together with

$$b(u) = \frac{1}{2} \nabla_u \begin{bmatrix} g^1(u) & b^{1,2}(u) \\ -b^{1,2}(u) & g^2(u) \end{bmatrix} \quad (3.61)$$

this induces a hydrodynamic Poisson bracket if

$$g^2 = b^{1,2} = 0 \quad (3.62)$$

or if

$$-g^2(u) = g^1(u) = b^{1,2}(u) = g^0(u^1 + u^2) \quad (3.63)$$

for all  $u \in \mathcal{U}^n$ , see `hydro_smooth_counterexample.ipynb`. We may stitch together the above families of Poisson brackets, as long as they are separated from each other, as for example as follows:

$$g^1(u) = g^0(u^1 + u^2) \quad (3.64)$$

$$-g^2(u) = \begin{cases} g^1(u) & \text{if } u^1 + u^2 < -\varepsilon, \\ 0 & \text{if } u^1 + u^2 \geq -\varepsilon \end{cases} \quad (3.65)$$

$$g^0(x) = \begin{cases} \neq 0 & \text{if } |x| > \varepsilon, \\ 0 & \text{if } |x| \leq \varepsilon \end{cases}, \quad (3.66)$$

with  $\varepsilon > 0$  and  $g^0$  constructed appropriately. We now have

$$\text{rank}(g) = \begin{cases} 2 & \text{if } u^1 + u^2 < -\varepsilon, \\ 0 & \text{if } |u^1 + u^2| \leq \varepsilon, \\ 1 & \text{if } u^1 + u^2 > \varepsilon \end{cases} \quad (3.67)$$

and thus

$$\overline{\cup_{k=0}^{n-1} \mathcal{O}_k} = \{u^1, u^2 \in \mathcal{U}^n \mid (u^1 + u^2) \in [-\varepsilon, \infty)\} \neq \mathcal{U}^n, \quad (3.68)$$

which completes the proof of the first part.

For the second part we use the following factoid (see e.g. [Hog13], chapter 2.4, fact 25):

A square matrix  $A$  has rank  $k$  if and only if  $A$  has a nonsingular  $k \times k$  submatrix, and every  $(k+1) \times (k+1)$  submatrix of  $A$  is singular.

As the determinant of a matrix with analytic coefficients is itself analytic, square submatrices of  $g$  that are singular on some open set  $\mathcal{O}_m \subset \mathcal{U}^n$  stay singular on the whole of the  $n$ -dimensional target manifold  $\mathcal{U}^n$ . For the same reason, square submatrices of  $g$  that are nonsingular on some open set  $\mathcal{O}_m \subset \mathcal{U}^n$  can not be singular on all of any open set. Given any  $x \in \mathcal{U}^n$ , this includes *any* open set  $\mathcal{O}' \subset \mathcal{U}^n$  containing  $x$ . Thus, every neighborhood of every point  $x \in \mathcal{U}^n$  contains at least one point from  $\mathcal{O}_m$ , which is thus dense in  $\mathcal{U}^n$ , i.e. its closure equals  $\mathcal{U}^n$  (see e.g. [Bou95], chapter 1, definition 10).  $\square$

**Remark.** The phrasing in [Bog07b] is slightly ambiguous. On page 540 he claims that “if the functions  $g^{i,j}(u)$  are smooth, [...] the [target manifold  $\mathcal{U}^n$ ] is the closure of the union  $\cup_{k=0}^{n-1} \mathcal{O}_k$ .”

Two sentences before, he talks about “any degenerate  $(2,0)$ -tensor  $g^{i,j}(u)$ ”, suggesting that the requirement for  $g$  to be Poisson can be dropped. The following theorem however relies on  $g$  being Poisson, so we must interpret the whole

section to carry this requirement implicitly. If indeed  $g$  were not required to be Poisson, an even simpler counterexample could have been constructed, as we are not constrained by the considerable requirements induced by the Jacobi identity. Next, the exact meaning of  $g$  being degenerate has not been stated in [Bog07b]. If it were to mean that  $\mathcal{O}_n$  is empty, the above statement would be trivially true, but then there would be no need of closing the union. The statement closest to a definition of degeneracy can be found on page 539: > In Section 2, we disclose the invariant meaning of the Poisson brackets for any values of  $k$  with degenerate  $(2,0)$ -tensor  $g^{i,j}(u)$  that has a constant rank  $g^{i,j} = m < n$  in an open domain  $\mathcal{O}_m$ . Note that the phrasing does not imply that  $\mathcal{O}_m$  is nonempty, that  $\mathcal{O}_m$  is defined as in 3.59, or that  $\mathcal{O}_n$  is empty. Again, if the requirement were not that at least one  $\mathcal{O}_m$  as defined in 3.59 with  $m < n$  is open (and nonempty), a simpler counterexample with a  $g$  that is Poisson could have been constructed.

Finally, at the end of Section 1, page 540 in [Bog07b] a further constraint is mentioned, namely  $1 < m < n$ . This constraint only seems to apply to Corollary 1 in [Bog07b], where it is explicitly invoked. Even if it were implied in the contested statement, we could simply construct a  $\hat{g}$  which is block-diagonal, with the  $g$  from the proof as one block and another matrix with arbitrary rank as the other block.

Cf. figure 3.1 for all constructions. For  $k < m$  we have  $\overline{\mathcal{O}_k} \cap \overline{\mathcal{O}_m} \subset \mathcal{O}_k$  and thus

- for the left figure  $\mathcal{O}_1$  is open but  $g$  is not Poisson,
- for the middle figure  $g$  is Poisson but  $\mathcal{O}_1$  is not open as it shares a boundary with  $\mathcal{O}_2$ ,
- for the right figure,  $g$  is Poisson and  $\mathcal{O}_1$  is open
- for all figures,  $\mathcal{O}_0$  is not open and  $\mathcal{O}_2$  is open.

**Theorem 3.3.15.** *Let  $L_u$  and  $\mathcal{O}_k$  be as in ?? and 3.59. Then*

- An invariant non-degenerate metric is defined on the distribution  $L_u$ ,
- The distribution  $L_u$  is invariant under any  $(1,1)$ -tensor  $A_j^i(u)$  (3.44) and operator  $b_k^{i,j}(u)$  for  $j = \text{const}$ ,
- In the open domain  $\mathcal{O}_m$ , the distribution  $L_u$  is involutive and defines an  $m$ -dimensional foliation  $\mathcal{F}_m$ ,
- The metric on the leaves of  $\mathcal{F}_m$  is flat.

*Proof.* See [Bog07b], theorem 1. □

### 3.3.2.2 in multiple spatial dimensions

### 3.3.3 of Lie-Poisson type

#### 3.3.3.1 in one spatial dimension

[BN85]

#### 3.3.3.2 in multiple spatial dimensions




Figure 3.1: Constructions for PROPOSITION. Colors and numbers indicate  $\text{Rank}(g(u))$ .

## 3.4 Hyperbolic Poisson brackets

Throughout, we will assume *vanishing of boundary terms*

$$\int \frac{\partial}{\partial x} (f(u(x))) dx = 0. \quad (3.69)$$

### 3.4.1 of constant type

For constant Poisson brackets, just like for linear hyperbolic systems of balance laws, many things simplify. Nevertheless, they have attracted some interest over the years, mainly in the context of multi-Hamiltonian systems, see e.g. [GN90]. Olver and Nutku examined one particular class of one-dimensional,  $2 \times 2$  constant hydrodynamic Poisson brackets in [ON88], following [Nut85] and [Nut87]. The same class has been connected to the shallow water equations in [CM82], to Poisson's equation in [Nut87], to the Born-Infeld equation in [ANN<sup>+</sup>89] and to Riemann invariants in [Ney89]. This class has been further investigated in [Car95]. Benjamin and Bowman considered discontinuous solutions of general one-dimensional constant Poisson brackets in [Ben87].

Here, we will only state some immediate results connecting constant hydrodynamic Poisson brackets to (possibly hyperbolic) systems of quasilinear first-order PDEs with an additional conservation law.

We adapt the customary definition of constant Poisson brackets on finite dimensional vector spaces as given e.g. in [LGPV13], definition 6.1:

**Definition 3.4.1.** *A hydrodynamic Poisson bracket  $\pi$  on the infinite-dimensional function space  $C^\infty(\mathcal{M}^d; \mathcal{U}^n)$  is called constant, if for each pair of linear functionals  $F$  and  $G$  of the form  $I = \int f_i(x) u^i(x) dx$ , their Poisson bracket  $\pi(F, G)$  is a constant function on  $C^\infty(\mathcal{M}^d; \mathcal{U}^n)$ .*

This translates to the equivalent conditions  $g^\mu = \text{symmconst}$  and  $b^\mu = 0$ , hence we have as evolution equation

$$u_t = g^\mu \nabla^2 h u_\mu. \quad (3.70)$$

As can be immediately verified from relations ??, any constant skew-symmetric bracket is a Poisson bracket. Furthermore, from lemma 3.2.10 we get:

**Corollary 3.4.2.** *Any constant hydrodynamic Poisson bracket induces a system of conservation laws with an additional balance law for the energy density. If the energy density is strictly convex, the system is symmetrizable.*

*Proof.* Follows from lemma 3.2.10 and  $b^\mu = 0$ . □

**Corollary 3.4.3.** *In one spatial dimension ( $d = 1$ ), all constant hydrodynamic Poisson brackets with positive definite matrix  $g$  induce a symmetrizable system of conservation laws for any hydrodynamic Hamiltonian.*

**Remark.** • The symmetrizer is  $g^{-1}$



- For  $d > 1$  the system is generally not symmetrizable.

**Corollary 3.4.4.** Any linear system of first-order PDEs

$$u_t + A^\mu u_\mu = 0 \quad (3.71)$$

that admits an additional conservation law for  $\eta(u)$  with non-singular and constant Hessian can be written using a constant hydrodynamic Poisson bracket with  $g^\mu = A^\mu \nabla^{-2} \eta$ .

*Proof.* From the existence of a conservation law for  $\eta(u)$  we get from the integrability condition (cf. equation (3.7))

$$\nabla^2 \eta A^\mu = (\nabla^2 \eta A^\mu)^T = (A^\mu)^T \nabla^2 \eta \quad (3.72)$$

and hence after multiplication with  $\nabla^{-2} \eta$  from the left and from the right

$$g^\mu = A^\mu \nabla^{-2} \eta = \nabla^{-2} \eta (A^\mu)^T = (A^\mu \nabla^{-2} \eta)^T = (g^\mu)^T = \text{symm.const}, \quad (3.73)$$

thus the  $g^\mu$  induce a Poisson bracket.  $\square$

**Remark.** • Generally we can assume  $\nabla^2 \eta = \text{const}$  if the system of first-order PDEs is linear.

- If  $\nabla^2 \eta$  is singular, the  $A^\mu$  are  $2 \times 2$  block-diagonal in coordinates in which  $\nabla^2 \eta$  is diagonal. The blocks associated with the non-zero eigenvalues of  $\nabla^2 \eta$  can be written using a constant hydrodynamic Poisson bracket, while we can say nothing about the other blocks.

We note that, while all linear hyperbolic systems of conservation laws with an additional conservation law admit a constant hydrodynamic Poisson bracket, not all linear evolution equations induced by constant hydrodynamic Poisson brackets are hyperbolic. Consider for example

$$g = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, h(u^1, u^2) = u^1 u^2 \implies A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}. \quad (3.74)$$

Similarly, only in one spatial dimension is positive definiteness of the matrices  $g^\mu$  sufficient for hyperbolicity. Consider for example

$$g^1 = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}, g^2 = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \implies g^1 - g^2 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad (3.75)$$

i.e. we may choose  $h$  as above such that in direction  $\xi = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  the system matrix

$A(\xi)$  is not real diagonalizable.

In fact, we may show the following:

**Corollary 3.4.5.** In one spatial dimension ( $d = 1$ ) a constant hydrodynamic Poisson bracket induces hyperbolic PDEs for all hydrodynamic functionals if and only if its metric  $g$  is definite or identically zero. In multiple spatial dimensions, in addition all metrics  $g^\mu$  have to be scalar multiples of each other.

*Proof.* For the one-dimensional case, we may simply construct a Hamiltonian as above in coordinates in which the metric is diagonal and contains only 1, 0 or  $-1$ . For the multi-dimensional case we note that from the 1D result it follows that for every *direction*  $\xi \in \mathcal{S}^{[d-1]}$  the matrix  $A(\xi)$  has to be either definite or identically zero. From continuity of eigenvalues of  $A(\xi)$  (see e.g. [Kat95], theorem 6.8) it follows that for all  $i, j$ , every curve connecting  $e_i$  and  $\pm e_j$  on  $\mathcal{S}^{[d-1]}$  has to pass a point  $\xi$  where  $A(\xi) = 0$ . This however can only happen, if  $A(e_i)$  is a scalar multiple of  $A(e_j)$ .  $\square$

Finally we want to note the following simple result, which will help us relate non-constant hydrodynamic Poisson brackets to constant ones:

**Corollary 3.4.6.** *Let  $g$  be the constant metric associated with a constant hydrodynamic Poisson bracket  $\pi$ . After an invertible change of variables*

$$\hat{u} = \hat{u}(u), \quad (3.76)$$

*we have as transformed expressions*

$$\hat{g}^{\mu, \hat{i}, \hat{j}} = \hat{u}_i^{\hat{i}} g^{\mu, i, j} \hat{u}_j^{\hat{j}} \quad (3.77)$$

$$\hat{b}_{\hat{k}}^{\mu, \hat{i}, \hat{j}} = \hat{u}_i^{\hat{i}} g^{\mu, i, j} \hat{u}_{j, k}^{\hat{j}} u_{\hat{k}}^k \quad (3.78)$$

where

$$\hat{u}_i^{\hat{i}} = [\nabla \hat{u}]_i^{\hat{i}} \quad (3.79)$$

$$\hat{u}_{j, k}^{\hat{j}} = [\nabla^2 \hat{u}]_{j, k}^{\hat{j}} \quad (3.80)$$

$$u_{\hat{k}}^k = [\hat{\nabla} u]_{\hat{k}}^k \quad (3.81)$$

*Proof.* A straightforward computation yields

$$\pi(F, H) = \left\langle F_i, g^{\mu, i, j} \partial_\mu (H_j) + \overbrace{b_k^{\mu, i, j}}^0 u_\mu^k H_j \right\rangle \quad (3.82)$$

$$= \left\langle \hat{F}_{\hat{i}} \hat{u}_i^{\hat{i}}, g^{\mu, i, j} \partial_\mu (\hat{u}_j^{\hat{j}} \hat{H}_{\hat{j}}) \right\rangle \quad (3.83)$$

$$= \left\langle \hat{F}_{\hat{i}}, \underbrace{\hat{u}_i^{\hat{i}} g^{\mu, i, j} \hat{u}_j^{\hat{j}}}_{\hat{g}^{\mu, \hat{i}, \hat{j}}} \partial_\mu (\hat{H}_{\hat{j}}) + \underbrace{\hat{u}_i^{\hat{i}} g^{\mu, i, j} \hat{u}_{j, k}^{\hat{j}} u_{\hat{k}}^k}_{\hat{b}_{\hat{k}}^{\mu, \hat{i}, \hat{j}}} \hat{u}_\mu^{\hat{k}} \hat{H}_{\hat{j}} \right\rangle \quad (3.84)$$

$$= \hat{\pi}(\hat{F}, \hat{H}). \quad (3.85)$$

$\square$

### 3.4.2 with non-degenerate metric

#### 3.4.2.1 in one spatial dimension

In [PPK20], the following result is stated (Theorem 2.10):

**Theorem 3.4.7.** *Consider a one-dimensional Hamiltonian system of hydrodynamic type with non-degenerate metric. Assuming that the energy of the system is of hydrodynamic type, convex and a proper scalar, it follows that the evolution equations can be regarded as a first-order quasilinear symmetric hyperbolic PDE system.*

This appears to be not correct. Consider the following diagonal metric  $g$  with matrix representation

$$g(u) = \begin{bmatrix} g^1(u^1) & 0 \\ 0 & g^2(u^2) \end{bmatrix}. \quad (3.86)$$

We get the corresponding differential operator

$$Lv = g(u) \frac{\partial v}{\partial x} + \frac{1}{2} \frac{\partial}{\partial u^k} (g(u)) u_x^k v \quad (3.87)$$

$$= \begin{bmatrix} g^1 & 0 \\ 0 & g^2 \end{bmatrix} \begin{bmatrix} \frac{\partial v_1}{\partial u^1} & \frac{\partial v_1}{\partial u^2} \\ \frac{\partial v_2}{\partial u^1} & \frac{\partial v_2}{\partial u^2} \end{bmatrix} \begin{bmatrix} u_x^1 \\ u_x^2 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} \frac{\partial g^1}{\partial u^1} u_x^1 & 0 \\ 0 & \frac{\partial g^2}{\partial u^2} u_x^2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \quad (3.88)$$

$$= \begin{bmatrix} g^1 \frac{\partial v_1}{\partial u^1} u_x^1 + g^1 \frac{\partial v_1}{\partial u^2} u_x^2 + \frac{\partial g^1}{\partial u^1} u_x^1 \frac{1}{2} v_1 \\ g^2 \frac{\partial v_2}{\partial u^1} u_x^1 + g^2 \frac{\partial v_2}{\partial u^2} u_x^2 + \frac{\partial g^2}{\partial u^2} u_x^2 \frac{1}{2} v_2 \end{bmatrix} \quad (3.89)$$

$$= \begin{bmatrix} g^1 \frac{\partial v_1}{\partial u^1} + \frac{1}{2} \frac{\partial g^1}{\partial u^1} v_1 & g^1 \frac{\partial v_1}{\partial u^2} \\ g^2 \frac{\partial v_2}{\partial u^1} & g^2 \frac{\partial v_2}{\partial u^2} + \frac{1}{2} \frac{\partial g^2}{\partial u^2} v_2 \end{bmatrix} \begin{bmatrix} u_x^1 \\ u_x^2 \end{bmatrix}, \quad (3.90)$$

defined by its action on a generic field  $v = \hat{v} \circ u \in C^\infty(\mathbb{R}; \mathbb{R}^2)$ . The induced bracket is Poisson, as can be verified by inserting

$$g^{i,j} = \delta^{i,j} g^i \quad (3.91)$$

$$b_k^{i,j} = \frac{1}{2} \frac{\partial g^{i,j}}{\partial u^k} = \frac{1}{2} \delta_k^{i,j} \frac{\partial g^i}{\partial u^k} \quad (3.92)$$

into relations ???. Here  $\delta_k^{i,j}$  is a generalization of the Kronecker delta, which is 1 exactly if all indices are equal and 0 otherwise. Inner products of  $g$  and  $b$  retain this structure, hence are symmetric in all free indices, and thus all terms in the Jacobi identity vanish.

Recall that (global) hyperbolicity means that the matrix in front of the spatial gradient of the unknown fields is diagonalizable over the real numbers

(everywhere). A necessary condition for a *real-valued matrix*

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad (3.93)$$

to be real diagonalizable is for the discriminant of its characteristic polynomial to be non-negative:

$$(a - d)^2 + 4bc \geq 0. \quad (3.94)$$

A sufficient condition for the above real-valued matrix  $A$  to be symmetric positive definite is

$$b = c, |b| < \min(a, d).$$

Let us now apply the above differential operator to the field associated with the variational derivative of some (strictly) convex hydrodynamic functional  $H(u) = \int h(u(x)) dx$ . We get

$$L^{i,j} \frac{\partial h}{\partial u^j} = \begin{bmatrix} g^1 \frac{\partial^2 h}{\partial u^1 \partial u^1} + \frac{1}{2} \frac{\partial g^1}{\partial u^1} \frac{\partial h}{\partial u^1} & g^1 \frac{\partial^2 h}{\partial u^1 \partial u^2} \\ g^2 \frac{\partial^2 h}{\partial u^2 \partial u^1} & g^2 \frac{\partial^2 h}{\partial u^2 \partial u^2} + \frac{1}{2} \frac{\partial g^2}{\partial u^2} \frac{\partial h}{\partial u^2} \end{bmatrix} \begin{bmatrix} u_x^1 \\ u_x^2 \end{bmatrix} \quad (3.95)$$

$$= A \begin{bmatrix} u_x^1 \\ u_x^2 \end{bmatrix} \quad (3.96)$$

and  $\nabla_i \nabla_j h = [g^{-1} A]_{i,j}$  where  $\nabla_i$  represents the covariant derivative associated with the Levi-Civita connection of the metric  $g$ .

We observe:

- the metric  $g$  is flat and the above differential operator induces a Poisson bracket
- the metric  $g$  is non-degenerate (everywhere), if  $g^1 \neq 0 \neq g^2$  (everywhere)
- the energy functional  $H$  is of hydrodynamic type
- the energy functional/density is (strictly) convex if  $\left| \frac{\partial^2 h}{\partial u^1 \partial u^2} \right| < \min(\frac{\partial^2 h}{\partial u^1 \partial u^1}, \frac{\partial^2 h}{\partial u^2 \partial u^2})$
- the energy density  $h$  is a proper scalar, if  $u^1$  and  $u^2$  are proper scalars.

However

- the signs  $\sigma(g^1)$  and  $\sigma(g^2)$  do not matter for non-degeneracy
- the first order partial derivatives of the energy density  $\frac{\partial h}{\partial u^1}$  and  $\frac{\partial h}{\partial u^2}$  do not matter for convexity
- the matrix defined by the entries  $\nabla_i \nabla_j h$ , while symmetric, is **not** necessarily positive definite

and we can thus choose our non-degenerate metric  $g$  and construct a (strictly) convex energy density  $h$  such that for some point  $u^*$




Figure 3.2: "Excess hyperbolicity"  $((a-d)^2 + 4bc)$  for  $g^i = (-1)^i(1 + u^{i^2})$  and  $h(u) = \left\langle u, \frac{1}{2} \begin{bmatrix} 1 & 1-\varepsilon \\ 1-\varepsilon & 1 \end{bmatrix} u + \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\rangle$ ,  $\varepsilon = 0.1$ . White corresponds to non-hyperbolic regions.

- $g^1 < 0 < g^2$
- $0 < \left| \frac{\partial^2 h}{\partial u^1 \partial u^2} \right| < \min\left(\frac{\partial^2 h}{\partial u^1 \partial u^1}, \frac{\partial^2 h}{\partial u^2 \partial u^2}\right)$
- $\frac{1}{2} \frac{\partial g^i}{\partial u^i} \frac{\partial h}{\partial u^i} = -1 g^i \frac{\partial^2 h}{\partial u^i \partial u^i}$  for  $i = 1, 2$ .

The first two points ensure that the product of the off-diagonal entries is (strictly) negative, whilst the last point ensures that the diagonal entries vanish, hence yielding non-hyperbolicity at  $u^*$  and in a neighbourhood around  $u^*$ .

See also figure 3.2 for a slightly different example.

We may wonder whether there is something peculiar about the above constructions. It is obvious that the above construction would not have worked, if the metric were definite or if  $\frac{\partial^2 h}{\partial u^1 \partial u^2} = 0$ . Are there special Poisson brackets or Hamiltonians which guarantee hyperbolicity? The following Lemma gives the answer:

**Lemma 3.4.8.** *Consider a Poisson bracket of hydrodynamic type with non-degenerate metric  $g$  in one space dimension and with arbitrary number of fields  $n \geq 2$ .*

- *If the metric is definite, every Hamiltonian of hydrodynamic type induces a system of hyperbolic PDEs.*
- *On the other hand, for every (non-constant) Hamiltonian of hydrodynamic type, we can construct a Poisson bracket with non-degenerate metric such*

that the induced system of PDEs is non-hyperbolic on a set of non-zero measure.

*Proof.* We know that the metric  $g$  is flat. If it is definite, for every point  $p \in \mathcal{U}^n$  we may find local coordinates  $\hat{u}$  such that

$$g_{i,j} = \pm \delta_{i,j} \quad (3.97)$$

on a neighborhood of  $p$ . On this neighborhood  $g$  is constant, hence the Christoffel symbols of the second kind (cf. equation (2.1)) vanish and thus  $b_k^{i,j}$  also does.

In these coordinates, the evolution equation reads

$$\hat{u}_t = \hat{L} \hat{\nabla} h = g \hat{\nabla}^2 \hat{h} \hat{u}_x \quad (3.98)$$

$$= \pm \hat{\nabla}^2 \hat{h} \hat{u}_x. \quad (3.99)$$

Due to symmetry of the Hessian, the system is hyperbolic. In fact, if we could ensure that the Hessian of the energy density were positive definite in those flat coordinates  $\hat{u}$ , we would also immediately get hyperbolicity for indefinite metrics. However, given a Hamiltonian of hydrodynamic type, we can in general guarantee neither this sufficient condition nor a weaker one.

Let us first look at the  $2 \times 2$  case. Let

$$\hat{h}_{i,\hat{j}} = \left[ \hat{\nabla}^2 \hat{h} \right]_{\hat{i},\hat{j}} \quad (3.100)$$

denote the entries of the Hessian in flat coordinates  $\hat{u}$  where the metric is constantly

$$\hat{g} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (3.101)$$

Then

$$A = \hat{g} \hat{\nabla}^2 \hat{h} \quad (3.102)$$

is real diagonalizable if and only if

$$\left| \hat{h}_{1,1} + \hat{h}_{2,2} \right| \geq 2 \left| \hat{h}_{1,2} \right|, \quad (3.103)$$

which is weaker than convexity of  $h$ .

The question now is, given some energy density  $h$ , which is possibly (strictly) convex in non-flat, “original” coordinates  $u$ , can we somehow guarantee hyperbolicity based on properties of the energy density alone, independently of the Poisson bracket? The answer, as stated in the lemma, is no:

Consider coordinates

$$\begin{bmatrix} u^1 \\ u^2 \end{bmatrix} = \begin{bmatrix} \hat{u}^1 \\ f(\hat{u}^1, \hat{u}^2) \end{bmatrix} \quad (3.104)$$

for some smooth real-valued function  $f$  with  $f' \neq 0$ . Denoting as entries of the Jacobian  $u_{\hat{i}}^i = \hat{\nabla}_{\hat{i}} u^i$  we get

$$\hat{h}_{\hat{i},\hat{j}} = h_{i,j} u_{\hat{i}}^i u_{\hat{j}}^j + h_i u_{\hat{i},\hat{j}}^i \quad (3.105)$$

$$= \begin{bmatrix} h_{i,j} u_1^i u_1^j + h_i u_{1,1}^i & h_{i,j} u_1^i u_2^j + h_i u_{1,2}^i \\ h_{i,j} u_2^i u_1^j + h_i u_{2,1}^i & h_{i,j} u_2^i u_2^j + h_i u_{2,2}^i \end{bmatrix} \quad (3.106)$$

$$= \begin{bmatrix} h_2 f_{1,1} + f_1^2 h_{2,2} + 2f_1 h_{1,2} + h_{1,1} & h_2 f_{1,2} + f_1 f_2 h_{2,2} + f_2 h_{1,2} \\ h_2 f_{2,1} + f_1 f_2 h_{2,2} + f_2 h_{1,2} & h_2 f_{2,2} + f_2^2 h_{2,2} \end{bmatrix}. \quad (3.107)$$

We get

$$|h_2 f_{1,1} + h_2 f_{2,2} + f_1^2 h_{2,2} + 2f_1 h_{1,2} + f_2^2 h_{2,2} + h_{1,1}| \geq 2|h_2 f_{1,2} + f_1 f_2 h_{2,2} + f_2 h_{1,2}| \quad (3.108)$$

as a necessary condition for hyperbolicity. Without loss of generality we will assume  $h_2(0) \neq 0$ . Note that given some function  $f$ , we may construct a function  $\hat{f}(x) = \frac{\hat{f}(cx)}{c}$  such that  $\hat{f}_i(0) = f_i(0)$  but  $\hat{f}_{i,j}(0) = c f_{i,j}(0)$  for an arbitrary constant  $c \neq 0$ . We achieve violation of the above inequality by choosing an  $f$  such that

$$2|f_{1,2}| \geq |f_{1,1} + f_{2,2}| \quad (3.109)$$

at the origin and then constructing an appropriate  $\hat{f}$  with a large enough  $c$ . One possible choice of  $f$  is

$$f(x, y) = \frac{1}{2} (1 + (1 + x)^2) (y + y^3). \quad (3.110)$$

Let us now look at the case  $n > 2$ . Given a non-constant energy density  $h$ , we will construct a Poisson bracket with non-degenerate indefinite metric  $g$  such that the induced PDE is non-hyperbolic on a set of non-zero measure. As above we will assume  $h_2(0) \neq 0$ . We will extend the constructions above. Let the metric in flat coordinates be

$$\hat{g} = \begin{bmatrix} 1 & 0 & 0_{1,[n-2]} \\ 0 & -1 & 0_{1,[n-2]} \\ 0_{[n-2],1} & 0_{[n-2],1} & I_{[n-2],[n-2]} \end{bmatrix} \quad (3.111)$$

and consider coordinates

$$\begin{bmatrix} u^1 \\ u^2 \\ u^3 \\ \vdots \end{bmatrix} = \begin{bmatrix} \hat{u}^1 \\ f(\hat{u}^1, \hat{u}^2) \\ \hat{u}^3 \\ \vdots \end{bmatrix}. \quad (3.112)$$

Then we get

$$\hat{h}_{\hat{i},\hat{j}} = \begin{cases} h_2 f_{1,1} + f_1^2 h_{2,2} + 2f_1 h_{1,2} + h_{1,1} & \text{if } \hat{i} = \hat{j} = 1, \\ h_2 f_{1,2} + f_1 f_2 h_{2,2} + f_2 h_{1,2} & \text{if } \text{sorted}(\hat{i}, \hat{j}) = 1, 2, \\ h_2 f_{2,2} + f_2^2 h_{2,2} & \text{if } \hat{i} = \hat{j} = 2, \\ h_{1,\hat{j}} + h_{2,\hat{j}} f_1 & \text{if } \min(\hat{i}, \hat{j}) = 1 \text{ and } \max(\hat{i}, \hat{j}) \geq 3, \\ h_{2,\hat{j}} f_2 & \text{if } \min(\hat{i}, \hat{j}) = 2 \text{ and } \max(\hat{i}, \hat{j}) \geq 3, \\ h_{\hat{i},\hat{j}} & \text{else} \end{cases} \quad (3.113)$$

Note that for the  $f$  given above, at the origin we have  $f_{1,1} = f_{2,2} = f_1 = 0$  and  $f_2 = 1$ , hence

$$\hat{h}_{\hat{i},\hat{j}}(0) = \begin{cases} h_{1,2}(0) + f_{1,2}(0)h_2(0) & \text{if } \text{sorted}(\hat{i}, \hat{j}) = 1, 2, \\ h_{\hat{i},\hat{j}}(0) & \text{else} \end{cases} \quad (3.114)$$

For arbitrary  $n > 2$ , we of course do not have the luxury of an explicit expression for the eigenvalues of a general matrix. However, we may still construct a matrix such that we can ensure that it has at least one eigenvalue pair with non-zero imaginary part. We will use Gerschgorin's theorem (cf. e.g. [Ste90], chapter IV, theorem 2.1), ensuring that at least two Gerschgorin disks are each separated from all others while not touching the real line.

Consider

$$\hat{g}\hat{\nabla}^2\hat{h} = \underbrace{\begin{bmatrix} h_{1,1} & 0 & h_{1,3} & \dots \\ 0 & -h_{2,2} & 0 & \dots \\ h_{1,3} & 0 & h_{3,3} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}}_{A_{\text{Symm}}} + \underbrace{\begin{bmatrix} 0 & \hat{h}_{1,2} & 0 & \dots \\ -\hat{h}_{1,2} & 0 & -h_{2,3} & \dots \\ 0 & h_{2,3} & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}}_{A_{\text{Skew}}} \quad (3.115)$$

at the origin. The matrix  $A_{\text{Skew}}$  has non-zero eigenvalues

$$\lambda_{1,2} = \pm i \left| \hat{h}_{1,2} + \sum_{j=3}^n h_{2,j} \right| \quad (3.116)$$

and, by virtue of being antisymmetric, is unitarily diagonalizable (over the complex numbers) with matrix  $Q$ . After again replacing  $f$  by  $\hat{f}$ , we see that the only entry of  $\hat{g}\hat{\nabla}^2\hat{h}$  that depends on  $c$  is  $\hat{h}_{1,2}$ , which only appears in the antisymmetric second part. We now apply Gerschgorin's theorem to  $Q\hat{g}\hat{\nabla}^2\hat{h}Q^*$ .

The absolute values of all entries of  $QA_{\text{Symm}}Q^*$  can be bounded by  $n^2\|A_{\text{Symm}}\|_\infty$ , independently of  $c$ .  $QA_{\text{Skew}}Q^*$  on the other hand will be diagonal and have as only non-zero entries two purely imaginary diagonal entries, whose absolute values will *grow* asymptotically linearly with  $c$ . Hence, the radii of all Gerschgorin disks can be bounded independently of  $c$ , as can the absolute values of all but



two centers of the Gerschgorin disks. For the centers whose absolute values can not be bounded independently of  $c$ , we can bound the distance to  $\lambda_{1,2}$  independently of  $c$ . However, as we can make  $|\lambda_{1,2}|$  arbitrarily large, this means that we can choose  $c$  such that two Gerschgorin disks are isolated from all others and do not touch the real line. This is sufficient to conclude that, at the origin,  $\hat{g}\hat{\nabla}^2\hat{h}$  has at least one eigenvalue pair with non-zero imaginary part, and hence the induced PDE is non-hyperbolic at the origin and in a neighborhood around the origin. This concludes the proof.  $\square$

**Remark.**     • *For  $n = 1$  every first order quasilinear PDE is hyperbolic and Poisson.*

- *For constant Hamiltonians the evolution equation is trivially zero.*
- *For  $n > 2$ , the constructed metric need not be Lorentzian, but it simplifies the presentation.*
- *If  $h_2(0) = 0$  we simply permute and shift the coordinate system.*

#### 3.4.2.2 in multiple spatial dimensions

#### 3.4.3 with degenerate metric

##### 3.4.3.1 in one spatial dimension

##### 3.4.3.2 in multiple spatial dimensions

#### 3.4.4 of Lie-Poisson type

##### 3.4.4.1 in one spatial dimension

##### 3.4.4.2 in multiple spatial dimensions

### 3.5 Hyperbolic gradient flows

## Chapter 4

# Structure preserving integrators

## 4.1 Symplectic integrators

## 4.2 Energy preserving integrators

### 4.3 Poisson integrators

## 4.4 Multi-symplectic integrators

## 4.5 Dissipation preserving integrators



## 4.6 Conformal symplectic integrators

## 4.7 Metriplectic integrators

## 4.8 Finite volume methods

## 4.9 Symmetric integrators

4.9.1 The discontinuous Galerkin time stepping method

4.9.2 The adjoint of the discontinuous Galerkin time stepping method

4.9.3 The symmetric discontinuous Galerkin time stepping method

4.9.4 The symmetric-dissipative discontinuous Galerkin time stepping method

## Chapter 5

## Conclusion

# Bibliography

- [AD10] Stephen C. Anco and Amanullah Dar. Conservation laws of inviscid non-isentropic compressible fluid flow in  $n > 1$  spatial dimensions. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 466(2121):2605–2632, mar 2010. URL: <https://doi.org/10.1098/rspa.2009.0579>, doi:10.1098/rspa.2009.0579.
- [AK98] Vladimir I. Arnold and Boris A. Khesin. *Topological Methods in Hydrodynamics*. Springer New York, 1998. URL: <https://doi.org/10.1007/2Fb97593>, doi:10.1007/b97593.
- [Ale98] Losik Michor Alekseevsky, Kriegl. Choosing roots of polynomials smoothly. 1998. URL: <http://arxiv.org/abs/math/9801026>.
- [and07] Boris Kolev and. Poisson brackets in hydrodynamics. *Discrete & Continuous Dynamical Systems - A*, 19(3):555–574, 2007. URL: <https://doi.org/10.3934/2Fdcds.2007.19.555>, doi:10.3934/dcds.2007.19.555.
- [ANN<sup>+</sup>89] Metin Arik, Fahrnis Neyzi, Yavuz Nutku, Peter J. Olver, and John M. Verosky. Multi-hamiltonian structure of the born–infeld equation. *Journal of Mathematical Physics*, 30(6):1338–1344, jun 1989. URL: <https://doi.org/10.1063/2F1.528314>, doi:10.1063/1.528314.
- [Arn66] Vladimir Arnold. Sur la géométrie différentielle des groupes de lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits. *Annales de l'institut Fourier*, 16(1):319–361, 1966. URL: <https://doi.org/10.5802/2Faif.233>, doi:10.5802/aif.233.
- [Arn89] V. I. Arnold. *Mathematical Methods of Classical Mechanics*. Springer New York, 1989. URL: <https://doi.org/10.1007/2F978-1-4757-2063-1>, doi:10.1007/978-1-4757-2063-1.
- [BdG12] Dietrich Burde and Willem de Graaf. Classification of novikov algebras. *Applicable Algebra in Engineering, Communication and Computing*, 24(1):1–15, nov 2012. URL: <https://doi.org/10.1007/2Fs00200-012-0180-x>, doi:10.1007/s00200-012-0180-x.

- [Ben87] Discontinuous solutions of one-dimensional hamiltonian systems. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 413(1845):263–295, oct 1987. URL: <https://doi.org/10.1098%2Frspa.1987.0115>, doi:10.1098/rspa.1987.0115.
- [BGS06] Sylvie Benzoni-Gavage and Denis Serre. *Multi-dimensional hyperbolic partial differential equations*. Oxford University Press, nov 2006. URL: <https://doi.org/10.1093%2Facprof%3Aoso%2F9780199211234.001.0001>, doi:10.1093/acprof:oso/9780199211234.001.0001.
- [Blu83] George W. Bluman. On mapping linear partial differential equations to constant coefficient equations. *SIAM Journal on Applied Mathematics*, 43(6):1259–1273, dec 1983. URL: <https://doi.org/10.1137%2F0143084>, doi:10.1137/0143084.
- [BM01a] Chengming Bai and Daoji Meng. The classification of novikov algebras in low dimensions. *Journal of Physics A: Mathematical and General*, 34(8):1581–1594, feb 2001. URL: <https://doi.org/10.1088%2F0305-4470%2F34%2F8%2F305>, doi:10.1088/0305-4470/34/8/305.
- [BM01b] Chengming Bai and Daoji Meng. The classification of novikov algebras in low dimensions: invariant bilinear forms. *Journal of Physics A: Mathematical and General*, 34(39):8193–8197, sep 2001. URL: <https://doi.org/10.1088%2F0305-4470%2F34%2F39%2F401>, doi:10.1088/0305-4470/34/39/401.
- [BN85] A. A. Balinskii and S. P. Novikov. Poisson brackets of hydrodynamic type, frobenius algebras and lie algebras. 1985. URL: <http://mi.mathnet.ru/eng/dan47130>.
- [Bog96] Oleg I. Bogoyavlenskij. Necessary conditions for existence of non-degenerate hamiltonian structures. *Communications in Mathematical Physics*, 182(2):253–289, dec 1996. URL: <https://doi.org/10.1007%2Fbf02517890>, doi:10.1007/bf02517890.
- [Bog99] Oleg I Bogoyavlenskij. Courant’s problems and their extensions. In *Hyperbolic problems: theory, numerics, applications*, pages 97–104. Springer, 1999.
- [Bog06a] Oleg I. Bogoyavlenskij. Algebraic identities for the nijmegen tensors. *Differential Geometry and its Applications*, 24(5):447–457, sep 2006. URL: <https://doi.org/10.1016%2Fj.difgeo.2006.02.009>, doi:10.1016/j.difgeo.2006.02.009.
- [Bog06b] Oleg I. Bogoyavlenskij. Schouten tensor and bi-hamiltonian systems of hydrodynamic type. *Journal of Mathematical Physics*,

- 47(2):023504, feb 2006. URL: <https://doi.org/10.1063%2F1.2167920>, doi:10.1063/1.2167920.
- [Bog07a] Oleg I. Bogoyavlenskij. Invariant dynamical system for a poisson bracket of hydrodynamic type. *Physics Letters A*, 365(1-2):105–110, may 2007. URL: <https://doi.org/10.1016%2Fj.physleta.2006.12.046>, doi:10.1016/j.physleta.2006.12.046.
- [Bog07b] Oleg I. Bogoyavlenskij. Invariant foliations for the poisson brackets of hydrodynamic type. *Physics Letters A*, 360(4-5):539–544, jan 2007. URL: <https://doi.org/10.1016%2Fj.physleta.2006.08.061>, doi:10.1016/j.physleta.2006.08.061.
- [Bog07c] Oleg I. Bogoyavlenskij. Tensor invariants of the poisson brackets of hydrodynamic type. *Communications in Mathematical Physics*, 277(2):369–384, oct 2007. URL: <https://doi.org/10.1007%2Fs00220-007-0370-8>, doi:10.1007/s00220-007-0370-8.
- [Bog11] Oleg I Bogoyavlenskij. Necessary conditions for the existence of conservation laws for systems of partial differential equations. *Mathematics and Theoretical Physics*, page 359, 2011.
- [Bou95] Nicolas Bourbaki. *General Topology*. Springer Berlin Heidelberg, 1995. URL: <https://doi.org/10.1007%2F978-3-642-61701-0>, doi:10.1007/978-3-642-61701-0.
- [BP01] A V Balandin and G V Potemin. On non-degenerate differential-geometric poisson brackets of third order. *Russian Mathematical Surveys*, 56(5):976–977, oct 2001. URL: <https://doi.org/10.1070%2Frm2001v056n05abeh000441>, doi:10.1070/rm2001v056n05abeh000441.
- [BPZ14] Thomas Bridges, Jonathan Pennant, and Sergey Zelik. Degenerate hyperbolic conservation laws with dissipation: Reduction to and validity of a class of burgers-type equations. *Archive for Rational Mechanics and Analysis*, 214(2):671–716, jul 2014. URL: <https://doi.org/10.1007%2Fs00205-014-0772-7>, doi:10.1007/s00205-014-0772-7.
- [BR01] Thomas J. Bridges and Sebastian Reich. Multi-symplectic integrators: numerical schemes for hamiltonian PDEs that conserve symplecticity. *Physics Letters A*, 284(4-5):184–193, jun 2001. URL: <https://doi.org/10.1016%2Fs0375-9601%2801%2900294-8>, doi:10.1016/s0375-9601(01)00294-8.
- [BR08] O. Bogoyavlenskij and P. Reynolds. Form-invariant poisson brackets of hydrodynamic type with several spatial variables. *Journal of Mathematical Physics*, 49(5):053520, may 2008. URL: <https://doi.org/10.1063%2F1.2924215>, doi:10.1063/1.2924215.



- [BR10] O. I. Bogoyavlenskij and A. P. Reynolds. Criteria for existence of a hamiltonian structure. *Regular and Chaotic Dynamics*, 15(4-5):431–439, aug 2010. URL: <https://doi.org/10.1134/S1560354710040039>, doi:10.1134/s1560354710040039.
- [BR11] D. A. Berdinsky and I. P. Rybnikov. On orthogonal curvilinear coordinate systems in constant curvature spaces. *Siberian Mathematical Journal*, 52(3):394–401, may 2011. URL: <https://doi.org/10.1134/S0037446611030025>, doi:10.1134/s0037446611030025.
- [Bri66] David R. Brillinger. The analyticity of the roots of a polynomial as functions of the coefficients. *Mathematics Magazine*, 39(3):145, may 1966. URL: <https://doi.org/10.2307/2F2689304>, doi:10.2307/2689304.
- [Bue96] Peter Bueken. Multi-hamiltonian structures for a class of degenerate completely integrable systems. *Journal of Mathematical Physics*, 37(6):2851–2862, jun 1996. URL: <https://doi.org/10.1063/1.531543>, doi:10.1063/1.531543.
- [Cal17] Gianluca Calcagni. *Classical and Quantum Cosmology*. Springer International Publishing, 2017. URL: <https://doi.org/10.1007/978-3-319-41127-9>, doi:10.1007/978-3-319-41127-9.
- [Car93] P. Carbonaro. Exceptional  $2 \times 2$  quasilinear hyperbolic systems. *Physics Letters A*, 182(4-6):363–368, nov 1993. URL: [https://doi.org/10.1016/0375-9601\(93\)90409-s](https://doi.org/10.1016/0375-9601(93)90409-s), doi:10.1016/0375-9601(93)90409-s.
- [Car95] P. Carbonaro. Exceptional hyperbolic systems of hamiltonian form. *European Journal of Applied Mathematics*, 6(2):157–167, apr 1995. URL: <https://doi.org/10.1017/S0956792500001753>, doi:10.1017/s0956792500001753.
- [Cas14] Matteo Casati. On deformations of multidimensional poisson brackets of hydrodynamic type. *Communications in Mathematical Physics*, 335(2):851–894, nov 2014. URL: <https://doi.org/10.1007/s00220-014-2219-2>, doi:10.1007/s00220-014-2219-2.
- [CC16] C. E. Caligan and C. Chandre. Conservative dissipation: How important is the jacobi identity in the dynamics? *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 26(5):053101, may 2016. URL: <https://doi.org/10.1063/1.4948411>, doi:10.1063/1.4948411.
- [CCS17] Guido Carlet, Matteo Casati, and Sergey Shadrin. Poisson cohomology of scalar multidimensional dubrovin–novikov brackets. *Journal of Geometry and Physics*, 114:404–419, apr 2017. URL: <https://doi.org/10.1016/j.geomphys.2016.12.008>, doi:10.1016/j.geomphys.2016.12.008.

- [CH89] R. Courant and D. Hilbert. *Methods of Mathematical Physics*. Wiley, apr 1989. URL: <https://doi.org/10.1002/2F9783527617210>, doi:10.1002/9783527617210.
- [CM82] J. Cavalcante and H.P. McKean. The classical shallow water equations: Symplectic geometry. *Physica D: Nonlinear Phenomena*, 4(2):253–260, jan 1982. URL: [https://doi.org/10.1016/0167-2789\(82\)90066-5](https://doi.org/10.1016/0167-2789(82)90066-5), doi:10.1016/0167-2789(82)90066-5.
- [CM15] Felipe Contatto and Dunajski Maciej. First integrals of affine connections and hamiltonian systems of hydrodynamic type. *Journal of Integrable Systems*, 1(1):xyw009, sep 2015. URL: <https://doi.org/10.1093/integr/xyw009>, doi:10.1093/integr/xyw009.
- [Coo91a] David B. Cooke. Classification results and the darboux theorem for low-order hamiltonian operators. *Journal of Mathematical Physics*, 32(1):109–119, jan 1991. URL: <https://doi.org/10.1063/1.529133>, doi:10.1063/1.529133.
- [Coo91b] David B. Cooke. Compatibility conditions for hamiltonian pairs. *Journal of Mathematical Physics*, 32(11):3071–3076, nov 1991. URL: <https://doi.org/10.1063/1.529053>, doi:10.1063/1.529053.
- [Daf87] Constantine M. Dafermos. Estimates for conservation laws with little viscosity. *SIAM Journal on Mathematical Analysis*, 18(2):409–421, mar 1987. URL: <https://doi.org/10.1137/0518031>, doi:10.1137/0518031.
- [Daf10] Constantine M. Dafermos. *Hyperbolic Conservation Laws in Continuum Physics*. Springer Berlin Heidelberg, 2010. URL: <https://doi.org/10.1007/978-3-642-04048-1>, doi:10.1007/978-3-642-04048-1.
- [DiP78] R.J. DiPerna. Entropy and the uniqueness of solutions to hyperbolic conservation laws. In *Nonlinear Evolution Equations*, pages 1–15. Elsevier, 1978. URL: [https://doi.org/10.1016/0167-2789\(82\)90066-5](https://doi.org/10.1016/0167-2789(82)90066-5), doi:10.1016/0167-2789(82)90066-5.
- [DKN01] B. A. Dubrovin, I. M. Krichever, and S. P. Novikov. Integrable systems.i. In *Dynamical Systems IV*, pages 177–332. Springer Berlin Heidelberg, 2001. URL: [https://doi.org/10.1007/978-3-662-06791-8\\_3](https://doi.org/10.1007/978-3-662-06791-8_3), doi:10.1007/978-3-662-06791-8\_3.
- [DLZ06a] Boris Dubrovin, Si-Qi Liu, and Youjin Zhang. On hamiltonian perturbations of hyperbolic systems of conservation laws i: Quasitriviality of bi-hamiltonian perturbations. *Communications on Pure*

- and *Applied Mathematics*, 59(4):559–615, 2006. URL: <https://doi.org/10.1002%2Fcpa.20111>, doi:10.1002/cpa.20111.
- [DLZ06b] Boris Dubrovin, Si-Qi Liu, and Youjin Zhang. On hamiltonian perturbations of hyperbolic systems of conservation laws i: Quasitriviality of bi-hamiltonian perturbations. *Communications on Pure and Applied Mathematics*, 59(4):559–615, 2006. URL: <https://doi.org/10.1002%2Fcpa.20111>, doi:10.1002/cpa.20111.
- [DM13] Mamadou Diagne and Bernhard Maschke. Port hamiltonian formulation of a system of two conservation laws with a moving interface. *European Journal of Control*, 19(6):495–504, dec 2013. URL: <https://doi.org/10.1016%2Fj.ejcon.2013.09.001>, doi:10.1016/j.ejcon.2013.09.001.
- [DMS04] Luca Degiovanni, Franco Magri, and Vincenzo Sciacca. On deformation of poisson manifolds of hydrodynamic type. *Communications in Mathematical Physics*, 253(1):1–24, nov 2004. URL: <https://doi.org/10.1007%2Fs00220-004-1190-8>, doi:10.1007/s00220-004-1190-8.
- [DN83a] B. A. Dubrovin and S. P. Novikov. The hamiltonian formalism of one-dimensional systems of hydrodynamic type and the bogolyubov-whitham averaging method. 1983. URL: <https://pdfs.semanticscholar.org/5103/72fc0865f2f2c2e8f7b50e87daca444a242d.pdf>.
- [DN83b] B. A. Dubrovin and S. P. Novikov. On poisson brackets of hydrodynamic type. 1983. URL: [https://iris.sissa.it/retrieve/handle/20.500.11767/80242/79796/dubrovin\\_novikov\\_1983\\_dan.pdf](https://iris.sissa.it/retrieve/handle/20.500.11767/80242/79796/dubrovin_novikov_1983_dan.pdf).
- [DN89] B A Dubrovin and S P Novikov. Hydrodynamics of weakly deformed soliton lattices. differential geometry and hamiltonian theory. *Russian Mathematical Surveys*, 44(6):35–124, dec 1989. URL: <https://doi.org/10.1070%2Frm1989v044n06abeh002300>, doi:10.1070/rm1989v044n06abeh002300.
- [DN96] B. A. DUBROVIN and S. P. NOVIKOV. HAMILTONIAN FORMALISM OF ONE-DIMENSIONAL SYSTEMS OF HYDRODYNAMIC TYPE, AND THE BOGOLYUBOV-WHITMAN AVERAGING METHOD. In *30 Years of the Landau Institute — Selected Papers*, pages 382–386. WORLD SCIENTIFIC, aug 1996. URL: [https://doi.org/10.1142%2F9789814317344\\_0051](https://doi.org/10.1142%2F9789814317344_0051), doi:10.1142/9789814317344\_0051.
- [Doy93a] Philip W. Doyle. Differential geometric poisson bivectors in one space variable. *Journal of Mathematical Physics*, 34(4):1314–1338,

- apr 1993. URL: <https://doi.org/10.1063%2F1.530213>, doi:10.1063/1.530213.
- [Doy93b] Philip W. Doyle. Differential geometric poisson bivectors in one space variable. *Journal of Mathematical Physics*, 34(4):1314–1338, apr 1993. URL: <https://doi.org/10.1063%2F1.530213>, doi:10.1063/1.530213.
- [Doy94] Philip W. Doyle. Symmetry classes of quasilinear systems in one space variable. *Journal of Nonlinear Mathematical Physics*, 1(3):225–266, jan 1994. URL: <https://doi.org/10.2991%2Fjnmp.1994.1.3.1>, doi:10.2991/jnmp.1994.1.3.1.
- [DPRZ17] Michael Dumbser, Ilya Peshkov, Evgeniy Romenski, and Olindo Zanotti. High order ADER schemes for a unified first order hyperbolic formulation of newtonian continuum mechanics coupled with electro-dynamics. *Journal of Computational Physics*, 348:298–342, nov 2017. URL: <https://doi.org/10.1016%2Fj.jcp.2017.07.020>, doi:10.1016/j.jcp.2017.07.020.
- [Dub01] Zhang Dubrovin. Normal forms of hierarchies of integrable pdes, frobenius manifolds and gromov - witten invariants. 2001. URL: <http://arxiv.org/abs/math/0108160>.
- [DV80] I.E. Dzyaloshinskii and G.E. Volovick. Poisson brackets in condensed matter physics. *Annals of Physics*, 125(1):67–97, mar 1980. URL: <https://doi.org/10.1016%2F0003-4916%2880%2990119-0>, doi:10.1016/0003-4916(80)90119-0.
- [Eis50] Luther Pfahler Eisenhart. *Riemannian Geometry*. Princeton University Press, dec 1950. URL: <https://doi.org/10.1515%2F9781400884216>, doi:10.1515/9781400884216.
- [Eva10] Lawrence Evans. *Partial Differential Equations*. American Mathematical Society, mar 2010. URL: <https://doi.org/10.1090%2Fgsm%2F019>, doi:10.1090/gsm/019.
- [Fer01] E V Ferapontov. Compatible poisson brackets of hydrodynamic type. *Journal of Physics A: Mathematical and General*, 34(11):2377–2388, mar 2001. URL: <https://doi.org/10.1088%2F0305-4470%2F34%2F11%2F328>, doi:10.1088/0305-4470/34/11/328.
- [FL71] K. O. Friedrichs and P. D. Lax. Systems of conservation equations with a convex extension. *Proceedings of the National Academy of Sciences*, 68(8):1686–1688, aug 1971. URL: <https://doi.org/10.1073%2Fpnas.68.8.1686>, doi:10.1073/pnas.68.8.1686.

- [FLS14] Evgeny V. Ferapontov, Paolo Lorenzoni, and Andrea Savoldi. Hamiltonian operators of dubrovin-novikov type in 2d. *Letters in Mathematical Physics*, 105(3):341–377, dec 2014. URL: <https://doi.org/10.1007%2Fs11005-014-0738-6>, doi:10.1007/s11005-014-0738-6.
- [FMS08] E. V. Ferapontov, A. Moro, and V. V. Sokolov. Hamiltonian systems of hydrodynamic type in  $2 + 1$  dimensions. *Communications in Mathematical Physics*, 285(1):31–65, jun 2008. URL: <https://doi.org/10.1007%2Fs00220-008-0522-5>, doi:10.1007/s00220-008-0522-5.
- [Gar71] Clifford S. Gardner. Korteweg-de vries equation and generalizations. IV. the korteweg-de vries equation as a hamiltonian system. *Journal of Mathematical Physics*, 12(8):1548–1551, aug 1971. URL: <https://doi.org/10.1063%2F1.1665772>, doi:10.1063/1.1665772.
- [GBB17] Michael Groß, Matthias Bartelt, and Peter Betsch. Structure-preserving time integration of non-isothermal finite viscoelastic continua related to variational formulations of continuum dynamics. *Computational Mechanics*, 62(2):123–150, oct 2017. URL: <https://doi.org/10.1007%2Fs00466-017-1489-x>, doi:10.1007/s00466-017-1489-x.
- [GD80a] I. M. Gel'fand and I. Ya. Dorfman. Hamiltonian operators and algebraic structures related to them. *Functional Analysis and Its Applications*, 13(4):248–262, 1980. URL: <https://doi.org/10.1007%2Fbf01078363>, doi:10.1007/bf01078363.
- [GD80b] I. M. Gel'fand and I. Ya. Dorfman. The schouten bracket and hamiltonian operators. *Functional Analysis and Its Applications*, 14(3):223–226, jul 1980. URL: <https://doi.org/10.1007%2Fbf01086188>, doi:10.1007/bf01086188.
- [GD82] I. M. Gel'fand and I. Ya. Dorfman. Hamiltonian operators and infinite-dimensional lie algebras. *Functional Analysis and Its Applications*, 15(3):173–187, 1982. URL: <https://doi.org/10.1007%2Fbf01089922>, doi:10.1007/bf01089922.
- [GGZ09] Metin Gurses, Gusein Sh. Guseinov, and Kostyantyn Zheltukhin. Dynamical systems and poisson structures. *Journal of Mathematical Physics*, 50(11):112703, nov 2009. URL: <https://doi.org/10.1063%2F1.3257919>, doi:10.1063/1.3257919.
- [Gli65] James Glimm. Solutions in the large for nonlinear hyperbolic systems of equations. *Communications on Pure and Applied Mathematics*, 18(4):697–715, nov 1965. URL: <https://doi.org/10.1002%2Fcpa.3160180408>, doi:10.1002/cpa.3160180408.

- [GM19] E. V. Glukhov and O. I. Mokhov. On algebraic-geometry methods for constructing flat diagonal metrics of a special form. *Russian Mathematical Surveys*, 74(4):761–763, aug 2019. URL: <https://doi.org/10.1070%2Frm9891>, doi:10.1070/rm9891.
- [GMR96] S. K. Godunov, T. Yu. Mikhaïlova, and E. I. Romenskiĭ. Systems of thermodynamically coordinated laws of conservation invariant under rotations. *Siberian Mathematical Journal*, 37(4):690–705, jul 1996. URL: <https://doi.org/10.1007%2Fbf02104662>, doi:10.1007/bf02104662.
- [GN90] H. Gumral and Y. Nutku. Multi-hamiltonian structure of equations of hydrodynamic type. *Journal of Mathematical Physics*, 31(11):2606–2611, nov 1990. URL: <https://doi.org/10.1063%2F1.529012>, doi:10.1063/1.529012.
- [GN94] H Gumral and Y Nutku. Bi-hamiltonian structures of d-boussinesq and benney-lax equations. *Journal of Physics A: Mathematical and General*, 27(1):193–200, jan 1994. URL: <https://doi.org/10.1088%2F0305-4470%2F27%2F1%2F013>, doi:10.1088/0305-4470/27/1/013.
- [GO97] Miroslav Grmela and Hans Christian Ottinger. Dynamics and thermodynamics of complex fluids. i. development of a general formalism. *Physical Review E*, 56(6):6620–6632, dec 1997. URL: <https://doi.org/10.1103%2Fphysreve.56.6620>, doi:10.1103/physreve.56.6620.
- [God61] S. K. Godunov. An interesting class of quasi-linear systems. 1961.
- [God62] S K Godunov. THE PROBLEM OF a GENERALIZED SOLUTION IN THE THEORY OF QUASILINEAR EQUATIONS AND IN GAS DYNAMICS. *Russian Mathematical Surveys*, 17(3):145–156, jun 1962. URL: <https://doi.org/10.1070%2Frm1962v017n03abeh004116>, doi:10.1070/rm1962v017n03abeh004116.
- [Gri85] N I Grinberg. On poisson brackets of hydrodynamic type with a degenerate metric. *Russian Mathematical Surveys*, 40(4):231–232, aug 1985. URL: <https://doi.org/10.1070%2Frm1985v040n04abeh003662>, doi:10.1070/rm1985v040n04abeh003662.
- [GSW00] A M Grundland, M B Sheftel, and P Winternitz. Invariant solutions of hydrodynamic-type equations. *Journal of Physics A: Mathematical and General*, 33(46):8193–8215, nov 2000. URL: <https://doi.org/10.1088%2F0305-4470%2F33%2F46%2F304>, doi:10.1088/0305-4470/33/46/304.

- [GTLF18] Binfang Gao, Kai Tian, Q. P. Liu, and Lujuan Feng. Conservation laws of the generalized riemann equations. *Journal of Nonlinear Mathematical Physics*, 25(1):122–135, jan 2018. URL: <https://doi.org/10.1080%2F14029251.2018.1440746>, doi:10.1080/14029251.2018.1440746.
- [Haa11] F. Haas. Comment on “dynamical systems and poisson structures” [j. math. phys. 50, 112703 (2009)]. *Journal of Mathematical Physics*, 52(12):124101, dec 2011. URL: <https://doi.org/10.1063%2F1.3664755>, doi:10.1063/1.3664755.
- [HMRW85] Darryl D. Holm, Jerrold E. Marsden, Tudor Ratiu, and Alan Weinstein. Nonlinear stability of fluid and plasma equilibria. *Physics Reports*, 123(1-2):1–116, jul 1985. URL: <https://doi.org/10.1016%2F0370-1573%2885%2990028-6>, doi:10.1016/0370-1573(85)90028-6.
- [Hog13] Leslie Hogben, editor. *Handbook of Linear Algebra*. Chapman and Hall/CRC, nov 2013. URL: <https://doi.org/10.1201%2Fb16113>, doi:10.1201/b16113.
- [Hun06] J.C.R. Hunt. NONLINEAR AND WAVE THEORY CONTRIBUTIONS OF t. BROOKE BENJAMIN (1929–1995). *Annual Review of Fluid Mechanics*, 38(1):1–25, jan 2006. URL: <https://doi.org/10.1146%2Fannurev.fluid.38.050304.092028>, doi:10.1146/annurev.fluid.38.050304.092028.
- [Hun07] John D. Hunter. Matplotlib: A 2d graphics environment. *Computing in Science & Engineering*, 9(3):90–95, 2007. URL: <https://doi.org/10.1109%2Fmcse.2007.55>, doi:10.1109/mcse.2007.55.
- [Kac90] Victor G. Kac. *Infinite-Dimensional Lie Algebras*. Cambridge University Press, sep 1990. URL: <https://doi.org/10.1017%2Fcbo9780511626234>, doi:10.1017/cbo9780511626234.
- [Kah80] Chapter 7 infinite-dimensional manifolds. In *Introduction to Global Analysis*, pages 206–225. Elsevier, 1980. URL: <https://doi.org/10.1016%2Fs0079-8169%2808%2961124-1>, doi:10.1016/s0079-8169(08)61124-1.
- [Kat75] Tosio Kato. The cauchy problem for quasi-linear symmetric hyperbolic systems. *Archive for Rational Mechanics and Analysis*, 58(3):181–205, 1975. URL: <https://doi.org/10.1007%2Fbf00280740>, doi:10.1007/bf00280740.
- [Kat95] Tosio Kato. *Perturbation Theory for Linear Operators*. Springer Berlin Heidelberg, 1995. URL: <https://doi.org/10.1007%2F978-3-642-66282-9>, doi:10.1007/978-3-642-66282-9.

- [KH10] Martin Kroger and Markus Hutter. Automated symbolic calculations in nonequilibrium thermodynamics. *Computer Physics Communications*, 181(12):2149–2157, dec 2010. URL: <https://doi.org/10.1016%2Fj.cpc.2010.07.050>, doi:10.1016/j.cpc.2010.07.050.
- [KHO01] Martin Kroger, Markus Hutter, and Hans Christian Ottinger. Symbolic test of the jacobi identity for given generalized ‘poisson’ bracket. *Computer Physics Communications*, 137(2):325–340, jun 2001. URL: <https://doi.org/10.1016%2Fs0010-4655%2801%2900161-8>, doi:10.1016/s0010-4655(01)00161-8.
- [KM97] Andreas Kriegl and Peter Michor. *The Convenient Setting of Global Analysis*. American Mathematical Society, sep 1997. URL: <https://doi.org/10.1090%2Fsurv%2F053>, doi:10.1090/surv/053.
- [Kri02] Michor Kriegl, Losik. Choosing roots of polynomials smoothly, ii. 2002. URL: <http://arxiv.org/abs/math/0208228>.
- [KVV17] Joseph Krasil'shchik, Alexander Verbovetsky, and Raffaele Vito. *The Symbolic Computation of Integrability Structures for Partial Differential Equations*. Springer International Publishing, 2017. URL: <https://doi.org/10.1007%2F978-3-319-71655-8>, doi:10.1007/978-3-319-71655-8.
- [KW09] Boris Khesin and Robert Wendt. *The Geometry of Infinite-Dimensional Groups*. Springer Berlin Heidelberg, 2009. URL: <https://doi.org/10.1007%2F978-3-540-77263-7>, doi:10.1007/978-3-540-77263-7.
- [Lan95] Serge Lang, editor. *Differential and Riemannian Manifolds*. Springer New York, 1995. URL: <https://doi.org/10.1007%2F978-1-4612-4182-9>, doi:10.1007/978-1-4612-4182-9.
- [Lan02] *Introduction to Differential Manifolds*. Springer-Verlag, 2002. URL: <https://doi.org/10.1007%2Fb97450>, doi:10.1007/b97450.
- [Lax53] Peter D. Lax. Nonlinear hyperbolic equations. *Communications on Pure and Applied Mathematics*, 6(2):231–258, may 1953. URL: <https://doi.org/10.1002%2Fcpa.3160060204>, doi:10.1002/cpa.3160060204.
- [Lee11] John M. Lee. *Introduction to Topological Manifolds*. Springer New York, 2011. URL: <https://doi.org/10.1007%2F978-1-4419-7940-7>, doi:10.1007/978-1-4419-7940-7.
- [Lee12] John M. Lee. *Introduction to Smooth Manifolds*. Springer New York, 2012. URL: <https://doi.org/10.1007%2F978-1-4419-9982-5>, doi:10.1007/978-1-4419-9982-5.



- [Lee18] John M. Lee. *Introduction to Riemannian Manifolds*. Springer International Publishing, 2018. URL: <https://doi.org/10.1007/2F978-3-319-91755-9>, doi:10.1007/978-3-319-91755-9.
- [LGPV13] Camille Laurent-Gengoux, Anne Pichereau, and Pol Vanhaecke. *Poisson Structures*. Springer Berlin Heidelberg, 2013. URL: <https://doi.org/10.1007/2F978-3-642-31090-4>, doi:10.1007/978-3-642-31090-4.
- [LO00] Yi A Li and Peter J Olver. Well-posedness and blow-up solutions for an integrable nonlinearly dispersive model wave equation. *Journal of Differential Equations*, 162(1):27–63, mar 2000. URL: <https://doi.org/10.1006/2Fjdeq.1999.3683>, doi:10.1006/jdeq.1999.3683.
- [LZ11] Si-Qi Liu and Youjin Zhang. Jacobi structures of evolutionary partial differential equations. *Advances in Mathematics*, 227(1):73–130, may 2011. URL: <https://doi.org/10.1016/2Fj.aim.2011.01.015>, doi:10.1016/j.aim.2011.01.015.
- [Mag78] Franco Magri. A simple model of the integrable hamiltonian equation. *Journal of Mathematical Physics*, 19(5):1156–1162, may 1978. URL: <https://doi.org/10.1063/2F1.523777>, doi:10.1063/1.523777.
- [Maj84] A. Majda. *Compressible Fluid Flow and Systems of Conservation Laws in Several Space Variables*. Springer New York, 1984. URL: <https://doi.org/10.1007/2F978-1-4612-1116-7>, doi:10.1007/978-1-4612-1116-7.
- [Man79] Yu. I. Manin. Algebraic aspects of nonlinear differential equations. *Journal of Soviet Mathematics*, 11(1):1–122, jan 1979. URL: <https://doi.org/10.1007/2Fbf01084246>, doi:10.1007/bf01084246.
- [MF90] O I Mokhov and E V Ferapontov. Non-local hamiltonian operators of hydrodynamic type related to metrics of constant curvature. *Russian Mathematical Surveys*, 45(3):218–219, jun 1990. URL: <https://doi.org/10.1070/2Frm1990v045n03abeh002351>, doi:10.1070/rm1990v045n03abeh002351.
- [MG80] Philip J. Morrison and John M. Greene. Noncanonical hamiltonian density formulation of hydrodynamics and ideal magnetohydrodynamics. *Physical Review Letters*, 45(10):790–794, sep 1980. URL: <https://doi.org/10.1103/2Fphysrevlett.45.790>, doi:10.1103/physrevlett.45.790.
- [Mok89] O. I. Mokhov. Dubrovin-novikov type poisson brackets (DN-brackets). *Functional Analysis and Its Applications*, 22(4):336–338, 1989. URL: <https://doi.org/10.1007/2Fbf01077434>, doi:10.1007/bf01077434.

- [Mok92] O.I. Mokhov. Hamiltonian systems of hydrodynamic type and constant curvature metrics. *Physics Letters A*, 166(3-4):215–216, jun 1992. URL: <https://doi.org/10.1016%2F0375-9601%2892%2990365-s>, doi:10.1016/0375-9601(92)90365-s.
- [Mok98] O I Mokhov. Symplectic and poisson structures on loop spaces of smooth manifolds, and integrable systems. *Russian Mathematical Surveys*, 53(3):515–622, jun 1998. URL: <https://doi.org/10.1070%2Frm1998v053n03abeh000019>, doi:10.1070/rm1998v053n03abeh000019.
- [Mok00] O I Mokhov. Compatible and almost compatible metrics. *Russian Mathematical Surveys*, 55(4):819–821, aug 2000. URL: <https://doi.org/10.1070%2Frm2000v055n04abeh000318>, doi:10.1070/rm2000v055n04abeh000318.
- [Mok01] O. I. Mokhov. *Functional Analysis and Its Applications*, 35(2):100–110, 2001. URL: <https://doi.org/10.1023%2Fa%3A1017575131419>, doi:10.1023/a:1017575131419.
- [Mok02a] O. I. Mokhov. *Theoretical and Mathematical Physics*, 132(1):942–954, 2002. URL: <https://doi.org/10.1023%2Fa%3A1019659324655>, doi:10.1023/a:1019659324655.
- [Mok02b] O. I. Mokhov. *Theoretical and Mathematical Physics*, 133(2):1557–1564, 2002. URL: <https://doi.org/10.1023%2Fa%3A1021155028895>, doi:10.1023/a:1021155028895.
- [Mok02c] Oleg I. Mokhov. Compatible flat metrics. *Journal of Applied Mathematics*, 2(7):337–370, 2002. URL: <https://doi.org/10.1155%2Fs1110757x02203149>, doi:10.1155/s1110757x02203149.
- [Mok06] O I Mokhov. The classification of multidimensional poisson brackets of hydrodynamic type. *Russian Mathematical Surveys*, 61(2):356–358, apr 2006. URL: <https://doi.org/10.1070%2Frm2006v061n02abeh004319>, doi:10.1070/rm2006v061n02abeh004319.
- [Mok08] O. I. Mokhov. The classification of nonsingular multidimensional dubrovin-novikov brackets. *Functional Analysis and Its Applications*, 42(1):33–44, jan 2008. URL: <https://doi.org/10.1007%2Fs10688-008-0004-8>, doi:10.1007/s10688-008-0004-8.
- [Mok12] Oleg I. Mokhov. Multidimensional poisson brackets of hydrodynamic type and flat pencils of metrics. Technical report, 2012. URL: <https://doi.org/10.7546%2Fjgsp-10-2007-51-72>, doi:10.7546/jgsp-10-2007-51-72.

[Mok17] O I Mokhov. Pencils of compatible metrics and integrable systems. *Russian Mathematical Surveys*, 72(5):889–937, oct 2017. URL: <https://doi.org/10.1070%2Frm9792>, doi:10.1070/rm9792.

[Mor80] Philip J. Morrison. The maxwell-vlasov equations as a continuous hamiltonian system. *Physics Letters A*, 80(5-6):383–386, dec 1980. URL: <https://doi.org/10.1016%2F0375-9601%2880%2990776-8>, doi:10.1016/0375-9601(80)90776-8.

[Mor84] Philip J. Morrison. Bracket formulation for irreversible classical fields. *Physics Letters A*, 100(8):423–427, feb 1984. URL: <https://doi.org/10.1016%2F0375-9601%2884%2990635-2>, doi:10.1016/0375-9601(84)90635-2.

[Mor17] P. J. Morrison. Structure and structure-preserving algorithms for plasma physics. *Physics of Plasmas*, 24(5):055502, apr 2017. URL: <https://doi.org/10.1063%2F1.4982054>, doi:10.1063/1.4982054.

[MP18] O. I. Mokhov and N. A. Pavlenko. Classification of the associativity equations with a first-order hamiltonian operator. *Theoretical and Mathematical Physics*, 197(1):1501–1513, oct 2018. URL: <https://doi.org/10.1134%2Fs0040577918100070>, doi:10.1134/s0040577918100070.

[MR98] Ingo Muller and Tomasso Ruggeri. *Rational Extended Thermodynamics*. Springer New York, 1998. URL: <https://doi.org/10.1007%2F978-1-4612-2210-1>, doi:10.1007/978-1-4612-2210-1.

[MSP<sup>+</sup>17] Aaron Meurer, Christopher P. Smith, Mateusz Paprocki, Ondřej Čertík, Sergey B. Kirpichev, Matthew Rocklin, AMiT Kumar, Sergiu Ivanov, Jason K. Moore, Sartaj Singh, Thilina Rathnayake, Sean Vig, Brian E. Granger, Richard P. Muller, Francesco Bonazzi, Harsh Gupta, Shivam Vats, Fredrik Johansson, Fabian Pedregosa, Matthew J. Curry, Andy R. Terrel, Štěpán Roučka, Ashutosh Saboo, Isuru Fernando, Sumith Kulal, Robert Cimrman, and Anthony Scopatz. SymPy: symbolic computing in python. *PeerJ Computer Science*, 3:e103, jan 2017. URL: <https://doi.org/10.7717%2Fpeerj-cs.103>, doi:10.7717/peerj-cs.103.

[Ney89] Fahrnunisa Neyzi. Diagonalization and hamiltonian structures of hyperbolic systems. *Journal of Mathematical Physics*, 30(8):1695–1698, aug 1989. URL: <https://doi.org/10.1063%2F1.528255>, doi:10.1063/1.528255.

[Nov82] S P Novikov. The hamiltonian formalism and a many-valued analogue of morse theory. *Russian Mathematical Surveys*, 37(5):1–56, oct 1982. URL: <https://doi.org/10.1070/RMS1982v037n05&lang=en>.

- [//doi.org/10.1070%2Frm1982v037n05abeh004020](https://doi.org/10.1070%2Frm1982v037n05abeh004020), doi:  
10.1070/rm1982v037n05abeh004020.
- [Nov85] S P Novikov. The geometry of conservative systems of hydrodynamic type. the method of averaging for field-theoretical systems. *Russian Mathematical Surveys*, 40(4):85–98, aug 1985. URL: <https://doi.org/10.1070%2Frm1985v040n04abeh003615>, doi:10.1070/rm1985v040n04abeh003615.
- [Nut85] Y. Nutku. On a new class of completely integrable nonlinear wave equations. i. infinitely many conservation laws. *Journal of Mathematical Physics*, 26(6):1237–1242, jun 1985. URL: <https://doi.org/10.1063%2F1.526530>, doi:10.1063/1.526530.
- [Nut87] Y. Nutku. On a new class of completely integrable nonlinear wave equations. II. multi-hamiltonian structure. *Journal of Mathematical Physics*, 28(11):2579–2585, nov 1987. URL: <https://doi.org/10.1063%2F1.527749>, doi:10.1063/1.527749.
- [OG97] Hans Christian Ottinger and Miroslav Grmela. Dynamics and thermodynamics of complex fluids. II. illustrations of a general formalism. *Physical Review E*, 56(6):6633–6655, dec 1997. URL: <https://doi.org/10.1103%2Fphysreve.56.6633>, doi:10.1103/physreve.56.6633.
- [Olv80] Peter J. Olver. On the hamiltonian structure of evolution equations. *Mathematical Proceedings of the Cambridge Philosophical Society*, 88(1):71–88, jul 1980. URL: <https://doi.org/10.1017%2Fs0305004100057364>, doi:10.1017/s0305004100057364.
- [Olv86] Peter J. Olver. *Applications of Lie Groups to Differential Equations*. Springer US, 1986. URL: <https://doi.org/10.1007%2F978-1-4684-0274-2>, doi:10.1007/978-1-4684-0274-2.
- [ON88] Peter J. Olver and Yavuz Nutku. Hamiltonian structures for systems of hyperbolic conservation laws. *Journal of Mathematical Physics*, 29(7):1610–1619, jul 1988. URL: <https://doi.org/10.1063%2F1.527909>, doi:10.1063/1.527909.
- [OST20] Hans Christian Ottinger, Henning Struchtrup, and Manuel Torrilhon. Formulation of moment equations for rarefied gases within two frameworks of non-equilibrium thermodynamics: RET and GENERIC. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378(2170):20190174, mar 2020. URL: <https://doi.org/10.1098%2Frsta.2019.0174>, doi:10.1098/rsta.2019.0174.
- [Ott01] Felix Otto. THE GEOMETRY OF DISSIPATIVE EVOLUTION EQUATIONS: THE POROUS MEDIUM EQUATION. *Communications in Partial Differential Equations*, 26(1-2):101–174, jan

2001. URL: <https://doi.org/10.1081%2Fpde-100002243>, doi: 10.1081/pde-100002243.
- [Ott05] Hans Christian Ottinger. *Beyond Equilibrium Thermodynamics*. John Wiley & Sons, Inc., jan 2005. URL: <https://doi.org/10.1002%2F0471727903>, doi:10.1002/0471727903.
- [P<sup>+</sup>12] Emanuele Parodi et al. Classification problems for hamiltonian evolutionary equations and their discretizations. 2012.
- [Pav03] Maxim V. Pavlov. Integrable hydrodynamic chains. *Journal of Mathematical Physics*, 44(9):4134, 2003. URL: <https://doi.org/10.1063%2F1.1597946>, doi:10.1063/1.1597946.
- [PGR14] Ilya Peshkov, Miroslav Grmela, and Evgeniy Romenski. Irreversible mechanics and thermodynamics of two-phase continua experiencing stress-induced solid–fluid transitions. *Continuum Mechanics and Thermodynamics*, 27(6):905–940, oct 2014. URL: <https://doi.org/10.1007%2Fs00161-014-0386-1>, doi: 10.1007/s00161-014-0386-1.
- [PKEG16] Michal Pavelka, Václav Klika, Oğul Esen, and Miroslav Grmela. A hierarchy of poisson brackets in non-equilibrium thermodynamics. *Physica D: Nonlinear Phenomena*, 335:54–69, nov 2016. URL: <https://doi.org/10.1016%2Fj.physd.2016.06.011>, doi: 10.1016/j.physd.2016.06.011.
- [PKG14] Michal Pavelka, Václav Klika, and Miroslav Grmela. Time reversal in nonequilibrium thermodynamics. *Physical Review E*, 90(6), dec 2014. URL: <https://doi.org/10.1103%2Fphysreve.90.062131>, doi:10.1103/physreve.90.062131.
- [PKG18] Michal Pavelka, Václav Klika, and Miroslav Grmela. *Multiscale Thermo-Dynamics*. De Gruyter, aug 2018. URL: <https://doi.org/10.1515%2F9783110350951>, doi:10.1515/9783110350951.
- [Pol87] M B Polyak. On one-dimensional hamiltonian systems of hydrodynamic type with explicit dependence on the spatial variable. *Russian Mathematical Surveys*, 42(3):229–230, jun 1987. URL: <https://doi.org/10.1070%2Frm1987v042n03abeh001433>, doi:10.1070/rm1987v042n03abeh001433.
- [PPK20] Michal Pavelka, Ilya Peshkov, and Václav Klika. On hamiltonian continuum mechanics. *Physica D: Nonlinear Phenomena*, 408:132510, jul 2020. URL: <https://doi.org/10.1016%2Fj.physd.2020.132510>, doi:10.1016/j.physd.2020.132510.
- [PPRG18] Ilya Peshkov, Michal Pavelka, Evgeniy Romenski, and Miroslav Grmela. Continuum mechanics and thermodynamics in the

- hamilton and the godunov-type formulations. *Continuum Mechanics and Thermodynamics*, 30(6):1343–1378, jan 2018. URL: <https://doi.org/10.1007%2Fs00161-018-0621-2>, doi: 10.1007/s00161-018-0621-2.
- [PS16] V. P. Pavlov and V. M. Sergeev. Fluid dynamics and thermodynamics as a unified field theory. *Proceedings of the Steklov Institute of Mathematics*, 294(1):222–232, aug 2016. URL: <https://doi.org/10.1134%2Fs0081543816060146>, doi: 10.1134/s0081543816060146.
- [RB11] A.P. Reynolds and O.I. Bogoyavlenskij. Lie algebra structures for four-component hamiltonian hydrodynamic type systems. *Journal of Geometry and Physics*, 61(12):2400–2409, dec 2011. URL: <https://doi.org/10.1016%2Fj.geomphys.2011.07.013>, doi:10.1016/j.geomphys.2011.07.013.
- [Rey10] P. Reynolds. Hamiltonian systems of hydrodynamic type. 2010. URL: <http://hdl.handle.net/1974/6254>.
- [Rom01] E. I. Romensky. Thermodynamics and hyperbolic systems of balance laws in continuum mechanics. In *Godunov Methods*, pages 745–761. Springer US, 2001. URL: [https://doi.org/10.1007%2F978-1-4615-0663-8\\_75](https://doi.org/10.1007%2F978-1-4615-0663-8_75), doi:10.1007/978-1-4615-0663-8\_75.
- [RS15] Tommaso Ruggeri and Masaru Sugiyama. *Rational Extended Thermodynamics beyond the Monatomic Gas*. Springer International Publishing, 2015. URL: <https://doi.org/10.1007%2F978-3-319-13341-6>, doi:10.1007/978-3-319-13341-6.
- [Rug] Tommaso Ruggeri. Global existence of smooth solutions and stability of the constant state for dissipative hyperbolic systems with applications to extended thermodynamics. In *Trends and Applications of Mathematics to Mechanics*, pages 215–224. Springer Milan. URL: [https://doi.org/10.1007%2F88-470-0354-7\\_17](https://doi.org/10.1007%2F88-470-0354-7_17), doi:10.1007/88-470-0354-7\_17.
- [Sav14] Savoldi. On deformations of one-dimensional poisson structures of hydrodynamic type with degenerate metric. 2014. URL: <http://arxiv.org/abs/1410.3361>.
- [Sav15] Andrea Savoldi. Degenerate first-order hamiltonian operators of hydrodynamic type in 2d. *Journal of Physics A: Mathematical and Theoretical*, 48(26):265202, jun 2015. URL: <https://doi.org/10.1088%2F1751-8113%2F48%2F26%2F265202>, doi:10.1088/1751-8113/48/26/265202.

- [Sav16a] A. Savoldi. On poisson structures of hydrodynamic type and their deformations. 2016. URL: [https://repository.lboro.ac.uk/articles/On\\_Poisson\\_structures\\_of\\_hydrodynamic\\_type\\_and\\_their\\_deformations/9373796](https://repository.lboro.ac.uk/articles/On_Poisson_structures_of_hydrodynamic_type_and_their_deformations/9373796).
- [Sav16b] Andrea Savoldi. On deformations of one-dimensional poisson structures of hydrodynamic type with degenerate metric. *Journal of Geometry and Physics*, 104:246–276, jun 2016. URL: <https://doi.org/10.1016%2Fj.geomphys.2016.03.002>, doi:10.1016/j.geomphys.2016.03.002.
- [Sch10] Rudolf Schmid. Infinite-dimensional lie groups and algebras in mathematical physics. *Advances in Mathematical Physics*, 2010:1–35, 2010. URL: <https://doi.org/10.1155%2F2010%2F280362>, doi:10.1155/2010/280362.
- [Ser91] Denis Serre. Richness and the classification of quasilinear hyperbolic systems. In *Multidimensional Hyperbolic Problems and Computations*, pages 315–333. Springer New York, 1991. URL: [https://doi.org/10.1007%2F978-1-4613-9121-0\\_24](https://doi.org/10.1007%2F978-1-4613-9121-0_24), doi:10.1007/978-1-4613-9121-0\_24.
- [Ser99] Denis Serre. *Systems of Conservation Laws 1*. Cambridge University Press, may 1999. URL: <https://doi.org/10.1017%2F9780511612374>, doi:10.1017/cbo9780511612374.
- [Sev91] Michael Sever. Hyperbolic systems of conservation laws with some special invariance properties. *Israel Journal of Mathematics*, 75(1):81–104, feb 1991. URL: <https://doi.org/10.1007%2Fbf02787183>, doi:10.1007/bf02787183.
- [Sev95] M. Sever. Hyperbolic systems of conservation laws with a strict riemann invariant. *Journal of Differential Equations*, 122(2):239–266, nov 1995. URL: <https://doi.org/10.1006%2Fjdeq.1995.1147>, doi:10.1006/jdeq.1995.1147.
- [SH01] D Serre and Y Horie. Systems of conservation laws 2: Geometric structures, oscillations and mixed problems. *Applied Mechanics Reviews*, 54(4):B58–B59, jul 2001. URL: <https://doi.org/10.1115%2F1.1383671>, doi:10.1115/1.1383671.
- [She83] M. B. Sheftel'. On the infinite-dimensional noncommutative lie-bcklund algebra associated with the equations of one-dimensional gas dynamics. *Theoretical and Mathematical Physics*, 56(3):878–891, sep 1983. URL: <https://doi.org/10.1007%2Fbf01086255>, doi:10.1007/bf01086255.
- [She86] M. B. Sheftel'. Integration of hamiltonian systems of hydrodynamic type with two dependent variables with the aid of the lie—backlund

- group. *Functional Analysis and Its Applications*, 20(3):227–235, jul 1986. URL: <https://doi.org/10.1007%2Fbf01078475>, doi: 10.1007/bf01078475.
- [Sle84] M Slemrod. Dynamics of first order phase transitions. *Phase transformations and material instabilities in solids*, pages 163–203, 1984.
- [SO20] Xiaocheng Shang and Hans Christian Ottinger. Structure-preserving integrators for dissipative systems based on reversible–irreversible splitting. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 476(2234):20190446, feb 2020. URL: <https://doi.org/10.1098%2Frspa.2019.0446>, doi:10.1098/rspa.2019.0446.
- [Ste90] G. W. Stewart. Matrix perturbation theory, 1990.
- [Str00] I. A. B. Strachan. Degenerate bi-hamiltonian structures of the hydrodynamic type. *Theoretical and Mathematical Physics*, 122(2):247–255, feb 2000. URL: <https://doi.org/10.1007%2Fbf02551201>, doi:10.1007/bf02551201.
- [Str19] Ian A. B. Strachan. A construction of multidimensional dubrovin-novikov brackets. *Journal of Nonlinear Mathematical Physics*, 26(2):202–213, mar 2019. URL: <https://doi.org/10.1080%2F14029251.2019.1591716>, doi:10.1080/14029251.2019.1591716.
- [Tad84] Eitan Tadmor. Skew-selfadjoint form for systems of conservation laws. *Journal of Mathematical Analysis and Applications*, 103(2):428–442, oct 1984. URL: <https://doi.org/10.1016%2F0022-247x%2884%2990139-2>, doi:10.1016/0022-247x(84)90139-2.
- [Thi00] Thiffeault. Classification, casimir invariants, and stability of lie-poisson systems. 2000. URL: <http://arxiv.org/abs/math-ph/0009017>.
- [TM98] JEAN-LUC THIFFEAULT and P. J. MORRISON. Invariants and labels in lie-poisson systems. *Annals of the New York Academy of Sciences*, 867(1 NONLINEAR DYN):109–119, dec 1998. URL: <https://doi.org/10.1111%2Fj.1749-6632.1998.tb11253.x>, doi:10.1111/j.1749-6632.1998.tb11253.x.
- [TM00] Jean-Luc Thiffeault and P.J. Morrison. Classification and casimir invariants of lie-poisson brackets. *Physica D: Nonlinear Phenomena*, 136(3-4):205–244, feb 2000. URL: <https://doi.org/10.1016%2Fs0167-2789%2899%2900155-4>, doi:10.1016/s0167-2789(99)00155-4.



- [Tsa85] Sergey Petrovich Tsarev. Poisson brackets and one-dimensional hamiltonian systems of hydrodynamic type. In *Doklady Akademii Nauk*, volume 282, pages 534–537. Russian Academy of Sciences, 1985.
- [Tsa91] S P Tsarev. THE GEOMETRY OF HAMILTONIAN SYSTEMS OF HYDRODYNAMIC TYPE. THE GENERALIZED HODOGRAPH METHOD. *Mathematics of the USSR-Izvestiya*, 37(2):397–419, apr 1991. URL: <https://doi.org/10.1070%2Fim1991v037n02abeh002069>, doi:10.1070/im1991v037n02abeh002069.
- [Ver86] John M. Verosky. First-order conserved densities for gas dynamics. *Journal of Mathematical Physics*, 27(12):3061–3063, dec 1986. URL: <https://doi.org/10.1063%2F1.527236>, doi:10.1063/1.527236.
- [Ver87] John Verosky. The hamiltonian structures of the nonlinear schrodinger equation in the classical limit. *Journal of Mathematical Physics*, 28(5):1094–1096, may 1987. URL: <https://doi.org/10.1063%2F1.527553>, doi:10.1063/1.527553.
- [Vis18] Vishnoi. Geodesic convex optimization: Differentiation on manifolds, geodesics, and convexity. 2018. URL: <http://arxiv.org/abs/1806.06373>.
- [Vit19] R. Vitolo. Computing with hamiltonian operators. *Computer Physics Communications*, 244:228–245, nov 2019. URL: <https://doi.org/10.1016%2Fj.cpc.2019.05.012>, doi:10.1016/j.cpc.2019.05.012.
- [VM20] Francesco C. De Vecchi and Paola Morando. The geometry of differential constraints for a class of evolution PDEs. *Journal of Geometry and Physics*, page 103771, jun 2020. URL: <https://doi.org/10.1016%2Fj.geomphys.2020.103771>, doi:10.1016/j.geomphys.2020.103771.
- [Web15] G. M. Webb. Multi-symplectic, lagrangian, one-dimensional gas dynamics. *Journal of Mathematical Physics*, 56(5):053101, may 2015. URL: <https://doi.org/10.1063%2F1.4919669>, doi:10.1063/1.4919669.
- [Win51] Aurel Wintner. On riemann metrics of constant curvature. *American Journal of Mathematics*, 73(3):569, jul 1951. URL: <https://doi.org/10.2307%2F2372308>, doi:10.2307/2372308.
- [WM81] Alan Weinstein and Philip J. Morrison. Comments on: The maxwell-vlasov equations as a continuous hamiltonian system. *Physics Letters A*, 86(4):235–236, nov 1981. URL: [https://doi.org/10.1016%2F0378-4378\(81\)90001-1](https://doi.org/10.1016%2F0378-4378(81)90001-1), doi:10.1016/0378-4378(81)90001-1.

- doi.org/10.1016%2F0375-9601%2881%2990496-5, doi:10.1016/0375-9601(81)90496-5.
- [WZ06] G M Webb and G P Zank. Fluid relabelling symmetries, lie point symmetries and the lagrangian map in magnetohydrodynamics and gas dynamics. *Journal of Physics A: Mathematical and Theoretical*, 40(3):545–579, dec 2006. URL: <https://doi.org/10.1088%2F1751-8113%2F40%2F3%2F013>, doi:10.1088/1751-8113/40/3/013.
- [Zor16] Vladimir A. Zorich. *Mathematical Analysis II*. Springer Berlin Heidelberg, 2016. URL: <https://doi.org/10.1007%2F978-3-662-48993-2>, doi:10.1007/978-3-662-48993-2.