




Boundaries

- Regulate flows of organisms, materials, energy and information across landscapes
- Importance is disproportionate to area occupied
- Characteristics
 - Different from adjacent patches
 - Magnitude of difference and location matters
 - $\boldsymbol{-}$ Function depends on the flow
 - Steeper gradients than in the two adjoining areas
 - Wide or narrow
 - 3-dimensional
- Similar terms: edges, ecotones, gradients, clines, transition zones, interfaces
- "Edges are one of the most extensively researched areas of ecology" (Ries et al. 2004. Ann. Rev. Ecol. & Syst. 35:491-522.)

 enasso et al. 2003 BioScience 53:750-758







Ecological Boundaries: 4 Main Classes of Boundary Traits

- · Origin and maintenance
- Spatial Structure
- Function
- · Temporal dynamics

Strayer et al. 2003

Origination and maintenance


- Investigative or tangible
 - Did the boundary originate in a scientist's mind or in nature
- Causal or consequential
 - Fence vs a natural forest-field boundary
- Contemporary or relict
 - Arisen from forces still in operation or forces no longer operating
- Endogenous or exogenous origin
 - Does the boundary originate from processes operating within or outside of the system of patches (e.g. succession vs. lava flow)
- Endogenous or exogenous controls (maintenance or suppression)
 - Are internal or external forces maintaining the boundary


Strayer et al. 2003

Main Spatial Boundary Attributes (Structure)

- Grain
- Extent
- Thickness
- Geometry of adjacency
- Gradient (abruptness or steepness)
- Patch contrast
- Integrity (perforated vs. unbroken)
- Shape

Strayer et al. 2003





Temporal Dynamics

- Does the boundary change in structure or function over time?
- Mobility (is the boundary stationary or moving)?
- What is the age and history of the boundary?



Strayer et al. 2003

Functional View

- Corridor = increased rate of flow relative to diffuse flow in matrix. Channelization and movement.
- Boundary = Point of Interaction between landscape patches. Regulate flux

Puth and Wilson 2001

Boundary-Corridor Continuum

- Opposite ends of permeability
- Similar conceptually
- · Differ only in the direction and rates of flow
- Boundaries and corridors occupy small spatial extents relative to their large influence on ecological flows
- A linear feature can act as both a boundary and a corridor, depending on the mover or process
- Commonly manipulated by humans therefore they should be a central consideration to managers


Puth and Wilson 2001

The function and dynamics of landscape corridors





Taiga & muskeg, central Alaska





Natural processes can also influence habitat connectivity.


Sagebrush steppe /juniper woodland mosaic, southwestern Idaho, USA

Sagebrush steppe habitat is declining reducing the amount and connectivity of habitat for sagebrush steppe obligate species.



Fragmentation has three recognized components:

- 1- Habitat loss
- 2- Reduction in the size of patches
- 3- Decreased connectivity between patches (habitat isolation)





400 Size distribution for remnant patches of natural vegetation on the Fleurieu 350 Peninsula, South Australia. 300 Number of patches 250 200 150 100 50 < 10 11 - 100 100 - 1000 > 1000 Size of vegetation patch (ha)

Why are small populations vulnerable to decline and extinction?

<u>Demographic stochasticity</u> - random variation in population parameters such as birth rate, death rate and sex ratio.

For example, if a small population of a short-lived species experiences, by chance, a low birth rate in two successive years, the immediate probability of survival of the population may be greatly reduced.

Genetic stochasticity - random genetic processes that can lead to a loss of genetic variation and a reduced capacity for a population to resist recessive lethal alleles, or to respond to changing environmental conditions.

Inbreeding depression and genetic drift can contribute to a loss of genetic variation in small populations.

Source: Bennett 200

Why are small populations vulnerable to decline and extinction ? (continued)

Environmental stochasticity - random variation in environmental processes that can affect a population.

For example fluctuations in temperature, rainfall, food resources, and populations of predators and competitors.

<u>Natural catastrophes</u> such as floods, fire, drought, hurricanes and earthquakes, occur at irregular intervals and can have a major effect on population survival.

In fragmented environments, small isolated remnants can be totally burned and the local population of a species eliminated.

ource: Bennett 2003

Corridor terminology

General definition: A corridor is a narrow linear strip of land that differs in structure from the surrounding matrix and facilitates movement of species and process between habitats



<u>Similar terms:</u> wildlife corridor, greenway, greenbelt, shelterbelt, buffer strip, landscape bridge, wildlife underpass


Functions can be related to:


Movement of species
Gene flow
Movement of processes

Biodiversity protection
Site recolonization of locally
extinct species

Water management
Recreation
Environmental modification

Enhancement of agroforestry production








Habitat corridors are likely to be more effective means of promoting landscape connectivity:


- Where a large part of the landscape is modified and inhospitable to native species
- For species that are habitat specialists or have obligate dependence on undisturbed habitats
- For species with limited dispersal distances
- Where the goal is to maintain continuity of populations between habitats
- Where maintenance of ecosystem processes require continuous habitat for their function

Adapted from Bennett 2003





















| Landscape corridors can enhance connectivity | | | |
|--|--|--|--|
| Landscape configuration | Fine-scale (1 -10 km) | Mid-scale (10-100s km) | Broad-scale (100-1000s km) |
| Habitat corridor | Hedgerows, streams, forest corridors | Rivers & associated riparian vegetation, broad linkages between reserves | Major river systems, mountain ranges |
| Stepping stones | Patches of native vegetation, small wetlands | Series of small reserves, woodland patches | Wetlands along flight paths, alpine habitats |
| Habitat mosaic | Patchy vegetation in farmland, mosaics of gardens, parks and natural areas | Mosaics of regenerating forest in forest blocks, patchy vegetation resulting from frequent disturbance | Regional mosaics supporting different vegetation communities |











Observations about functionality of landscape corridors:

- No single corridor vegetation structure serves all species equally well and some landscape patterns may actually serve as a barrier to species movement.
- The function of corridors is dynamic as landscapes change through time responding to disturbances and succession.
- Corridors may enhance the movement of invasive or otherwise undesirable species.
- The movement of wildfire within the landscape may be either enhanced or restricted by corridors.
- In some cases, landscape structure has been specifically modified
 to serve other purposes (e.g. fire breaks, flood zones, walkways,
 greenbelts) and may or may not be effective as biological
 corridors. In some cases minor modifications can make them
 more effective biological corridors.

Landscape Ecol (2015) 30:21–50 DOI 10.1007/s10980-014-0107-y

REVIEW ARTICLE

What can studies of woodland fragmentation and creation tell us about ecological networks? A literature review and synthesis

Jonathan W. Humphrey · Kevin Watts · Elisa Fuentes-Montemayor Nicholas A. Macgregor · Andrew J. Peace · Kirsty J. Park Synthesised 104 studies, published 1990- 2013

- Development of ecological networks could help <u>reverse the effects of habitat</u> <u>fragmentation</u> on woodland biodiversity in temperate agricultural landscapes.
- Improving or <u>expanding existing habitat patches</u> is more effective compared to landscape-scale rehabilitation (e.g. creating new habitat or corridors in the landscape).
- Species responses (richness, diversity, occurrence) were strongly influenced by patch area, patch characteristics (e.g. stand structure) and isolation (e.g. distance between habitat patches).
- Very few studies have assessed all local and landscape variables together.
 Further information on impacts of different attributes of ecological networks in temperate agricultural landscapes is urgently needed.

Considerations in the design and management of conservation corridors

Biological issues

- Biological purpose
- Behavior and ecology of species
- Structural connectivity
- Quality of surrounding habitat
- Quality of corridor
- Corridor width and edge effects
- Location
- Monitoring capability

Adequacy of resources Support from local communities

Status and tenure of land

Management responsibility

Integration with other land

Socio-political issues

- management programs Community education and
- awareness
- Strategic approach to planning
- Most efficient use of conservation funds

Adapted from Bennett 2003