


Variation in climate


Differences in climate may include two types of variation:







- **Climatic variation**- changes in climate from one year to the next
- **Climatic change**- changes in climate over time, changes in the mean values (temperature, precipitation, precipitation seasonality, etc.)






Human-induced global changes



- Climate change
- Increased human population density and resource demand
- Increasing intensive and extensive land use
- Nitrogen fixation and agricultural application
- Fragmentation (smaller and more isolated habitats)
- Increased wildland-urban interface issues
- Fire exclusion & fuel accumulation
- Exotic species introductions
- Altered communities, processes and structures
- Species extinction













- Climate Envelope Models**
1. Describe the "climate envelope" which is the climate where a species currently lives (precipitation, temperature, etc. etc.)
 2. Then map the geographic shift of that "envelope" under a climate change scenario. Different time periods can be mapped.
 3. Several scenarios are often mapped because of uncertainty in climate predictions.








Conservation Approaches
Relationship between three conservation approaches that land managers may undertake and possible responses of individual species to environmental disruptions.

Component redundancy- suggests that in natural systems greater ecosystem resilience, despite changing climates, may be achieved by increasing species and community redundancy,

Functional redundancy- is the idea that different components of a system can fulfill the same functions, thereby producing the same result.

Increased connectivity- suggests that success is achieved by ensuring that suitable habitats are always within easy reach of one another.

Dunwiddie et al. 2009



Seeding and planting sagebrush

- Seeding sagebrush generally unsuccessful
- Seed sources often do not contain information about source or climate of where the seed was obtained.
- Match the climate of origin of the seed source to the place where it is planted (more important than subspecies, especially if seeds come from far away)



Germino 2017



Abies grandis – Abies concolor complex





Table 2. List of correlations between the environmental variables and factors F1, F2 and F3 from the discriminant analysis.

Variable	F1	F2	F3
Bio-1 Annual Mean Temperature	-0.293	-0.147	0.524
Bio-2 Mean Diurnal Range Temp.	0.661	0.410	-0.482
Bio-3 Isothermality (Bio2/Bio7) (*100)	0.300	0.503	-0.005
Bio-4 Temperature Seasonality	0.464	-0.089	-0.399
Bio-5 Max. Temp. of Warmest Month	-0.137	0.112	0.005
Bio-6 Min. Temp. of Coldest Month	-0.620	-0.091	0.547
Bio-7 Temperature Annual Range	0.551	0.146	-0.543
Bio-8 Mean Temp. of Wettest Quarter	0.791	-0.243	0.082
Bio-9 Mean Temp. of Driest Quarter	-0.729	0.131	0.222
Bio-10 Mean Temp. of Warmest Quarter	-0.084	-0.191	0.402
Bio-11 Mean Temp. of Coldest Quarter	-0.399	-0.020	0.576
Bio-12 Annual Precipitation	-0.481	-0.307	0.487
Bio-13 Precipitation of Wettest Month	-0.437	-0.209	0.516
Bio-15 Precipitation Seasonality	-0.272	0.220	0.539
Bio-16 Precipitation of Wettest Quarter	-0.476	-0.220	0.509
Bio-17 Precipitation of Driest Quarter	-0.134	-0.784	-0.060
Bio-18 Precipitation of warmest Quarter	0.624	-0.589	-0.162
Bio-19 Precipitation of Coldest Quarter	-0.385	-0.130	0.516
Solar Insolation	0.806	0.350	-0.173

Ott et al. 2015

Adaptation to new environments



Ott et al. 2015

Conclusions

- Hybrids occupy distinct habitats that are variably intermediate between *A. grandis* and *A. concolor* (parental taxa)
- Hybrid habitats have lower precipitation in the summertime than associated with either progenitor.
- We hypothesize that moisture stress and high levels of solar insolation in the summertime associated with hybrid populations have played an important role in the observed patterns of introgression of key traits: leaf angle, specific leaf area and abundance of adaxial stomates.
- Phenotypic plasticity may allow hybrids to adapt to an environment with a higher degree of variability in moisture stress.
- Hybrids may have an adaptive advantage over *A. grandis*.

Ott et al. 2015

References

- Amiro et al. 2001
 Ayres, M.P. and Lombardero, M.J. 2000. Assessing the consequences of global change for forest disturbance from herbivores and pathogens. *Science of the Total Environment*. 262: 263-286.
- Bachelet, D., Nelson, R.P., Lenihan, J.M. and Drapek, R.J. 2001. Climate change effects on vegetation distribution and carbon budget in the united states. *Ecosystems*. 4:164-185.
- Bachelet, D. and R.P. Nelson. 2000. Chapter 2: Biome redistribution under climate change. USDA Forest Service Gen Tech Rep. RMRS-GTR-59.
- Bartlein et al. 1997
 Canadian Forest Service. 2001. Forest fire: context for the Canadian Forest Service's science program. 2001. Available [Online]: <http://www.nrcan-nrcan.gc.ca/cfs-scf/science/context_fire/index_e.html>. Accessed November 2001.
- Carter. 1964. Effects of drought on mesquite. *Journal of Range Management*. 17:275-276.
- Dale, V.H., Joyce, L.A., McNulty, S., Neilson, P., Ayres, M.P., Flanagan, M.D., Benson, P.J., Irland, L.C., Lugo, A.E., Peterson, C.J., Simberloff, D., Swanson, F.J., Stocks, B.J. and Wotton, B.M. 2001. Climate change and forest disturbances. *BioScience*. 51:723-734.
- Dell'Osso et al. 2004
 Dunwiddie, P.W. S.A. Hall, M.W. Ingraham, J.D. Bakker, K.S. Nelson, R. Fuller and E. Gray. 2009. Rethinking Conservation Practice in Light of Climate Change. *Ecological Restoration* 27:320-329.
- Flanagan et al. 1998
 Frelich, L.E. and P.B. Reich. 2010. Will environmental changes reinforce the impact of global warming on the prairie-forest border of central North America? *Frontiers in Ecology and the Environment* 8:371-378.

References

- Gates, D.M. 1993. Climate change and its biological consequences. Sinauer, Sunderland MA. 280p.
- Hansen, A.J., R.P. Neilson, V.H. Dale, C.H. Flather, L.R. Iverson, D.J. Currie, S. Shafer, R. Cook, P.J. Bartlein. 2001. Global change in forests: responses of species, communities, and biomes. *BioScience*. 51:765-779.
- He et al. 2002
 Herbel, Ares & Wright. 1972. Drought effects on a semidesert grassland range. *Ecology*. 53:1084-1093.
- Mackenzie et al. 2004
 Mattson & Haack. 1987. The role of drought in outbreaks of plant-eating insects. *BioScience*. 37:110-118.
- McMatty, J.P. 2001. Ecological Consequences of Recent Climate Change. *Conservation Biology*. 15:320-331.
- Nelson 1993

References

- Peterson, C.J. 2000. Catastrophic wind damage to north American forests and the potential impact of climate change. *Science of the Total Environment*. 262: 287-311.
- Polley, H.W. 1999. Implications of rising atmospheric carbon dioxide concentrations for rangelands. *Journal of Range Management*. 52:561-577.
- Price and Rind 1994
 Rapp 2004
 Rehefeld, Crookston, Warwell and Evans. 2006. Empirical analysis of plant-climate relationships for the western United States. *International J. Plant Sci.* 167:1123-1150. http://forest.moscowfsl.wsu.edu/gems/Biogeography.pdf
- Simms 1988
 Swetnam & Betancourt. 1990. Fire-southern oscillation relations in the southwestern United States. *Science*. 249:1017-1020.
- UN Intergovernmental Panel on Climate Change 2000
 Veblen et al. 2000
 Whittlock et al. 2003
 World Health Organization (www.who.int/globalchange/climate/summary/en/)
 Young. 1956. Effects of the 1949-1954 drought on the ranges of Texas. *Journal of Range Management*. 9:139-145.