

An Agent Based Model of Diel Vertical Migration in *Mysis diluviana*



Nick J. Strayer¹, Jason D. Stockwell², Brian P. O'Malley³, Sture Hansson⁴

¹College of Engineering and Mathematical Sciences, ^{2,3}Rubenstein Ecosystem Science Laboratory, ⁴Stockholm University


Results

- Mysis survival rate is highly sensitive to perturbations in their feeding efficiency.
- Water temperature limits migration extent for the middle of the year whereas light levels do for the rest.

Introduction

- *Mysis diluviana* (*Mysis*) is a small macro-invertebrate crustacean that lives in Lake Champlain and other deep glacial lakes.
- *Mysis* migrate up and down the water column daily in a process called "diel vertical migration."
- We used publicly available data and previous studies to construct a model that simulates the environmental pressures on a *Mysis* at an hourly time interval over the course of a year.


Mysocline




- Light intensity and thermocline data combine to paint a picture of mysis migration extent.

Agent Based Modeling

- Simulates an individual *Mysis*.
- Every hour draws are taken from probability distributions to decide if the *mysis* migrates, how much they feed, etc.
- Good for complex scenarios with noisy input variables (e.g. cloud cover for light intensity).
- Many individuals are simulated to get a sense of population level trends.




Energy Conditions



- Each line represents an individual being simulated over the year.
- Lines that drop to or below zero are individuals who have starved


Mortality Rates



- There is greater sensitivity to changes in reward units than migration cost.
- Results follow expected trends based upon ecological principles.

Interactive Application

Mysis Condition



- Utilizes R Shiny Servers to allow user to interact with model parameters without tinkering with code.
- Rapidly facilitates insights into effects of changing parameters.
- Hosted on the web for anyone to use. (nstrayer.shinyapps.io/mysisApp)
- Helps make research reproducible by others.

Future Directions

- Probe the possibility of multiple stable migration patterns.
- Dig in to specific aspects of the model. E.g. predation risk, benthic food availability
- Utilize real data in model inputs such as thermocline depth and food availability.

References

- Jensen O. P., Hrabik T. R., Martell S. J. D., Walters C. J., Kitchell J. F., Diel vertical migration in the Lake Superior pelagic community. II. Modeling trade-offs at an intermediate trophic level, *Can. J. Fish. Aquat. Sci.* 63: 2296-2307 (2006)
- Grimm V. et al, A standard protocol for describing individual-based and agent-based models, *Ecological Modeling*, V. 198(1-2): 115-126 (2006)

Acknowledgements

- Office of Undergraduate Research.
- The entire UVM Honors College staff for support.
- Peter Euclide for *Mysis* insights.
- Professors James Bagrow & Daniel Benteil thesis advising
- Many more.