

The Companion Frequency and Orbital Distribution of M-Dwarf Binaries

Nicholas Susemihl¹, Michael Meyer¹

¹ – Department of Astronomy, University of Michigan, Ann Arbor

Motivation & Goals

- M-Dwarf multiplicity is not well constrained
- We seek to **fit a model to the surface density distribution** of M-Dwarfs using point estimates from surveys over a range of orbital separations
- **Calculate a constrained frequency**
 - Specific ranges of mass ratio (q) and semi-major axis (a) from sensitivity of data
 - Assume companion mass ratio distribution does not depend on orbital separation
 - Only account for binary systems (no triples, quadruples, etc.)
- **Extrapolate constrained frequency** to find new value covering broad ranges of mass ratio and semi-major axis
- Compare to other stellar multiplicity estimates

Methods


- Collected data from five M-Dwarf multiplicity surveys

Reference	Semi-Major Axis Sensitivity Range (AU)	Multiplicity Estimate ($q \geq 0.6$)
Delfosse et. al. (1998)	0.00 - 4.63	0.04 ± 0.018
Fischer & Marcy (1992)	0.04 - 4.00	0.08 ± 0.034
Cortes-Contreras et. al. (2016)	2.60 - 29.5	0.07 ± 0.012
Janson et. al. (2012)	3.00 - 227	0.18 ± 0.016
Ward-Duong et. al. (2015) A	3.00 - 100	0.11 ± 0.022
Ward-Duong et. al. (2015) B	100 - 10,000	0.07 ± 0.017
- Used MCMC with chi-squared likelihood to fit a log-normal model to separation distribution with three parameters ($\log_{10} \mu$, $\log_{10} \sigma$, A)
$$\phi = A * \frac{e^{-(\log_{10} a - \log_{10} \mu)^2 / 2 \log_{10} \sigma^2}}{\log_{10} \sigma * \sqrt{2\pi}}$$
- Referenced mass ratio distribution from Reggiani & Meyer (2013)
$$\psi = q^{0.25 \pm 0.29}$$
- Calculated frequency
$$f = \int_{q_{min}}^{q_{max}} \psi dq * \int_{\log_{10} a_{min}}^{\log_{10} a_{max}} \phi d \log_{10} a$$
- Analyzed Goodness of Fit
 - $\chi^2_{red.} (3 d.o.f.) = 2.14$
 - $P(X \geq \chi^2_{red.}) = 0.09$


Despite this low probability, we do not reject the null hypothesis that the data came from this model

Results

- **Constrained Frequency** ($0.60 \leq q \leq 1.00$, $0.00 \leq a \leq 10,000$): **0.239 ± 0.04**
- **Overall Frequency** ($0.10 \leq q \leq 1.00$, $0.00 \leq a \leq \infty$): **0.481 ± 0.08**



Analysis of Fit



Conclusions

- Peak in orbital surface density distribution of M-Dwarf binaries occurs at about 40 AU (larger than the estimate from Winters et al. 2018 and slightly smaller than Raghavan et al. 2010 for FGK stars)
- **About half of all M-Dwarfs have a low mass companion**, many of which may be brown dwarfs after extrapolating down to this regime
- Comparisons
 - Extrapolated results from surveys of other spectral type over constrained regions of mass ratio and separation

Survey	Raghavan et. al. (2010) - FGK Stars	De Rosa et. al. (2013) - A Stars
Multiplicity Fraction $q = [0.6, 1]$, $a = [0.00, 10000 \text{ AU}]$	$f = 0.230 \pm 0.032$	$f = 0.238 \pm 0.026$

- **Overall, multiplicity fraction does not vary significantly across spectral types** over $q = [0.6, 1]$ and $a = [0.00, 10000 \text{ AU}]$

Acknowledgements

We thank the Formation and Evolution of Planetary Systems group (<https://sites.lsa.umich.edu/feps/>) for their consistent support. We would also like to thank Dr. Max Moe and Dr. Kimberly Ward-Duong for their insightful advice.

Literature Cited

Cortes-Contreras et. al. 2016, *Astro. & Astrophys.*, FC23
De Rosa et. al. 2013, *Mon. Not. R. Astro. Soc.* 437, 1216-1240
Delfosse et. al. 1998, *Astro. & Astrophys.*, 344, 897-910
Fischer and Marcy 1992, *The Astrophys. Journal*, 396, 178-194

Janson et. al. 2012, *The Astrophys. Journal*, 754, 44-70
Raghavan et. al. 2010, *The Astrophys. Journal*, 190, 1-42
Reggiani and Meyer 2013, *Astro. & Astrophys.*, 553, A124
Ward-Duong et. al. 2015, *Mon. Not. R. Astro. Soc.* 000, 1-34
Winters et. al. 2019, *The Astron. Journal*, 157, 6, 216

