

References from electronic databases (**n = 10124**)
(as **n = 10093** surveys)
English searches (**n = 7572**)
Non-English searches (**n = 2552**)

References from other sources (**n = 1434**)
Citation searching (**n = 7**)
Grey literature:
2023 Union Conference abstracts (**n=1427**)

References removed (**n = 1316**)
Duplicates identified manually (**n = 249**)
Duplicates identified by Covidence (**n = 1067**)

Studies screened
(**n = 10211**)

Surveys excluded (**n = 9993**)

Studies included
in Horton et al.
2016 review
(**n = 77**)

WHO National
Prevalence
Survey
Reports (**n = 34**)

Studies sought for retrieval
(**n = 107**)
English language (**n =103**)
Non-English language (**n = 4**)

Studies not retrieved (**n = 0**)

Studies assessed for eligibility (**n = 218**)

Studies excluded (**n = 118**)
Editorial (**n = 4**)
Wrong setting (**n = 1**)
Wrong outcomes (**n = 40**)
Wrong comparator (**n = 4**)
Wrong intervention (**n = 7**)
Wrong study design (**n = 10**)
Wrong patient population (**n = 34**)
Did not report sex-stratified TB prevalence
estimates (**n=18**)

Studies included in review (**n=100**)
English language (**n =100**)
Non-English language (**n =0**)
Surveys included primary quantitative analysis* (**n = 102**)