

Age distribution totals across 64 TB prevalence surveys with standard age groups													
Description	15-24 years (N)	25-34 years (N)	35-44 years (N)	45-54 years (N)	55-64 years (N)	65+ years (N)	15-24 years (%)	25-34 years (%)	35-44 years (%)	45-54 years (%)	55-64 years (%)	65+ years (%)	Total (N)
abnormal	13,029	17,383	23,149	26,174	25,127	31,748	9.5	12.7	16.9	19.2	18.4	23.2	136,610
bacteriological	588	954	1,207	1,095	893	1,153	10	16.2	20.5	18.6	15.2	19.6	5,890
culture	90	154	207	169	134	184	9.6	16.4	22.1	18	14.3	19.6	938
eligible	528,446	429,047	349,151	259,405	159,307	149,021	28.2	22.9	18.6	13.8	8.5	8	1,874,377
participants	543,831	449,243	378,674	287,517	193,444	181,250	26.7	22.1	18.6	14.1	9.5	8.9	2,033,959
presumptive	16,552	19,782	21,479	20,594	18,709	23,556	13.7	16.4	17.8	17.1	15.5	19.5	120,672
prevalent	165	284	366	333	362	469	8.3	14.4	18.5	16.8	18.3	23.7	1,979
smear	249	356	463	427	327	387	11.3	16.1	21	19.3	14.8	17.5	2,209
sputum	9,027	9,213	9,858	9,388	9,140	9,507	16.1	16.4	17.6	16.7	16.3	16.9	56,133
symptoms	10,815	13,109	14,679	13,781	11,889	13,110	14	16.9	19	17.8	15.4	16.9	77,383