

### Sex distribution totals across 107 TB prevalence surveys

| Description     | Female (N) | Female (%) | Male (N)  | Male (%) | Total (N) |
|-----------------|------------|------------|-----------|----------|-----------|
| eligible        | 1,771,757  | 52.5       | 1,605,227 | 47.5     | 3,376,984 |
| participants    | 2,207,018  | 54.3       | 1,857,416 | 45.7     | 4,064,434 |
| presumptive     | 103,463    | 50.3       | 102,192   | 49.7     | 205,655   |
| symptoms        | 68,793     | 43.9       | 87,887    | 56.1     | 156,680   |
| abnormal        | 93,710     | 47.5       | 103,458   | 52.5     | 197,168   |
| sputum          | 66,718     | 53.4       | 58,303    | 46.6     | 125,021   |
| radiologic      | 57         | 47.9       | 62        | 52.1     | 119       |
| bacteriological | 2,821      | 32.3       | 5,917     | 67.7     | 8,738     |
| smear           | 1,321      | 29.5       | 3,163     | 70.5     | 4,484     |
| culture         | 1,246      | 28.4       | 3,135     | 71.6     | 4,381     |
| prevalent       | 1,171      | 33.2       | 2,353     | 66.8     | 3,524     |