




Number of prevalence surveys by country



Number of prevalence surveys by country




Number of prevalence surveys by country




Number of prevalence surveys by World Bank region

Survey region




Number of prevalence surveys by stratification

Prevalence stratification




Number of prevalence surveys by representativeness

Scope of prevalence review



Number of prevalence surveys by publication year



Number of prevalence surveys by start year

Survey start year

2020


2010

2000




Number of prevalence surveys by screening algorithm

Screening algorithm




Number of prevalence surveys by symptoms screened

Symptom included




Number of prevalence surveys by definition of abnormal chest X-ray


Abnormal chest X-ray definition




TB prevalence definition includes children




TB prevalence definition includes people on TB treatment




TB prevalence definition includes people identified through contact tracing




TB diagnostic method




Combination of TB diagnostic methods



Survey start years by World Bank region




Study publication years by World Bank region



Sex distribution totals across 108 TB prevalence surveys

Description	Female (N)	Female (%)	Male (N)	Male (%)	Total (N)
eligible	1,771,757	52.5	1,605,227	47.5	3,376,984
participants	2,249,595	54.2	1,898,264	45.8	4,147,859
presumptive	103,463	50.3	102,192	49.7	205,655
symptoms	68,793	43.9	87,887	56.1	156,680
abnormal	93,710	47.5	103,458	52.5	197,168
sputum	66,718	53.4	58,303	46.6	125,021
radiologic	57	47.9	62	52.1	119
bacteriological	2,821	32.3	5,917	67.7	8,738
smear	1,357	28.7	3,370	71.3	4,727
culture	1,322	27.3	3,516	72.7	4,838
prevalent	1,171	33.2	2,353	66.8	3,524


Male to female ratio of participants



Rurality distribution totals across 48 TB prevalence surveys

Description	Rural (N)	Rural (%)	Urban (N)	Urban (%)	Total (N)
eligible	1,163,242	60.1	773,579	39.9	1,936,821
participants	1,191,926	61.5	746,572	38.5	1,938,498
presumptive	67,731	65.3	36,036	34.7	103,767
symptoms	42,141	65.9	21,789	34.1	63,930
abnormal	76,372	63.5	43,858	36.5	120,230
sputum	18,135	58.8	12,725	41.2	30,860
radiologic	17	94.4	1	5.6	18
bacteriological	3,070	60.1	2,037	39.9	5,107
smear	1,191	66.8	593	33.2	1,784
culture	793	63.7	452	36.3	1,245
prevalent	637	62.3	386	37.7	1,023

Crude urban to rural ratio of participants




HIV status distribution totals across 17 TB prevalence surveys

Description	HIV negative (N)	HIV negative (%)	HIV positive (N)	HIV positive (%)	Total (N)
participants	128,042	88.7	16,306	11.3	144,348
presumptive	25,585	89.4	3,021	10.6	28,606
symptoms	15,136	91.4	1,433	8.6	16,569
abnormal	3,049	76.1	956	23.9	4,005
sputum	15,648	85.7	2,616	14.3	18,264
bacteriological	1,335	77.3	391	22.7	1,726
smear	692	89.8	79	10.2	771
culture	65	38.9	102	61.1	167
prevalent	886	86.6	137	13.4	1,023

Age distribution totals across 64 TB prevalence surveys with standard age groups													
Description	15-24 years (N)	25-34 years (N)	35-44 years (N)	45-54 years (N)	55-64 years (N)	65+ years (N)	15-24 years (%)	25-34 years (%)	35-44 years (%)	45-54 years (%)	55-64 years (%)	65+ years (%)	Total (N)
abnormal	13,029	17,383	23,149	26,174	25,127	31,748	9.5	12.7	16.9	19.2	18.4	23.2	136,610
bacteriological	588	954	1,207	1,095	893	1,153	10	16.2	20.5	18.6	15.2	19.6	5,890
culture	90	154	207	169	134	184	9.6	16.4	22.1	18	14.3	19.6	938
eligible	528,446	429,047	349,151	259,405	159,307	149,021	28.2	22.9	18.6	13.8	8.5	8	1,874,377
participants	543,831	449,243	378,674	287,517	193,444	181,250	26.7	22.1	18.6	14.1	9.5	8.9	2,033,959
presumptive	16,552	19,782	21,479	20,594	18,709	23,556	13.7	16.4	17.8	17.1	15.5	19.5	120,672
prevalent	165	284	366	333	362	469	8.3	14.4	18.5	16.8	18.3	23.7	1,979
smear	249	356	463	427	327	387	11.3	16.1	21	19.3	14.8	17.5	2,209
sputum	9,027	9,213	9,858	9,388	9,140	9,507	16.1	16.4	17.6	16.7	16.3	16.9	56,133
symptoms	10,815	13,109	14,679	13,781	11,889	13,110	14	16.9	19	17.8	15.4	16.9	77,383

Was the study's sample population a true or close representation of the target population?



Was some form of random selection used to select the sample or was a census undertaken?



Was the likelihood of non-response bias minimal?



Were data collected directly from subjects (as opposed to a proxy)?



Was an acceptable case definition used in the study?



Was the study instrument that measured the parameter of interest shown to have reliability and validity?



Was the same mode of data collection used for all subjects?



Were the numerator and denominator for the parameter of interest appropriate?



Overall bias assessment High risk of bias Low risk of bias Moderate risk of bias Unknown

Bias assessment summary measure

Moderate risk of bias

44

Low risk of bias


48

High risk of bias

22

Unknown

1



0 10 20 30 40 50

Bias assessment summary measure by survey scope

