P8104 Homework 3

Nick Williams ntw2117

Problem 1

(1)

(2)

Problem 2

$$f_X(x) = 2xI\{0 < x < 1\}$$

(1)

$$Y=g(x)=4X-3$$

$$\frac{d}{dx}g(x)=4>0 \qquad \text{thus } g(x) \text{is an increasing function}$$

$$g^{-1}(y)=\frac{y+3}{4}$$

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|$$

$$= f_X(\frac{y+3}{4}) \left| \frac{1}{4} \right| \qquad \frac{d}{dy} g^{-1}(y) = \frac{1}{4}$$

$$= 2\frac{y+3}{4} \left(\frac{1}{4} \right)$$

$$= \frac{y+3}{8} \text{ for } -3 < y < 1$$

(2)

$$Y = g(x) = 3 - X^2$$

$$\frac{d}{dy}g(x) = -2x < 0 \quad \text{monotone } \forall \ 0 < x < 1$$

$$g^{-1}(y) = \sqrt{3 - y}$$

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|$$

$$= f_x(\sqrt{3-y}) \left| -\frac{1}{2\sqrt{3-y}} \right| \quad \frac{d}{dy} g^{-1}(y) = -\frac{1}{2\sqrt{3-y}}$$

$$= f_x(\sqrt{3-y}) \left(\frac{1}{2\sqrt{3-y}} \right)$$

$$= 2(\sqrt{3-y}) \left(\frac{1}{2\sqrt{3-y}} \right)$$

$$= 1 \text{ for } 2 < y < 3$$

Problem 3

(1)

(2)
$$X \sim U(-1,1) \text{ thus } f_X(x) = \frac{1}{2} \text{ for } -1 \le x \le 1$$

$$Y = g(x) = e^x \text{ thus } g^{-1}(y) = \log(x)$$

Because $\frac{d}{dx}g(x) = e^x > 0 \ \forall x \in [-1,1], \ g(x)$ is an increasing function and monotone; therefore

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right| \quad F_Y(y) = F_X(g^{-1}(y))$$

$$= \frac{1}{2} \left| \frac{1}{y} \right| \qquad \qquad = \frac{g^{-1}(y) + 1}{2}$$

$$= \frac{1}{2y} \qquad \qquad = \frac{\log(y) + 1}{2}$$

$$X \sim U(-1,1)$$
 thus $f_X(x) = \frac{1}{2}$ for $-1 \le x \le 1$
 $Z = g(x) = -\log(X+1)$ thus $g^{-1}(z) = e^{-z} - 1$

Because $\frac{d}{dx}g(x) = -\frac{1}{x+1} < 0 \ \forall x \in [-1,1], \ g(x)$ is a decreasing function and monotone; therefore

$$F_Z(z) = 1 - F_X(g^{-1}(z)) \qquad f_Z(z) = f_X(g^{-1}(z)) \left| \frac{d}{dz} g^{-1}(z) \right|$$

$$= 1 - F_X(e^{-z} - 1) \qquad \qquad = \frac{1}{2} \left| -e^{-z} \right|$$

$$= 1 - \frac{(e^{-z} - 1) + 1}{2} \qquad \qquad = \frac{e^{-z}}{2}$$

$$= -\frac{e^{-z}}{2}$$

Problem 4

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} \forall x \in (-\infty, \infty) \quad Y = g(x) = \log(|X + 1|)$$

g(x) is not monotone on $(-\infty, \infty)$, however it is monotone on $(-\infty, -1)$ and $(-1, \infty)$. We therefore can define:

$$A_0 = -1 \qquad \text{On } A_0 \ y = \{0\}$$

$$A_1 = (-\infty, -1) \qquad \text{On } A_1 \ y = \log(|x+1|)I\{x < -1\} \qquad g^{-1}(y) = -e^y - 1$$

$$A_2 = (-1, \infty) \qquad \text{On } A_2 \ y = \log(|x+1|)I\{x > -1\} \qquad g^{-1}(y) = e^y - 1$$

$$f_Y(y) = \sum_{i=1}^k f_X(g_i^{-1}(y)) \left| \frac{d}{dy} g_i^{-1}(y) \right| I\{y \in \Omega_Y\}$$

$$f_Y(y) = f_X(g_1^{-1}(y)) \left| \frac{d}{dy} g_1^{-1}(y) \right| + f_X(g_2^{-1}(y)) \left| \frac{d}{dy} g_2^{-1}(y) \right|$$

$$= f_X(-e^y - 1) |-e^y| + f_X(e^y - 1) |e^y|$$

$$= f_X(-e^y - 1) (e^y) + f_X(e^y - 1) (e^y)$$

$$= e^y \left(f_X(-e^y - 1) + f_X(e^y - 1) \right)$$

$$= e^y \left(\frac{1}{\sqrt{2\pi}} e^{\frac{-(-e^y - 1)^2}{2}} + \frac{1}{\sqrt{2\pi}} e^{\frac{-(e^y - 1)^2}{2}} \right) I\{y > 0\}$$

Problem 5

$$f_X(x) = cx^2 I\{-1 < x < 1\}$$

(1) Because $f_X(x) \ge 0 \forall x, c > 0$:

$$1 = \int_{-\infty}^{\infty} f_X(x) dx$$

$$= \int_{-\infty}^{\infty} cx^2 I\{-1 < x < 1\} dx$$

$$= c \int_{-1}^{1} x^2 dx$$

$$= c \frac{1}{3} x^3 \Big|_{-1}^{1}$$

$$= c \frac{1}{3} (1)^3 - c \frac{1}{3} (-1)^3$$

$$c = \frac{3}{2} > 0$$

(2) By the definition of a cumulative distribution function

$$F_X(x) = P(X \le x) = \int f_X(x)dx$$

$$= \int \frac{3}{2}x^2 I\{-1 < x < 1\}dx$$

$$= \frac{3}{6}x^3$$

$$= \frac{x^3}{2} \forall x \in (-1, 1)$$

(3) If $U \sim U(0,1)$, to find $F^1(U)$, we need to solve $y = \frac{x^3}{2}$ for x in terms of $y \in (0,1)$

$$F_X(F^{-1}(u)) = u$$

$$\frac{(F^{-1}(U))^3}{2} = (F^{-1}(U))^3 = 2u$$

$$F^{-1}(U) = 2u^{1/3} \ \forall U \in (0, 1)$$