


# P8110: Homework 3

Nick Williams  
ntw2117

## Problem 1



Based on the survival graph, it appears participants enrolled in the body-cleansing treatment are experiencing greater survival time compared to those enrolled in the routine cleansing treatment. For example, the survival function for those in the body-cleansing treatment never reaches 50% failure rate by the end of the study while the survival function for those in the routine cleansing treatment does.

## Problem 2

Test of Equality over Strata			
Test	Chi-Square	DF	Pr > Chi-Square
Log-Rank	3.7924	1	0.0515
Wilcoxon	2.8639	1	0.0906
-2Log(LR)	3.4994	1	0.0614

### log-rank test

1.  $H_0 : S_1(t) = S_0(t)$ , for all  $t \leq \tau$   
 $H_A : S_1(t) \neq S_0(t)$ , for all  $t \leq \tau$

2.

$$Q = 3.7924 \sim \chi_1^2$$

$$p = P(\chi_1^2 \leq 3.7924) = 0.0515$$

3.

$0.0515 > 0.05 \rightarrow$  fail to reject the null

4. At the 5% level of significance, using a log-rank test, there is insufficient evidence to conclude that differential survival time exists between participants enrolled in the body-cleansing treatment and those enrolled in the routine cleansing treatment

### Wilcoxon test

1.  $H_0 : S_1(t) = S_0(t)$ , for all  $t \leq \tau$   
 $H_A : S_1(t) \neq S_0(t)$ , for all  $t \leq \tau$

2.

$$Q = 2.8639 \sim \chi_1^2$$

$$p = P(\chi_1^2 \leq 2.8639) = 0.0906$$

3.

$0.0906 > 0.05 \rightarrow$  fail to reject the null

4. At the 5% level of significance, using a Wilcoxon rank test, there is insufficient evidence to conclude that differential survival time exists between participants enrolled in the body-cleansing treatment and those enrolled in the routine cleansing treatment.

### Difference between tests

The log-rank test is more sensitive to differences between survival functions at time points later in follow-up while the Wilcoxon test is more sensitive to differences earlier in time. This is because the Wilcoxon rank test applies a weight to each unique event time point according to the number of at risk individuals at that time point. This is why the p-value for the Wilcoxon test is larger in this case compared to the p-value from the log-rank test.

### Problem 3

Test of Equality over Strata			
Test	Chi-Square	DF	Pr > Chi-Square
Log-Rank	5.4722	3	0.1403
Wilcoxon	4.3055	3	0.2303
-2Log(LR)	3.8573	3	0.2773

1.  $H_0 : S_1(t) = S_2(t) = S_3(t) = S_4(t)$  for all  $t \leq \tau$   
 $H_A : \text{At least one } S_k \text{ is different for some } t \leq \tau$

2.

$$Q = 5.4722 \sim \chi_3^2$$

$$p = P(\chi_3^2 \leq 5.4722) = 0.1403$$

3.

$0.1403 > 0.05 \rightarrow$  fail to reject the null

4. At the 5% level of significance, using the generalized log-rank test, there is insufficient evidence to conclude that the survival functions are different among the 4 different groups of surface area burned.