

Patient's Name: _____
UHID: _____ IPID: _____ Age: __ Gender: Male/Female _____
D.O.A.: _____ Unit: _____

INFORMED CONSENT FOR ANAESTHESIA

When surgical intervention is required to solve the patient's current medical problems, appropriate anaesthesia is needed to reduce pain and fear during surgery. Although it provides more comfortable and safer physical condition under surgical stress in most of the situation, certain anesthesia related risk may still happen particularly in patient with prior medical ailments.

ANAESTHESIA

It is a technique which blocks pain, make you sleep and keep you safe during surgical / medical / investigative procedures. That way, people can have surgery without feeling pain. It makes surgery much safer and more comfortable.

TYPES OF ANAESTHESIA: There are 3 main types of anaesthesia are given depending on requirement of patient.

1. Local anaesthesia
2. Regional anaesthesia
3. General anaesthesia

1) LOCAL ANAESTHESIA: Local anaesthesia is a technique that renders part of the body insensitive to pain/surgical procedure done without affecting consciousness. It allows patients to undergo surgery with no pain and distress. In a Strict sense it is anaesthesia of a small part of the body such as a tooth or an area of skin.

➤ Benefits of local anaesthesia

The anaesthetic action extends for longer duration and therefore provides pain relief for several hours after the operation. Local anaesthesia is associated with reduced blood loss. It gives the patient a sense of feeling more "in control" as they are awake during the procedure.

The patient returns to normal activities like eating, walking on the same day. He may even be able to return home the same day following the surgery.

➤ Risks of local anaesthesia :(Depends on area of procedure and surgery)

Swelling at the site of injections or drips, Severe headache, blurred or double vision, dizziness or light heartedness, drowsiness, confusion anxiety, excitement, nervousness or restlessness convulsions (seizures), feeling hot, cold, or numb ringing or buzzing in the ears, shivering or trembling sweating pale skin, slow or irregular heartbeat breathing problems, unusual weakness or tiredness and occasionally cardiac arrest.

2) REGIONAL ANAESTHESIA: It is a anaesthesia affecting larger region of the body, such as a limb, arm or lower abdomen.

- **Intravenous Regional Anesthesia** - Technique: Drug injected into veins of arm or leg while using a tourniquet
- **Nerve Block**-Technique: Drug injected near nerves causing loss of sensation to the area of the operation.

- **Spinal/ Epidural Analgesia:** Technique -Drug injected through a needle/catheter placed either directly in the spinal canal or immediately outside the spinal canal or both.
- **Benefits of regional anaesthesia:**

Patient remains conscious, maintains his own airway, smooth recovery requiring less skilled nursing care as compared to general anaesthesia, reduction in surgical stress, earlier discharge and better pain relief.
- **Risks of regional anaesthesia:**

Muscle weakness in the anaesthetized limb, or difficulty in passing urine for a lower body block, Headache, which is usually short lived but can be severe and last some days. Damage to nearby blood vessels or organs very rarely. Risk of infection on or bleeding at the injection site. Rarely, nerves may be damaged resulting in long term weakness, pain, and altered sensation or even paralysis and infection to brain and spinal cord.
- 3) GENERAL ANAESTHESIA:** General anesthesia is used for surgical procedures that are extensive, those that are of uncertain duration, or when other anesthesia is not feasible. It is also used for emergency surgeries. Other determining factors include the type of surgery and patient's medical history.
- Technique:** Drug injected into the bloodstream, breathed into the lungs, or by other routes. Possible placement of a tube into the windpipe.
- General anaesthesia causes you to be unconscious throughout your operation. During the operation, your anaesthesiologist will closely monitor the functions of your heart, lungs, kidneys. This ensures the patient is not aware of events and does not feel pain during the operation. It is produced by drugs given through a vein and / or breathed from an anaesthesia machine. Once your surgery is completed, the effects of the medications will be reversed and you will regain consciousness. The anaesthetist uses special skills and equipment to monitor and manage the patient, to ensure their safety during anaesthesia and the operation. The patient may be shifted to ICU for further care including ventilator care is required.
- **Benefits of general anaesthesia:**

Quick onset, complete control of body functions, no memory of the surgery, effective for surgeries of longer duration, ability to perform surgeries in different parts of the body at the same time.
- **Risks of general anaesthesia:**
- ✓ **Common risks for ALL patients include:**

Bruising or swelling at the site of injections or drips. Nausea or vomiting (although the anaesthetist will limit or prevent this as far as possible). Sore throat from the gases and / or the breathing tube. You may notice temporary difficulty in speaking, which generally improves after some hours. Temporary muscle pains.

Temporary headache or blurred vision.
- 4) Monitored Anesthesia Care (MAC):** Is Given in Day care procedures like Endoscopy/Colonoscopy/CT/MRI and surgical cases like Circumcision, Cystoscopies etc..It includes with sedation or without sedation.
- Technique**
- With sedation-** Drug injected into the bloodstream, breathed into the lungs, or by other routes producing a semiconscious state.
- **Without sedation**
- **Benefits of MAC:** Reduced anxiety and pain, partial or total amnesia.
- **Risks of MAC:** An unconscious state, depressed breathing (requiring Ventilation), injury to blood vessels. Awareness, anxiety and/or procedure related discomfort. Probability to convert in to General Anesthesia.
- ✓ **Uncommon risks for ALL patients include:**

Awareness of activity in the operating room during anesthesia, particularly during certain operations and in some emergency situations. Eye abrasions causing pain and requiring treatment with medication and patching. Damage to teeth or dental work, lips or tongue.

✓ **Extremely rare risks for ALL patients. These may cause brain damage or death and include:**

Death or permanent disability related to anaesthesia is rare. Obstructions in the breathing passages that cannot be readily controlled, these can lead to severe difficulty with breathing. Allergy to drugs causing wheezing and rash and in rare cases, severe swelling, low blood pressure and poor circulation. Inherited muscle sensitivity to particular anaesthetic drugs (malignant hyperthermia) which cause a rapid rise in temperature, heart rate and breathing with high blood pressure and muscle rigidity. Heart attacks, strokes and pneumonia. While these are uncommon, the risks are higher for patients with diseases of the arteries or lungs and in smoker and patient with previous heart related ailments.

ADDITIONALLY, THE FOLLOWING FACTORS CAN INCREASE YOUR RISK OF COMPLICATIONS:

- Medical conditions involving your heart, lungs, kidneys or liver
- Medications, such as aspirin/clopidogrel, that can increase bleeding
- Smoking, which increases the likelihood of breathing problems
- Alcohol use, which may predispose you to liver damage
- Family history of adverse reactions to anaesthesia
- Food or drug allergies
- Sleep apnea
- Obesity
- Temporary mental confusion
- Lung infections
- Stroke
- Heart attack
- Abnormal airway (mouth and upper part of wind passage)

DECLARATION BY PATIENT / GUARDIAN:

- I acknowledge that the anaesthetist has informed me about the anaesthetic procedure, alternative treatments and answered my specific queries and concerns about this matter.
- I understand that an anaesthetist other than the consultant anaesthetist may administer the anaesthesia.
- I have been given enough time to read and understand whatever written in this form.
- I have explained all the above in the language _____ that I understand.

PROCEDURE/SURGERY PLANNED -

Type of anaesthesia :	
☐ General anaesthesia	☐ Regional anaesthesia
☐ MAC Sedation - ☐ Mild ☐ Moderate ☐ Deep	
☐ Local anaesthesia	

Individuals Risks (to be completed by the anaesthetist completing this form)

Examples of possible risks and complications specific to this patient: _____

BY SIGNING THIS FORM, I GIVE MY FULL, FREE, VOLUNTARY CONSENT WITHOUT ANY RESERVATION TO DR. _____ (name of doctor administering Anesthesia) or his/her associates, for administering anesthesia to ☐ MYSELF or ☐ MY PATIENT _____ (name of patient), being fully aware of the nature, risks and complications.

	Name	Signature	Date	Time
Patient				
Patient representative with relationship				
Reason for surrogate consent	Patient is unable to give consent because:			
Witness				
Interpreter (if necessary)				

only if Patient is a minor or unable to give consent

I, the undersigned Doctor, have explained the nature, potential risks and complications, expected benefits, expected post-anesthesia recovery, and possible alternatives to the patient/patient representative. I am confident that he / she has understood the information provided in this document.

Consent obtained by^

Dr. _____ Designation & Dept. _____ Signature: _____

^(being person administering Anesthesia or member of his /her team)