

HEALTHCARE-ASSOCIATED INFECTIONS (HAIS) - CARE BUNDLE FORM

Phlebitis prevention care bundle

Sl. No	Check list	Date:			Date:			Date:		
		M	E	N	M	E	N	M	E	N
0	Admitted with outside Peripheral Intravenous Catheter (PIVC) Yes: [], If yes mention date: No: []	Time	Time	Time	Time	Time	Time	Time	Time	Time
	Insertion Care:									
1	Assessed the need for the Peripheral Intravenous Catheter									
2	Skin preparation with 2% chlorhexidine or 70% alcohol									
3	IV cannula size									
4	Date and time of insertion mentioned over the tegaderm									
5	Number of pricks (mention 1, 2 or 3)									
	Maintenance Care:									
1	Review of Peripheral Intravenous Catheter.									
2	Hand hygiene performed before and after handling the PIVC line.									
3	Needle-less access ports connected.									
4	Scrub the IV caps with 70% alcohol swab before medication administration.									
5	Flushing the cannula before and after medication administration with 0.9% normal saline.									
6	Phlebitis score (mention score 0-5).									
7	Replacement of PVC every 96 hours / clinically indicated									
8	Prompt removal of unnecessary lines.									
	Employee code / Signature of the nurse.									

V.I.P Score (Visual Infusion Phlebitis score)	VIP score should be evaluated during each shift and documented on the bundle care	
I. V site appears healthy	0 →	No signs of phlebitis OBSERVE CANNULA
One of the following is evident: ● Slight pain or Slight redness near IV site	1 →	Possible first signs of phlebitis OBSERVE CANNULA
Two of the following are evident: ● Pain at IV site ● Redness ● Swelling	2 →	Early stage of Phlebitis RESITE CANNULA
All of the following are evident: ● Pain along path of cannula ● Erythema ● Induration	3 →	Medium stage of phlebitis RESITE CANNULA & CONSIDER TREATMENT
All of the following are evident and extensive: ● Pain along path of cannula ● Erythema ● Induration ● Palpable venous cord	4 →	Advanced stage of phlebitis or start of thrombophlebitis RESITE CANNULA & CONSIDER TREATMENT
All of the following are evident and extensive: ● Pain along path of cannula ● Erythema ● Induration ● Palpable venous cord ● Pyrexia	5 →	Advanced stage of thrombophlebitis RESITE CANNULA & INITIATE TREATMENT

Catheter associated urinary tract infection (CAUTI) care bundle.

Sl. No	Check list	Date:			Date:			Date:		
		M	E	N	M	E	N	M	E	N
0	Admitted with outside urinary catheter: Yes [], If yes mention date: No: []	Time	Time	Time	Time	Time	Time	Time	Time	Time
	Insertion Care: Foley's Silicon									
1	Accessed the need for a urinary catheter.									
2	Catheter size									
3	Aseptic technique followed.									
4	Gentle insertion of the catheter after application of new lubricant /anesthetic gel									
5	Date and time of insertion documented over the urinary bag and catheter.									
	Maintenance Care:									
1	Review on indwelling catheter necessity									
2	Hand hygiene performed before and after handling the catheter									
3	Catheter secured properly over the thigh									
4	Maintained a closed drainage system									
5	Uro bag kept below the bladder level and not touching the floor.									
6	Unobstructed urine flow (No kinks)									
7	Performed routine meatal care									
8	Uro bag emptied once it's 2/3 full.									
9	Replacement of urinary bag/ Indwelling catheter as per policy / clinically indicated.									
10	Prompt removal of catheter when no longer needed									
	Employee code/ Signature of the nurse									

Surgical site infection (SSI) care bundle.

Sl. No	Check list	Date:			Date:			Date:		
		M	E	N	M	E	N	M	E	N
0	Admitted with the history of previous surgery: Yes [] If yes, mention date of surgery : No []	Time	Time	Time	Time	Time	Time	Time	Time	Time
1	Hand hygiene performed before and after handling the surgical wound.									
2	Wound assessment done each shift .									
3	Aseptic techniques during wound dressing.									
4	Surgical dressing is clean and dry.									
5	Antibiotic followed as prescribed.									
6	Maintained normal blood glucose level.									
7	Maintained good nutritional status.									
8	Early mobilization and ambulation.									
9	No sign of redness, swelling, pain, purulent discharge, fever or chill.									
10	Health education given to the patient and family.									
	Employee code/ Signature of the nurse.									