Class 14: RNASeq Mini-Project

Nathalie Huang (PID: A19134713)

Table of contents

Background . . . 1
Data Import . . . 1
  Clean up (data tidying) . . . 2
DESeq Analysis . . . 3
  Setting up the DESeq object . . . 3
  Running DESeq . . . 6
  Getting results . . . 6
Volcano Plot . . . 7
Add Annotation . . . 9
Pathway Analysis . . . 11
  KEGG . . . 11
  GO . . . 20
  Reactome . . . 22

Background

The data for today's mini-project comes from a knock-down study of an important HOX gene.

Data Import

```
countData <- read.csv("GSE37704_featurecounts.csv", row.names=1)
head(countData)
```

```
                 length SRR493366 SRR493367 SRR493368 SRR493369 SRR493370
ENSG00000186092    918         0         0         0         0         0
```

```
ENSG00000279928    718         0         0         0         0         0
ENSG00000279457   1982        23        28        29        29        28
ENSG00000278566    939         0         0         0         0         0
ENSG00000273547    939         0         0         0         0         0
ENSG00000187634   3214       124       123       205       207       212
                 SRR493371
ENSG00000186092         0
ENSG00000279928         0
ENSG00000279457        46
ENSG00000278566         0
ENSG00000273547         0
ENSG00000187634       258
```

```
colData <- read.csv("GSE37704_metadata.csv", row.names=1)
head(colData)
```

```
              condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

Clean up (data tidying)

Q. Complete the code below to remove the troublesome first column from countData

```
countData <- as.matrix(countData[, 2:ncol(countData)])
head(countData)
```

```
                 SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000186092         0         0         0         0         0         0
ENSG00000279928         0         0         0         0         0         0
ENSG00000279457        23        28        29        29        28        46
ENSG00000278566         0         0         0         0         0         0
ENSG00000273547         0         0         0         0         0         0
ENSG00000187634       124       123       205       207       212       258
```

Q. Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns).

```r
countData <- countData[rowSums(countData) > 0, ]
head(countData)
```

```
                SRR493366 SRR493367 SRR493368 SRR493369 SRR493370 SRR493371
ENSG00000279457        23        28        29        29        28        46
ENSG00000187634       124       123       205       207       212       258
ENSG00000188976      1637      1831      2383      1226      1326      1504
ENSG00000187961       120       153       180       236       255       357
ENSG00000187583        24        48        65        44        48        64
ENSG00000187642         4         9        16        14        16        16
```

DESeq Analysis

Setting up the DESeq object

```r
library(DESeq2)
```

```
Loading required package: S4Vectors

Loading required package: stats4

Loading required package: BiocGenerics

Loading required package: generics


Attaching package: 'generics'

The following objects are masked from 'package:base':

    as.difftime, as.factor, as.ordered, intersect, is.element, setdiff,
    setequal, union


Attaching package: 'BiocGenerics'
```

```
The following objects are masked from 'package:stats':

    IQR, mad, sd, var, xtabs

The following objects are masked from 'package:base':

    anyDuplicated, aperm, append, as.data.frame, basename, cbind,
    colnames, dirname, do.call, duplicated, eval, evalq, Filter, Find,
    get, grep, grepl, is.unsorted, lapply, Map, mapply, match, mget,
    order, paste, pmax, pmax.int, pmin, pmin.int, Position, rank,
    rbind, Reduce, rownames, sapply, saveRDS, table, tapply, unique,
    unsplit, which.max, which.min


Attaching package: 'S4Vectors'

The following object is masked from 'package:utils':

    findMatches

The following objects are masked from 'package:base':

    expand.grid, I, unname

Loading required package: IRanges

Loading required package: GenomicRanges

Loading required package: Seqinfo

Loading required package: SummarizedExperiment

Loading required package: MatrixGenerics

Loading required package: matrixStats


Attaching package: 'MatrixGenerics'
```

```
The following objects are masked from 'package:matrixStats':

    colAlls, colAnyNAs, colAnys, colAvgsPerRowSet, colCollapse,
    colCounts, colCummaxs, colCummins, colCumprods, colCumsums,
    colDiffs, colIQRDiffs, colIQRs, colLogSumExps, colMadDiffs,
    colMads, colMaxs, colMeans2, colMedians, colMins, colOrderStats,
    colProds, colQuantiles, colRanges, colRanks, colSdDiffs, colSds,
    colSums2, colTabulates, colVarDiffs, colVars, colWeightedMads,
    colWeightedMeans, colWeightedMedians, colWeightedSds,
    colWeightedVars, rowAlls, rowAnyNAs, rowAnys, rowAvgsPerColSet,
    rowCollapse, rowCounts, rowCummaxs, rowCummins, rowCumprods,
    rowCumsums, rowDiffs, rowIQRDiffs, rowIQRs, rowLogSumExps,
    rowMadDiffs, rowMads, rowMaxs, rowMeans2, rowMedians, rowMins,
    rowOrderStats, rowProds, rowQuantiles, rowRanges, rowRanks,
    rowSdDiffs, rowSds, rowSums2, rowTabulates, rowVarDiffs, rowVars,
    rowWeightedMads, rowWeightedMeans, rowWeightedMedians,
    rowWeightedSds, rowWeightedVars

Loading required package: Biobase

Welcome to Bioconductor

    Vignettes contain introductory material; view with
    'browseVignettes()'. To cite Bioconductor, see
    'citation("Biobase")', and for packages 'citation("pkgname")'.


Attaching package: 'Biobase'

The following object is masked from 'package:MatrixGenerics':

    rowMedians

The following objects are masked from 'package:matrixStats':

    anyMissing, rowMedians
```

```
dds = DESeqDataSetFromMatrix(countData= countData,
                             colData= colData,
                             design=~ condition)
```

```
Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in
design formula are characters, converting to factors
```

```r
dds = DESeq(dds)
```

```
estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing
```

```r
dds
```

```
class: DESeqDataSet
dim: 15975 6
metadata(1): version
assays(4): counts mu H cooks
rownames(15975): ENSG00000279457 ENSG00000187634 ... ENSG00000276345
  ENSG00000271254
rowData names(22): baseMean baseVar ... deviance maxCooks
colnames(6): SRR493366 SRR493367 ... SRR493370 SRR493371
colData names(2): condition sizeFactor
```

Running DESeq

```r
res = results(dds)
```

Getting results

> Q. Call the summary() function on your results to get a sense of how many genes are up or down-regulated at the default 0.1 p-value cutoff.

```r
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
LFC > 0 (up)       : 4349, 27%
LFC < 0 (down)     : 4396, 28%
outliers [1]       : 0, 0%
low counts [2]     : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```

Volcano Plot

```r
library(ggplot2)
```

```r
ggplot(res) +
  aes(x = log2FoldChange,
      y = -log10(padj)) +
  geom_point()
```

```
Warning: Removed 1237 rows containing missing values or values outside the scale range
(`geom_point()`).
```



Q. Improve this plot by completing the below code, which adds color, axis labels and cutoff lines:

```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res))

# Color blue the genes with fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "blue"

# Color gray those with adjusted p-value more than 0.01
mycols[ res$padj > 0.01 ] <- "gray"

ggplot(res) +
  aes(x=log2FoldChange,
      y=-log10(padj)) +
  geom_point(color = mycols) +
  xlab("Log2(FoldChange)") +
  ylab("-Log(P-value)") +
  geom_vline(xintercept = c(-2,2)) +
  geom_hline(yintercept = -log10(0.01))
```

```
Warning: Removed 1237 rows containing missing values or values outside the scale range
(`geom_point()`).
```



Add Annotation

Q. Use the mapIDs() function multiple times to add SYMBOL, ENTREZID and GENENAME annotation to our results by completing the code below.

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

```
 [1] "ACCNUM"       "ALIAS"        "ENSEMBL"      "ENSEMBLPROT"  "ENSEMBLTRANS"
 [6] "ENTREZID"     "ENZYME"       "EVIDENCE"     "EVIDENCEALL"  "GENENAME"
[11] "GENETYPE"     "GO"           "GOALL"        "IPI"          "MAP"
[16] "OMIM"         "ONTOLOGY"     "ONTOLOGYALL"  "PATH"         "PFAM"
[21] "PMID"         "PROSITE"      "REFSEQ"       "SYMBOL"       "UCSCKG"
[26] "UNIPROT"
```

```r
res$symbol = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="SYMBOL",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```r
res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```r
res$name =   mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="GENENAME",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```r
head(res, 10)
```

```
log2 fold change (MLE): condition hoxa1 kd vs control sirna
Wald test p-value: condition hoxa1 kd vs control sirna
DataFrame with 10 rows and 9 columns
                   baseMean log2FoldChange     lfcSE       stat      pvalue
                  <numeric>      <numeric> <numeric>  <numeric>   <numeric>
ENSG00000279457   29.913579      0.1792571 0.3248216   0.551863 5.81042e-01
ENSG00000187634  183.229650      0.4264571 0.1402658   3.040350 2.36304e-03
ENSG00000188976 1651.188076     -0.6927205 0.0548465 -12.630158 1.43990e-36
ENSG00000187961  209.637938      0.7297556 0.1318599   5.534326 3.12428e-08
ENSG00000187583   47.255123      0.0405765 0.2718928   0.149237 8.81366e-01
ENSG00000187642   11.979750      0.5428105 0.5215598   1.040744 2.97994e-01
ENSG00000188290  108.922128      2.0570638 0.1969053  10.446970 1.51282e-25
ENSG00000187608  350.716868      0.2573837 0.1027266   2.505522 1.22271e-02
```

```
ENSG00000188157 9128.439422      0.3899088 0.0467163   8.346304 7.04321e-17
ENSG00000237330    0.158192      0.7859552 4.0804729   0.192614 8.47261e-01
                      padj      symbol      entrez                   name
                  <numeric> <character> <character>            <character>
ENSG00000279457 6.86555e-01          NA          NA                     NA
ENSG00000187634 5.15718e-03      SAMD11      148398 sterile alpha motif ..
ENSG00000188976 1.76549e-35       NOC2L       26155 NOC2 like nucleolar ..
ENSG00000187961 1.13413e-07      KLHL17      339451 kelch like family me..
ENSG00000187583 9.19031e-01     PLEKHN1       84069 pleckstrin homology ..
ENSG00000187642 4.03379e-01       PERM1       84808 PPARGC1 and ESRR ind..
ENSG00000188290 1.30538e-24        HES4       57801 hes family bHLH tran..
ENSG00000187608 2.37452e-02       ISG15        9636 ISG15 ubiquitin like..
ENSG00000188157 4.21963e-16        AGRN      375790                  agrin
ENSG00000237330          NA      RNF223      401934 ring finger protein ..
```

Q. Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

```
res = res[order(res$pvalue),]
write.csv(res, file="deseq_results.csv")
```

Pathway Analysis

KEGG

```
#Run in R console
#BiocManager::install( c("pathview", "gage", "gageData") )
```

```
library(pathview)
```

```
##############################################################################
Pathview is an open source software package distributed under GNU General
Public License version 3 (GPLv3). Details of GPLv3 is available at
http://www.gnu.org/licenses/gpl-3.0.html. Particullary, users are required to
formally cite the original Pathview paper (not just mention it) in publications
or products. For details, do citation("pathview") within R.

The pathview downloads and uses KEGG data. Non-academic uses may require a KEGG
license agreement (details at http://www.kegg.jp/kegg/legal.html).
##############################################################################
```

```r
library(gage)
```

```r
library(gageData)

data(kegg.sets.hs)
data(sigmet.idx.hs)

kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
[1] "10"   "1544" "1548" "1549" "1553" "7498" "9"

$`hsa00983 Drug metabolism - other enzymes`
 [1] "10"     "1066"   "10720"  "10941"  "151531" "1548"   "1549"   "1551"
 [9] "1553"   "1576"   "1577"   "1806"   "1807"   "1890"   "221223" "2990"
[17] "3251"   "3614"   "3615"   "3704"   "51733"  "54490"  "54575"  "54576"
[25] "54577"  "54578"  "54579"  "54600"  "54657"  "54658"  "54659"  "54963"
[33] "574537" "64816"  "7083"   "7084"   "7172"   "7363"   "7364"   "7365"
[41] "7366"   "7367"   "7371"   "7372"   "7378"   "7498"   "79799"  "83549"
[49] "8824"   "8833"   "9"      "978"

$`hsa00230 Purine metabolism`
  [1] "100"    "10201"  "10606"  "10621"  "10622"  "10623"  "107"    "10714"
  [9] "108"    "10846"  "109"    "111"    "11128"  "11164"  "112"    "113"
 [17] "114"    "115"    "122481" "122622" "124583" "132"    "158"    "159"
 [25] "1633"   "171568" "1716"   "196883" "203"    "204"    "205"    "221823"
 [33] "2272"   "22978"  "23649"  "246721" "25885"  "2618"   "26289"  "270"
 [41] "271"    "27115"  "272"    "2766"   "2977"   "2982"   "2983"   "2984"
 [49] "2986"   "2987"   "29922"  "3000"   "30833"  "30834"  "318"    "3251"
 [57] "353"    "3614"   "3615"   "3704"   "377841" "471"    "4830"   "4831"
 [65] "4832"   "4833"   "4860"   "4881"   "4882"   "4907"   "50484"  "50940"
 [73] "51082"  "51251"  "51292"  "5136"   "5137"   "5138"   "5139"   "5140"
 [81] "5141"   "5142"   "5143"   "5144"   "5145"   "5146"   "5147"   "5148"
 [89] "5149"   "5150"   "5151"   "5152"   "5153"   "5158"   "5167"   "5169"
 [97] "51728"  "5198"   "5236"   "5313"   "5315"   "53343"  "54107"  "5422"
[105] "5424"   "5425"   "5426"   "5427"   "5430"   "5431"   "5432"   "5433"
[113] "5434"   "5435"   "5436"   "5437"   "5438"   "5439"   "5440"   "5441"
```

```
[121] "5471"   "548644" "55276"  "5557"   "5558"   "55703"  "55811"  "55821"
[129] "5631"   "5634"   "56655"  "56953"  "56985"  "57804"  "58497"  "6240"
[137] "6241"   "64425"  "646625" "654364" "661"    "7498"   "8382"   "84172"
[145] "84265"  "84284"  "84618"  "8622"   "8654"   "87178"  "8833"   "9060"
[153] "9061"   "93034"  "953"    "9533"   "954"    "955"    "956"    "957"
[161] "9583"   "9615"
```

```r
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
     1266     54855      1465      2034      2150      6659
-2.422719  3.201955 -2.313738 -1.888019  3.344508  2.392288
```

```r
keggres = gage(foldchanges, gsets=kegg.sets.hs)
```

```r
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

```r
head(keggres$less)
```

```
                                        p.geomean stat.mean        p.val
hsa04110 Cell cycle                   8.995727e-06 -4.378644 8.995727e-06
hsa03030 DNA replication              9.424076e-05 -3.951803 9.424076e-05
hsa03013 RNA transport                1.375901e-03 -3.028500 1.375901e-03
hsa03440 Homologous recombination     3.066756e-03 -2.852899 3.066756e-03
hsa04114 Oocyte meiosis               3.784520e-03 -2.698128 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 8.961413e-03 -2.405398 8.961413e-03
                                            q.val set.size         exp1
hsa04110 Cell cycle                   0.001448312      121 8.995727e-06
hsa03030 DNA replication              0.007586381       36 9.424076e-05
hsa03013 RNA transport                0.073840037      144 1.375901e-03
hsa03440 Homologous recombination     0.121861535       28 3.066756e-03
hsa04114 Oocyte meiosis               0.121861535      102 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 0.212222694       53 8.961413e-03
```

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

```
'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14

Info: Writing image file hsa04110.pathview.png
```




```
pathview(gene.data=foldchanges, pathway.id="hsa04110", kegg.native=FALSE)
```

```
'select()' returned 1:1 mapping between keys and columns

Warning: reconcile groups sharing member nodes!
```

```
     [,1] [,2]
[1,] "9"  "300"
[2,] "9"  "306"
```

```
Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14
```

```
Info: Writing image file hsa04110.pathview.pdf
```

```r
# Focus on top 5 upregulated pathways
keggrespathways <- rownames(keggres$greater)[1:5]

# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```r
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14
```

```
Info: Writing image file hsa04640.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14
```

```
Info: Writing image file hsa04630.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14
```

```
Info: Writing image file hsa00140.pathview.png
```

```
'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14

Info: Writing image file hsa04142.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/nathaliehuang/Desktop/bimm143/class14

Info: Writing image file hsa04330.pathview.png
```


HEMATOPOIETIC CELL LINEAGE
Lymphoid Related Dendritic cell
γδ T cell
CD8 T cell
CD4 T cell
Regulatory T cell
NKT cell
Thymus
Pro T cell (DN2)
DN3
DN4
Intermediate single-positive cell (ISP)
Double-positive cell (DP)
NK cell Precursor
NK cell
Lymphoid stem cell, Double-negative cell (DN1)
Pro B Cell
Pre B I cell
Pre B II cell
Immature B cell
B Cell
Hematopoietic stem cell
CFU-Mast
Mast cell
CFU-Bas
Myeloblast
Basophilic Myelocyte
Basophil
CFU-E0
Eosinophilic Myelocyte
Eosinophil
CFU-M/DC
Myeloid Related Dendritic Cell
Monoblast
Promonocyte
Monocyte
Macrophage
Myeloid Stem Cell
CFU-GEMM
CFU-GM
CFU-G
Neutrophilic Myelocyte
Neutrophil
Bone marrow
BFU-E
CFU-E
Proerythroblast
Erythrocyte
BFU-MK
CFU-MK
Mega-karyocyte
Platelets
Data on KEGG graph
Rendered by Pathview


JAK-STAT SIGNALING PATHWAY
-1
0
1
ECS complex
Ubiquitin mediated proteolysis
Cytokine-cytokine receptor interaction
TC-PTP
SHP1
STAM
-p
-p
STAT dimerization
Interleukins
IL2family
IL3family
IL6family
IL10family
IL12/23
Hormone
Interferons
IFN-I/III
IFN-II
Growth factors
GF
Receptor
JAK1/3
JAK2
JAK1/2
TYK2
JAK1/3
TYK2
JAK2
TYK2
JAK2
JAK1
TYK2
JAK1/2
JAK1/2
+p
STAT
STAT1/3/5/6
STAT5
STAT1/3/6
STAT3
STAT3/4
STAT3/5
STAT1/2
STAT1/3
STAT1/3
IRF9
TC-PTP
PIAS
-p
+u
CBP/P300
SLIM
Proteasome
DNA
CIS
SOCS
Bcl-2
MCL1
Bcl-XL
PIM1
c-Myc
CycD
p21
AOX
GFAP
Apoptosis
Anti-apoptosis
Cell cycle
Cell-cycle progression
Cell-cycle inhibition
Lipid metabolism
Differentiation
MAPK signaling pathway
SHP2
GRB
SOS
Ras
Raf
Proliferation
Differentiation
PI3K
AKT
mTOR
Cell cycle
Cell survival
PI3K-AKT signaling pathway
Data on KEGG graph
Rendered by Pathview


STEROID HORMONE BIOSYNTHESIS
-1
0
1
Steroid biosynthesis
Cholesterol
Cholesterol sulfate
20α-Hydroxy-cholesterol
22β-Hydroxy-cholesterol
20α,22β-Dihydroxy-cholesterol
21-Hydroxy-pregnenolone
11-Deoxy-corticosterone
Corticosterone
11-Dehydro-corticosterone
Aldosterone hemiacetal
Aldosterone
18-Hydroxy-corticosterone
5α-Dihydro-deoxycorticosterone
Allotetrahydro-deoxycorticosterone
11β,21-Dihydroxy-3,20-oxo-5β-pregnan-18-al
3α,11β,21-Trihydroxy-20-oxo-5β-pregnan-18-al
3α,21-Dihydroxy-5β-pregnane-11,20-dione
3α,20α,21-Trihydroxy-5β-pregnane-11-one
11β,21-Dihydroxy-5β-pregnane-3,20-dione
Tetrahydro-corticosterone
21-Hydroxy-5β-pregnane-3,11,20-trione
3α-Hydroxy-5β-pregnane-20-one
Pregnanediol
5β-Pregnane-3,20-dione
4-Methylpentanal
Pregnenolone
7α-Hydroxy-pregnenolone
20α-Hydroxy-progesterone
Progesterone
11β-Hydroxy-progesterone
Pregnenolone-sulfate
5α-Pregnan-20α-ol-3-one
5α-Pregnan-3,20-dione
3α-Hydroxy-5α-pregnan-20-one
5α-Pregnane-3α,20α-diol
C21-Sterolds
17α,20α-Dihydroxy-pregn-4-en-3-one
17α,20α-Dihydroxy-cholesterol
17α-Hydroxy-pregnenolone
17α-Hydroxy-progesterone
21-Deoxycortisol
5α-Pregnan-17α-ol-3,20-dione
5α-Pregnane-3α,17α-diol-20-one
11β,17α,21-Trihydroxy-5β-pregnane-3,20-dione
Urocortisol
Cortol
17α,21-Dihydroxy-pregnenolone
11-Deoxycortisol
Cortisol
17α,21-Dihydroxy-5β-pregnane-3,11,20-trione
Cortisone
Urocortisone
Cortolone
11β,17α,21-Trihydroxy-pregnenolone
11β-Hydroxy-testosterone
Dehydro-epiandrosterone
Dehydroepiandro-steron sulfate
11β-Hydroxyandrost-4-ene-3,17-dione
Adrenosterone
Estrone 3-sulfate
7α-Hydroxydehydro-epiandrosterone
5β-Androstane-3,17-dione
Etiocholan-3α-ol-17-one
Etiocholan-3α-ol-17-one 3-glucuronide
Estrone glucuronide
Estradiol-17α
2-Methoxyestrone-3-sulfate
16α-Hydroxyandrost-4-ene-3,17-dione
16α-Hydroxydehydro-epiandrosterone
3-Oxo-13,17-secoandrost-4-ene-17,13α-lactone
2-Hydroxyestrone
2-Methoxyestrone
2-Methoxyestrone-3-glucuronide
19-Hydroxyandrost-4-ene-3,17-dione
19-Oxoandrost-4-ene-3,17-dione
Androst-4-ene-3,17-dione
Estrone
16-α-Hydroxyestrone
Androsterone
5α-Androstane-3,17-dione
Androsterone-glucuronide
7α-Hydroxy-androstenedione
C18-Sterolds
7α-Hydroxy-testosterone
5α-Dihydro-testosterone
Androstan-3alpha,17beta-diol
Testosterone
3β,17β-Dihydroxy-androst-5-ene
19-Hydroxy-testosterone
19-Oxo-testosterone
Estradiol-17β
16-Glucuronide-estriol
Estriol
2-Methoxy-estradiol-17β
2-Hydroxy-estradiol-17β
2-Methoxy-estradiol-17β-3-glucuronide
6β-Hydroxy-estradiol-17β
2-Methoxy-estradiol-17β-3-sulfate
Estradiol-17β-3-glucuronide
C19-Sterolds
5β-Dihydro-testosterone
Testosterone glucuronide
Estradiol-17β-3-sulfate
Data on KEGG graph
Rendered by Pathview


LYSOSOME
Exocytosis
Phagocytosis
bacterium
exosome
Phagosome
phagosome
endolysosome
phagolysosome
MVB
Endocytosis
Endocytosis
clathrin coat
AP-2
clathrin
M6P
ATPaV1
early endosome
M6P receptor
late endosome
multivesicular body (MVB)
mannose
mature lysosomal hydrase
AP-3
BLOC3
cytosol
pH~7.2
ATP
ADP
pH~5.0
acid hydrolase
lysosomal membrane protein
lysosome
VAMP8
SNAP29
Stx17
AP-1
recycling endosome
Receptor recycling
Receptor-dependent transport
clathrin coated vesicle
clathrin
CLN5
CLN3
Retromer
Rab7
PLEKHM
autolysosome
degradation and recycling
Transport of synthesized lysosomal enzymes
AP-1
AP-3
GGAs
AP-4
AP-5
SPG11
SPG15
MCOLN1
Activation of lysosomal sulfatase precursor
FGE
lysosomal hydrase precursor
M6P receptor
MPR
sortilin
LIMP2
mannose
from ER
CLN8
CLN6
LYSET
GNPT
NAGPA
cis Golgi network
Golgi body
trans Golgi network
transport vesicle
Snare interactions in vesicular transport
autophagosome
LMP
membrane repair & proton homeostasis
Autophagy
Regulation of autophagy
mitochondria
phagophore
plasma membrane
-1
0
1
Lysosomal membrane proteins
pH~7.2
transporters
ion channels, transporters
MCOLN1
LAPTM4B
SLRAMP
CLN7
GLUT12
TPC
TMEM175
ATP
ADP
ATPaV1
H+
pH~5.0
acidification regulators
DMXL1
ROGDI
WDR7
NCOA7
major lysosomal membrane proteins
LAMP1
LIMP2
fusion
mucolipin
lipid
NPC1
ABCA3/5
VPS13C
ORPs
sugar
GLUT6
NPC2
Lysosomal acid hydrolases
proteases
cathepsins
pepsin
LGMN
TPP1
glycosidases
GLA
GLB
GAA
GBA
IDUA
NAGA
NAGLU
GALC
GUSB
FUCA1
HEXA/B
MANB
GALAM
HPH
HYAL1
sulfatases
ARS
GALNS
GNS
IDS
SGSH
lipases
LIPA
LYPLA2
nuclease
DNase II
phosphatase
ACP2
ACP5
sphingomyelinase
SMPD1
ceramidase
ASAH1
aspartylglucosaminidase
AGA
Other lysosomal enzymes and activators
saposin
GM2A
CLN1
SNAREs
VAMP7
VAMP8
GTPase
Rab7
ARL8
tethering factors
HOPS
BORC
motor adaptors
nucleoside
SLC29A3
amino acid
PAT
FHT
CTNS
sialic acid
sialin
peptide
ABCB9
enzymes
PIGM
PIGV
LALP1
HGSNAT
lysosome
signaling complexes
mTORC1
RagA/B
RagC/D
Ragulator
AMPK
sensors
SLC38A9
LYCHOS
transcription factors
MITF
TFEB
TFE3
mTOR signaling pathway
immunity
TLR2
TLR7/8
TLR3
LITAF
Toll-like receptor signaling pathway
Glycosaminoglycan degradation
Other glycan degradation
Data on KEGG graph
Rendered by Pathview


-1
0
1
NOTCH SIGNALING PATHWAY
Fringe
γ-Secretase complex
PSE2
PSEN
NCSTN
APH-1
ADAM17
S2
S3
Delta
Notch
Serrate
+u
Deltex
Itch
Numb
Dvl
NICD
(Notch intracellular domain)
Co-activator
MAML
HATs
SKIP
CSL
DNA
Hes1/5
Hey
PreTα
NRARP
Co-repressor
Hairless
SMRT
CtBP
Gro/TLE
CIR
SHARP
HDAC
ATXN1/L
Hes1
NRARP
Ras/MAPK
MAPK signaling pathway
Gene expression
Data on KEGG graph
Rendered by Pathview

GO

```r
data(go.sets.hs)
data(go.subs.hs)

gobpsets = go.sets.hs[go.subs.hs$BP]

gobpres = gage(foldchanges, gsets=gobpsets)

lapply(gobpres, head)
```

```
$greater
                                          p.geomean stat.mean        p.val
GO:0007156 homophilic cell adhesion       8.519724e-05  3.824205 8.519724e-05
GO:0002009 morphogenesis of an epithelium 1.396681e-04  3.653886 1.396681e-04
GO:0048729 tissue morphogenesis           1.432451e-04  3.643242 1.432451e-04
GO:0007610 behavior                       1.925222e-04  3.565432 1.925222e-04
GO:0060562 epithelial tube morphogenesis  5.932837e-04  3.261376 5.932837e-04
GO:0035295 tube development               5.953254e-04  3.253665 5.953254e-04
                                              q.val set.size         exp1
GO:0007156 homophilic cell adhesion       0.1951953      113 8.519724e-05
GO:0002009 morphogenesis of an epithelium 0.1951953      339 1.396681e-04
GO:0048729 tissue morphogenesis           0.1951953      424 1.432451e-04
GO:0007610 behavior                       0.1967577      426 1.925222e-04
GO:0060562 epithelial tube morphogenesis  0.3565320      257 5.932837e-04
GO:0035295 tube development               0.3565320      391 5.953254e-04

$less
                                            p.geomean stat.mean        p.val
GO:0048285 organelle fission             1.536227e-15 -8.063910 1.536227e-15
GO:0000280 nuclear division              4.286961e-15 -7.939217 4.286961e-15
GO:0007067 mitosis                       4.286961e-15 -7.939217 4.286961e-15
GO:0000087 M phase of mitotic cell cycle 1.169934e-14 -7.797496 1.169934e-14
GO:0007059 chromosome segregation        2.028624e-11 -6.878340 2.028624e-11
GO:0000236 mitotic prometaphase          1.729553e-10 -6.695966 1.729553e-10
                                                q.val set.size         exp1
GO:0048285 organelle fission             5.841698e-12      376 1.536227e-15
GO:0000280 nuclear division              5.841698e-12      352 4.286961e-15
GO:0007067 mitosis                       5.841698e-12      352 4.286961e-15
GO:0000087 M phase of mitotic cell cycle 1.195672e-11      362 1.169934e-14
GO:0007059 chromosome segregation        1.658603e-08      142 2.028624e-11
GO:0000236 mitotic prometaphase          1.178402e-07       84 1.729553e-10
```

```
$stats
                                          stat.mean     exp1
GO:0007156 homophilic cell adhesion        3.824205 3.824205
GO:0002009 morphogenesis of an epithelium  3.653886 3.653886
GO:0048729 tissue morphogenesis            3.643242 3.643242
GO:0007610 behavior                        3.565432 3.565432
GO:0060562 epithelial tube morphogenesis   3.261376 3.261376
GO:0035295 tube development                3.253665 3.253665
```

Reactome

```r
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```r
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quote=
```

> Q. What pathway has the most significant "Entities p-value"? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

Most significant "Entities p-value": Cell Cycle, Mitotic and Cell Cycle Yes, the most significant pathways listed match my previous KEGG results. The KEGG results identified Cell Cycle–related pathways as the most significant. Factors that could cause differences between the two methods are different gene background sizes, how broad the pathways are, and pvalue calculations.