

Number Theoretic Transform

Jonghyun Kim

Korea University

January 28, 2026


Table of Contents

Number Theoretic Transform - Basics


Number Theoretic Transform - Advanced

Number Theoretic Transform - Basics


Modular Exponentiation Using CRT



Polynomial Multiplication Using NTT




Radix-2 NTT Layer




$$2a_0(x) = \hat{a}_0(x) + \hat{a}_1(x)$$

$$2a_1(x) = (\hat{a}_0(x) - \hat{a}_1(x))\zeta^{-1}$$

Radix-2 NTT Structure (1)




Radix-2 NTT Structure (2)



Total : $\mathcal{O}(nk) = \mathcal{O}(n \log n)$

Radix-2 Inverse NTT Structure



Condition for applying NTT (1)

- $\mathbb{Z}_q[x]/\langle x^n + 1 \rangle$, where n is a power of 2
 - ζ : primitive $2n$ -th root of unity modulo q
 - $\zeta^i \not\equiv 1 \pmod{q}$ for $i \in [1, 2n - 1]$
 - $\zeta^{2n} \equiv 1 \pmod{q}$
 - **Fact 1:** $\zeta^n + 1 \equiv 0 \pmod{q}$
 - $\zeta^{2n} - 1 \equiv (\zeta^n + 1)(\zeta^n - 1) \equiv 0 \pmod{q}$
 - By the definition of ζ , $\zeta^n - 1 \not\equiv 0 \pmod{q} \Rightarrow \zeta^n + 1 \equiv 0 \pmod{q}$
 - **Fact 2:** $\zeta^i \not\equiv \zeta^j \pmod{q}$ for $i, j \in [1, 2n]$ with $i \neq j$
 - If there exist i, j with $1 \leq i < j \leq 2n$ such that $\zeta^i \equiv \zeta^j \pmod{q}$, then $\zeta^{j-i} \equiv 1 \pmod{q}$, a contradiction since $j - i \in [1, 2n - 1]$.

Condition for applying NTT (2)

- $\mathbb{Z}_q[x]/\langle x^n + 1 \rangle$, where n is a power of 2
- $\boxed{\zeta: \text{primitive } 2n\text{-th root of unity modulo } q}$
 - $\zeta^i \not\equiv 1 \pmod{q}$ for $i \in [1, 2n - 1]$
 - $\zeta^{2n} \equiv 1 \pmod{q}$
- **Fact 1:** $\zeta^n + 1 \equiv 0 \pmod{q}$
- **Fact 2:** $\zeta^i \not\equiv \zeta^j \pmod{q}$ for $i, j \in [1, 2n]$ with $i \neq j$
- $$\begin{aligned} x^n + 1 &= x^n - \zeta^n && \because \text{Fact 1} \\ &= (x^{n/2} - \zeta^{n/2})(x^{n/2} + \zeta^{n/2}) \\ &= (x^{n/2} - \zeta^{n/2})(x^{n/2} - \zeta^{3n/2}) && \because \text{Fact 1} \\ &= (x^{n/4} - \zeta^{n/4})(x^{n/4} + \zeta^{n/4})(x^{n/4} - \zeta^{3n/4})(x^{n/4} + \zeta^{3n/4}) \\ &= (x - \zeta)(x - \zeta^3)(x - \zeta^5) \cdots (x - \zeta^{2n-1}) \end{aligned}$$

All the factors are distinct (\because **Fact 2**) \Rightarrow they are relatively prime.

Finding $2n$ -th root of unity modulo q

1. Find a generator $g \in \mathbb{Z}_q^* = \{1, \dots, q-1\}$.
 - $g^i \not\equiv 1 \pmod{q}$ for $i \in [1, q-2]$
 - $g^{q-1} \equiv 1 \pmod{q}$

$$\{1, \dots, q-1\} = \{g^1, \dots, g^{q-1}\}$$

2. Compute the integer $k = \frac{q-1}{2n}$, assuming that $2n|q-1$.
3. Output $\zeta = g^k \pmod{q}$ as a primitive $2n$ -th root of unity.
 - $\zeta^i \equiv (g^k)^i \not\equiv 1$ for $i \in [1, 2n-1]$ by the definition of g
 - $\zeta^{2n} \equiv (g^k)^{2n} \equiv g^{2nk} \equiv g^{q-1} \equiv 1 \pmod{q}$

Generator Test for $g \in \mathbb{Z}_q^* = \{1, \dots, q-1\}$ (1)

○ \mathbb{Z}_{13}^*

– $g^{q-1} \equiv g^{2^2 \cdot 3} \equiv 1 \pmod{13}$

$\Rightarrow g^{\{1,2,4,6\}} \stackrel{?}{\not\equiv} 1 \pmod{13} \Rightarrow g^{\{4,6\}} \stackrel{?}{\not\equiv} 1 \pmod{13}$

i	1	2	3	4	5	6	7	8	9	10	11	12
1^i	1	1	1	1	1	1	1	1	1	1	1	1
2^i	2	4	8	3	6	12	11	9	5	10	7	1
3^i	3	9	1	3	9	1	3	9	1	3	9	1
4^i	4	3	12	9	10	1	4	3	12	9	10	1
5^i	5	12	8	1	5	12	8	1	5	12	8	1
6^i	6	10	8	9	2	12	7	3	5	4	11	1
7^i	7	10	5	9	11	12	6	3	8	4	2	1
8^i	8	12	5	1	8	12	5	1	8	12	5	1
9^i	9	3	1	9	3	1	9	3	1	9	3	1
10^i	10	9	12	3	4	1	10	9	12	3	4	1
11^i	11	4	5	3	7	12	2	9	8	10	6	1
12^i	12	1	12	1	12	1	12	1	12	1	12	1

Generator Test for $g \in \mathbb{Z}_q^* = \{1, \dots, q-1\}$ (1)

1. Factorize $q-1$ as $q-1 = p_1^{r_1} \cdots p_\ell^{r_\ell}$
 - p_i are distinct primes.
2. For each $i \in \{1, \dots, \ell\}$:
 - If $g^{(q-1)/p_i} \equiv 1 \pmod{q}$, then return “ g is not a generator.”
3. Return “ g is a generator.”

Finding prime numbers q such that $2n|q - 1$ (SageMath)

```
def find_ntt_prime(n, bits):  
    qs = [];  
    k = 1;  
  
    while True:  
        q = 2*n*k+1;  
        if q > 2^bits:  
            break;  
        if q in Primes():  
            qs.append(q);  
        k += 1;  
    return qs;
```


Finding generators g of \mathbb{Z}_q^* (SageMath)

```
def find_generator(q):  
    Zq = IntegerModRing(q);  
    gs = range(1,q);  
  
    for x in list(factor(q-1)):  
        p = x[0];  
        t = [];  
        for g in gs:  
            if Zq(g)^((q-1)/p) != 1:  
                t.append(Zq(g));  
        gs = t;  
  
    return gs;
```


Finding primitive $2n$ -th root of unity modulo q (SageMath)

```
def find_w(q):  
    k = Integer((q-1)/(2*n));  
    ws = [g^k for g in find_generator(q)];  
    return sorted(list(set(ws)));
```


Generating Precomputation Table (1)



Generating Precomputation Table (2)

$$\begin{aligned}\hat{a}_0(x) &= a_0(x) + a_1(x)\zeta \\ \hat{a}_1(x) &= a_0(x) - a_1(x)\zeta \\ 2a_0(x) &= \hat{a}_0(x) + \hat{a}_1(x) \\ 2a_1(x) &= (\hat{a}_0(x) - \hat{a}_1(x))\zeta^{-1}\end{aligned}$$

$$\zeta^n \equiv -1 \pmod{q}$$



$$\zeta_i \zeta_{9-i} \equiv \zeta^n \equiv -1 \pmod{q} \Rightarrow \zeta_i^{-1} \equiv -\zeta_{9-i} \pmod{q}$$

Generating Precomputation Table (3) (SageMath)

```
level = Integer(log(n,2));

zetas = [];

tree = zero_matrix(ZZ,level+1,1 << level);
tree[0,0] = n;

for l in range(level):
    for i in range(1 << l):
        tree[l+1,2*i] = tree[l, i] / 2;
        tree[l+1,2*i+1] = tree[l+1,2*i] + n;

    zetas.append(Zq(w)^tree[l+1,2*i]);
```

Signed Montgomery Reduction

- Signed Montgomery Reduction [3]

- For $0 < q < \frac{\beta}{2}$:

$$\hat{a} = \text{Mont}(a) \equiv a\beta^{-1} \pmod{q}$$

- Constraints on a : $-\frac{\beta}{2}q \leq a < \frac{\beta}{2}q$
- Range of \hat{a} : $-q < \hat{a} < q$

- Montgomery Reduction for Multiplication


- Transform to Montgomery Form

- $\hat{a} = \text{Mont}(a \cdot (\beta^2 \bmod q)) \equiv a\beta \pmod{q}$
- $\hat{b} = \text{Mont}(b \cdot (\beta^2 \bmod q)) \equiv b\beta \pmod{q}$

- Montgomery Multiplication

- $\hat{a}\hat{b} \equiv ab\beta^2 \pmod{q}$
- $\text{Mont}(\hat{a}\hat{b}) \equiv \text{Mont}(ab\beta^2) \equiv ab\beta \pmod{q}$

NTT Using Montgomery Reduction



$$\text{Mont}(a_1(x) \times \underbrace{\zeta \beta \bmod q}_{\text{pre-computation}}) = a_1(x)\zeta \bmod q$$

Number Theoretic Transform - Advanced

Variants of NTT (1)

- Incomplete NTT

- $\mathbb{Z}_q[x]/\langle x^n + 1 \rangle \approx \prod_{i=1}^{n/2} \mathbb{Z}_q[x]/\langle x^2 - \zeta_i \rangle$
 - $n = 2^m$ for some $m \in \mathbb{N}$
 - ζ : primitive $(2n/2)$ -th root of unity modulo q
 - q : prime number with $q = (2n/2) \cdot k + 1$ for some $k \in \mathbb{N}$
 - Supports a larger set of modulus q for the NTT

- Example

- Complete NTT in CRYSTAL-KYBER (Round 1)
 - $n = 256, q = 7681$
- Incomplete NTT in CRYSTAL-KYBER (Round 2 & 3)
 - $n = 256, q = 3329$

Variants of NTT (2)

- Radix-2 NTT Layer for Cyclotomic Trinomial [2]

- $\mathbb{Z}_q[x]/\langle x^n - x^{n/2} + 1 \rangle \approx \prod_{i=1}^2 \mathbb{Z}_q[x]/\langle x^{n/2} - \zeta_i \rangle$
 - $n = 2^a 3^b$ for some $a, b \in \mathbb{N}$

- Radix-3 NTT Layer [1]

- $\mathbb{Z}_q[x]/\langle x^n - \zeta^3 \rangle \approx \prod_{i=1}^3 \mathbb{Z}_q[x]/\langle x^{n/3} - \zeta_i \rangle$
 - $n = 2^a 3^b$ for some $a, b \in \mathbb{N}$

Table: Combinations of NTT layers in NTRU+


n	q	Radix-2 for CT	Radix-3	Radix-2	d	ζ	$\ell = 3n/d$
576	3457	1	2	3	4	81	432
768	3457	1	1	5	4	22	576
864	3457	1	2	4	3	9	864
1152	3457	1	2	4	4	9	864

ζ : primitive ℓ -th root of unity modulo q

Radix-2 NTT Layer for Cyclotomic Trinomial (1)

- $R_q = \mathbb{Z}_q[x] / \langle x^n - x^{n/2} + 1 \rangle$
 - ψ : primitive 6-th root of unity modulo q
 - $\psi^i \not\equiv 1 \pmod{q}$ for $i \in [1, 5]$
 - $\psi^6 \equiv 1 \pmod{q}$
 - **Fact 1:** $\psi^2 - \psi + 1 \equiv 0 \pmod{q}$
 - $\psi^6 - 1 \equiv (\psi^3 - 1)(\psi + 1)(\psi^2 - \psi + 1) \equiv 0 \pmod{q}$
 - By the definition of ψ , $\psi^2 - \psi + 1 \equiv 0 \pmod{q}$
 - **Fact 1-1:** $\psi^3 + 1 \equiv (\psi + 1)(\psi^2 - \psi + 1) \equiv 0 \pmod{q}$
 - **Fact 2:** $x^2 - x + 1 = (x - \psi)(x - \psi^5)$
 - $(x - \psi)(x - \psi^5) \equiv x^2 - (\psi + \psi^5) + \psi^6 \pmod{q}$
 - **Fact 2-1:** $\psi + \psi^5 \equiv \psi - \psi^2 \equiv 1 \pmod{q}$
 - $\psi^6 \equiv 1 \pmod{q}$
 - $x^n - x^{n/2} + 1 = (x^{n/2} - \psi)(x^{n/2} - \psi^5)$ (\because **Fact 2**)


Radix-2 NTT Layer for Cyclotomic Trinomials (2)



Radix-3 NTT Layer (1)

- $R_q = \mathbb{Z}_q[x] / \langle x^3 - \zeta^3 \rangle$
 - ω : primitive 3-th root of unity modulo q
 - $\omega^i \not\equiv 1 \pmod{q}$ for $i \in [1, 2]$
 - $\omega^3 \equiv 1 \pmod{q}$
 - **Fact 1:** $\omega^2 + \omega + 1 \equiv 0 \pmod{q}$
 - $\omega^3 - 1 \equiv (\omega - 1)(\omega^2 + \omega + 1) \equiv 0 \pmod{q}$
 - By the definition of ω , $\omega^2 + \omega + 1 \equiv 0 \pmod{q}$
 - **Fact 2:** $x^3 - \zeta^3 = (x - \alpha)(x - \beta)(x - \gamma)$
 - $\alpha = \zeta, \beta = \zeta\omega, \gamma = \zeta\omega^2$
 - $(x - \alpha)(x - \beta)(x - \gamma) = x^3 - (\alpha + \beta + \gamma)x^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)x - \alpha\beta\gamma$
 - $\alpha + \beta + \gamma \equiv \zeta(1 + \omega + \omega^2) \equiv 0 \pmod{q}$ (\because **Fact 1**)
 - $\alpha\beta + \beta\gamma + \gamma\alpha \equiv \zeta(\omega + \omega^3 + \omega^2)$
 $\equiv \zeta(1 + \omega + \omega^2) \equiv 0 \pmod{q}$ (\because **Fact 1**)
 - $\alpha\beta\gamma \equiv \zeta^3\omega^3 \equiv \zeta^3 \pmod{q}$

Radix-3 NTT Layer (2)



Radix-3 NTT layer (3)

- NTT


$$\begin{pmatrix} \hat{a}_0(x) \\ \hat{a}_1(x) \\ \hat{a}_2(x) \end{pmatrix} = \begin{pmatrix} 1 & \alpha & \alpha^2 \\ 1 & \beta & \beta^2 \\ 1 & \gamma & \gamma^2 \end{pmatrix} \begin{pmatrix} a_0(x) \\ a_1(x) \\ a_2(x) \end{pmatrix}$$

- Inverse NTT

$$\begin{pmatrix} 1 & 1 & 1 \\ \alpha^{-1} & \beta^{-1} & \gamma^{-1} \\ \alpha^{-2} & \beta^{-2} & \gamma^{-2} \end{pmatrix} \begin{pmatrix} 1 & \alpha & \alpha^2 \\ 1 & \beta & \beta^2 \\ 1 & \gamma & \gamma^2 \end{pmatrix} \begin{pmatrix} a_0(x) \\ a_1(x) \\ a_2(x) \end{pmatrix} = 3 \begin{pmatrix} a_0(x) \\ a_1(x) \\ a_2(x) \end{pmatrix}$$

- $\alpha^2 + \beta^2 + \gamma^2 \equiv (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha) \equiv 0 \pmod{q}$
- $\alpha^{-1} + \beta^{-1} + \gamma^{-1} \equiv (\alpha\beta\gamma)^{-1}(\alpha\beta + \beta\gamma + \gamma\alpha) \equiv 0 \pmod{q}$
- $\alpha^{-2} + \beta^{-2} + \gamma^{-2}$
 $\equiv (\alpha^{-1} + \beta^{-1} + \gamma^{-1})^2 - 2(\alpha^{-1}\beta^{-1} + \beta^{-1}\gamma^{-1} + \gamma^{-1}\alpha^{-1})$
 $\equiv (\alpha^{-1} + \beta^{-1} + \gamma^{-1})^2 - 2\alpha^{-1}\beta^{-1}\gamma^{-1}(\alpha + \beta + \gamma) \equiv 0 \pmod{q}$


Radix-3 Inverse NTT Layer (1)



Example

- $\mathbb{Z}_q[x]/\langle x^{24} - x^{12} + 1 \rangle \approx \prod_{i=1}^{12} \mathbb{Z}_q[x]/\langle x^2 - \zeta_i \rangle$
- ζ : primitive 36-th root of unity modulo q .
 - $\zeta^{18} \equiv -1 \pmod{q}$, $\psi \equiv \zeta^6 \pmod{q}$, $\omega \equiv \zeta^{12} \pmod{q}$


$$\begin{aligned}
 - \quad x^{24} - x^{12} + 1 &= (x^{12} - \zeta^6)(x^{12} - \zeta^{30}) \\
 &= (x^4 - \zeta^2)(x^4 - \zeta^{14})(x^4 - \zeta^{26})(x^4 - \zeta^{10})(x^4 - \zeta^{22})(x^4 - \zeta^{34}) \\
 &= (x^2 - \zeta)(x^2 - \zeta^{19})(x^2 - \zeta^7)(x^2 - \zeta^{25})(x^2 - \zeta^{13})(x^2 - \zeta^{31}) \\
 &\quad (x^2 - \zeta^5)(x^2 - \zeta^{23})(x^2 - \zeta^{11})(x^2 - \zeta^{29})(x^2 - \zeta^{17})(x^2 - \zeta^{35})
 \end{aligned}$$



$$\begin{aligned}
 \zeta^2 \zeta^{10} &\equiv \zeta^{12} \equiv \omega \pmod{q} &\Rightarrow &\omega \zeta^{-2} \equiv \zeta^{10} \pmod{q} \\
 \zeta^4 \zeta^{20} &\equiv \zeta^{24} \equiv \omega^2 \pmod{q} &\Rightarrow &\omega^2 \zeta^{-4} \equiv \zeta^{20} \pmod{q}
 \end{aligned}$$

$$\zeta^1 \zeta^{17} \equiv \zeta^{18} \equiv -1 \pmod{q} \quad \Rightarrow \quad \zeta^{-1} \equiv -\zeta^{17} \pmod{q}$$

Radix-3 Inverse NTT Layer (2)



Helpful Scripts related with NTT

`https://github.com/ntruplus/ntt_for_ntruplus/`

Thank You for Your Attention!

Any Questions?

(I know this is shared online, so questions might be difficult.)

Contact: `yoswuk@korea.ac.kr`

References



Chenar Abdulla Hassan and Oğuz Yayla.

Radix-3 NTT-based polynomial multiplication for lattice-based cryptography.

[Cryptography ePrint Archive, Report 2022/726, 2022.](#)



Vadim Lyubashevsky and Gregor Seiler.

NTTRU: Truly fast NTRU using NTT.

IACR Transactions on Cryptographic Hardware and Embedded Systems, 2019(3):180–201, 2019.



Gregor Seiler.

Faster AVX2 optimized NTT multiplication for ring-LWE lattice cryptography.

[Cryptography ePrint Archive, Report 2018/039, 2018.](#)