An independent double-difference tomography implementation: synthetic reconstruction and limits of manuscript reproduction

Independent computational reproduction

October 2026

Abstract

Double-difference tomography uses paired seismic arrivals to reduce shared travel-time errors while estimating source locations and velocity jointly. Here we implement this inverse problem independently in C++ and test it against the written model of Zhang and Thurber (2003). The code combines variational ray tracing, trilinear velocity interpolation, joint linearization, and damped, smoothed least squares. We reconstruct the specified synthetic velocity profile and generate new event and station geometries. Consequently, the true-model panel is independently reproduced, whereas synthetic inversion panels are partial reproductions. The initial project instructions excluded Hayward observations. A later authorized acquisition supplies an independently selected, identified geometry for a partial Figure 4 counterpart. The catalog DD and absolute-data inversions have written numerical outputs. A partial velocity comparison is available; remaining methods and final diagnostics are pending. All numerical figures in this report use saved output from this implementation. The result is a reproducible method investigation, not a recovery of the published Hayward velocity model. The resumed work permits authoritative observational inputs. A later section records authoritative metadata investigation and ray-kernel corrections. The synthetic inverse solutions retain their original baseline provenance.

1 Scientific scope and evidence

The target paper is Zhang and Thurber (2003), *Bulletin of the Seismological Society of America* 93, 1875–1889, doi:10.1785/0120020190. Its written equations, experiment descriptions, and captions are the scientific specification. The publisher page returned HTTP 403; the complete manuscript transcription posted by an author on ResearchGate provided the text. The baseline calculations used no original model code, source observations, synthetic arrival files, plotting scripts, or figure files. No published plot was digitized. No published image appears in this report.

The initial exclusion of source observations had a direct scientific consequence. A physical forward model can generate new arrival times for a specified geometry, but it cannot recover the historical earthquake realization and picking errors from a prose description. The baseline therefore separated a reconstructible velocity specification, an independent synthetic inverse experiment, and unavailable field-data inversions. Later sections document the authorized authoritative-data retrieval and subsequent calculations. The panel audit preserves all eleven manuscript figures and records each remaining experimental difference. No supplement is listed for this article on the original SSA issue contents or its original abstract page. No supplementary figures were identified. The official issue contents marks supplements for other articles explicitly; `reference/specification_notes.md` records those checks.

2 Forward problem and joint inversion

Assume an isotropic, stationary medium and high-frequency geometric P rays. Coordinates are in km, velocity in km/s, and time in s; depth is positive downward. The predicted arrival of event i at station

k is

$$T_{ik} = \tau_i + \int_{\Gamma_{ik}} \frac{1}{v(\mathbf{x})}\, ds. \tag{1}$$

Here τ_i is origin time and Γ_{ik} is a stationary travel-time path with fixed endpoints. The calculation includes no reflected or converted phases, attenuation, or wave-amplitude effects. Paths remain inside the rectangular model domain. There are no periodic or reflecting boundary conditions.

Nodal velocities v_n are interpolated trilinearly. Write the update as $q_n = \delta \log(1/v_n)$, so that the nonlinear velocity update is $v_n^{\text{new}} = v_n \exp(-q_n)$. The absolute residual linearization is

$$r_{ik} \simeq \mathbf{g}_{ik} \cdot \delta\mathbf{x}_i + \delta\tau_i + \sum_n K_{ikn} q_n, \qquad K_{ikn} = v_n \int_{\Gamma_{ik}} \frac{w_n(\mathbf{x})}{v(\mathbf{x})^2}\, ds. \tag{2}$$

w_n is the interpolation weight. The source derivative $\mathbf{g}_{ik}$ is evaluated from the differentiated endpoint segment integral. Absolute arrivals constrain structure outside clusters and the common source shift.

For a measured event-pair difference d_{ijk} at a common station, use

$$r_{ijk}^{\text{DD}} = d_{ijk} - (T_{ik} - T_{jk}), \qquad A_{ijk}^{\text{DD}} = A_{ik} - A_{jk}. \tag{3}$$

Velocity sensitivities from both events enter the difference, subject to the 5 m sparsification rule described below. This includes changes in path anomalies with source location. Shared station timing errors cancel in differenced observations, whereas independent pick errors do not. Catalog differences share errors with each other and with absolute picks; our weighting follows the staged formulation without whitening this covariance.

The linear step minimizes a weighted residual norm with first-difference regularization of the final log slowness and damping of the column-scaled increment. A first-difference row divides the difference between adjacent nodes by their separation. Columns are normalized to unit Euclidean norm before damping rows are added. An independently written LSQR implementation solves the sparse system without forming normal equations. Absolute, catalog-difference, and precise-difference rows receive stage weights $(1, 0.1, 0.01)$, $(0.1, 1, 0.01)$, and $(0.001, 0.01, 1)$, respectively. Smoothing and damping choices are explicit in the configuration. After a rejected update, identical repetitions within the same stage are skipped, while every later weight stage is retained.

Each accepted nonlinear step must decrease the current weighted misfit plus roughness. Weights are frozen within that line search. Trial source locations are projected onto the grid domain; velocities remain positive by logarithmic parameterization. No published solution is used to choose an update. Sources at boundaries and failed updates are counted in the results because they can indicate model error or weak constraints.

2.1 Ray integration and numerical substitutions

The baseline ray tracer minimizes discretized Fermat time with limited-memory BFGS and Armijo backtracking, starting from a straight path. Each segment is split at grid-plane crossings and integrated with two-point Gaussian quadrature. This independent variational method replaces the paper's pseudo-bending implementation. It searches for a local stationary path, with no guarantee of finding every competing first-arrival branch. Search directions and convergence gradients respect active model boundaries. Near-coincident interior nodes are merged at a numerical tolerance far below ray spacing, with the quasi-Newton history reset after such topology changes. Independent rays are traced in parallel with OpenMP.

The reference configuration uses 1 km initial ray spacing and 180 ray iterations, 18 nonlinear iterations in three equal stages, smoothing weight 5, and scaled damping 0.1. These choices are not all specified by the paper. Distance and residual biweights, a 5 m differential path-coefficient cutoff, regularization scaling, line search, and the residual-scale estimator are fully defined in `METHODS.md` and

the source. A damping or smoothing number cannot be transferred between implementations without its scaling convention. Our equal-direction gradient penalty should therefore not be assumed identical to the original regularization.

3 Independent synthetic experiment

The true medium is constant with depth and along strike. Its cross-strike profile is 6 km/s for $x \leq 0$, falls linearly to 4 km/s at $x = 1$ km, remains 4 km/s through $x = 5$ km, rises linearly to 5 km/s at $x = 6$ km, and increases linearly to 6 km/s at $x = 38$ km. The paper describes sharp gradients over $x = 0$–1 and 5–6 km; linear interpolation across these two transitions is our explicit assumption. Figure 1 is generated from saved model sampling and reconstructs the written velocity specification with this assumption, without using published pixel values.

The inversion grid follows the coordinates in manuscript caption 2: $x = (-35, -15, 0, 2, 4, 6, 20, 35)$ km, $y = (-60, -40, -20, 0, 20, 40)$ km, and $z = (0, 3, 7, 11, 16)$ km. The forward grid adds the profile breakpoints $x = 1$ and 5 km. This prevents the data generator from using exactly the same coarse representation as the inverse problem.

We generate 120 sources and 32 surface stations independently with seed 2003. Half the sources sample a 12 km wide region around the trough, and half sample a broader 40 km interval; all sample the model volume in depth and along strike. Stations use a jittered surface array. Figure 2 shows the complete generated geometry. These are not the paper's Loma Prieta observations. The starting velocity is homogeneous at 5.5 km/s, substituting for the paper's referenced layered starting model, whose values are not tabulated in the target article. Independent starting source-coordinate errors have standard deviation 0.6 km, with an initial depth restriction of 0.5–15.5 km. Starting origin-time errors have standard deviation 0.05 s.

We round forward travel times to 0.01 s, add independent 0.04 s Gaussian pick noise and a station-constant uniform bias between −0.3 and 0.3 s. Catalog differences subtract these noisy picks. Precise synthetic differences subtract rounded forward times before added pick noise. This explicitly resolves an ambiguity in the text's noise ordering in favor of precise pair differences. Every event is observed at every station, and pairs within 20 km are selected. The saved arrival table separates exact time, rounding, station bias, pick noise, and final observed time. Subtracting two separately rounded arrivals permits differential rounding errors approaching 0.01 s. The paper quotes 0.005 s for its differential data; this discrepancy remains a limitation of our explicit rounding order.

3.1 Starting-model sensitivity

The target paper does not identify a side of the Dietz–Ellsworth model. USGS Open-File Report 93-578, Table 4, tabulates two gradient adaptations attributed to that work. We sample each named profile (LOM and LON) onto the same inversion-depth grid and repeat DD inversion using otherwise identical inputs. The adaptation footnote prevents identifying either as the confirmed target start. This comparison measures sensitivity to two written alternatives without inventing their values.

4 Validation and numerical results

All 32 implementation consistency checks passed. Across 32 deterministic source–receiver pairs, halving the ray segment spacing and increasing the iteration limit changed travel times by a median of 0.054 ms and a maximum of 0.892 ms. This check does not establish global ray optimality.

In the small known-model recovery check, the median three-dimensional location error decreased from 0.3035 to 0.0343 km; the final absolute-time RMS was 0.0034 s. Only 0.01 s time quantization remains in this check.

Start	Median 3D error	Mean $\lvert\Delta v\rvert$	Interior $\lvert\Delta v\rvert$	R_{abs}	R_{prec}
Homogeneous	0.8222	0.3050	0.2135	0.3526	0.4390
LOM	1.4316	0.4036	0.2589	0.3752	0.4370
LON	1.1469	0.3606	0.2444	0.3780	0.4348

Table 1: Starting-model sensitivity of the independent DD experiment. LOM and LON are written gradient adaptations in USGS OFR 93-578. All runs share geometry, true velocity, and observation noise. Median three-dimensional location errors are in km, mean absolute velocity errors are in km/s, and arrival residual RMS values are in s. The first velocity column uses all saved slices; the interior column restricts x to $[-15, 20]$ km, y to $[-40, 20]$ km, and depths to 3–11 km.




Figure 1: Manuscript Figure 1 reconstruction: true P velocity from the written sandwich profile, sampled at 7 km depth. The model is independent of depth; colors show velocity in km/s. All plotted samples are saved C++ output. The trough occupies $x = 1$–5 km and has velocity 4 km/s.




Figure 2: Partial reconstruction of manuscript Figure 2. Crosses are the written inversion grid, triangles are independent stations, and dots are independent true sources. The coordinates are experimental design choices, not estimates extracted from the published map.

Before the main inversions, the median three-dimensional source error was 1.0285 km. The homogeneous 5.50 km/s initial model had mean absolute velocity errors of 0.456 km/s over the full domain and 0.544 km/s in the interior, evaluated at the same saved slice samples as the recovered models.

The standard run ended with median three-dimensional source error 1.7069 km, 0 sources on a model boundary, 1 zero-update iterations, and nodal velocities ranging from 2.41 to 7.05 km/s. LSQR converged in 10 of 10 solves, with worst relative normal residual 8.35e-09. The dd run ended with median three-dimensional source error 0.8222 km, 0 sources on a model boundary, 1 zero-update iterations, and nodal velocities ranging from 3.74 to 6.41 km/s. LSQR converged in 18 of 18 solves, with worst relative normal residual 8.5e-09. The loc1d run ended with median three-dimensional source error 2.9856 km, 11 sources on a model boundary, 1 zero-update iterations, and nodal velocities ranging from 5.50 to 5.50 km/s. LSQR converged in 15 of 15 solves, with worst relative normal residual 9.64e-09. The loc3d run ended with median three-dimensional source error 0.1974 km, 14 sources on a model boundary, 0 zero-update iterations, and nodal velocities ranging from 4.00 to 6.00 km/s. LSQR converged in 18 of 18 solves, with worst relative normal residual 9.32e-09.

Relative to standard tomography, DD changed the full-domain mean absolute velocity error by -0.269 km/s and the median depth error by -0.996 km. Negative differences indicate improvement. These comparisons refer only to the independent experiment. The analytic homogeneous joint-inversion recovery gates **PASSED**. Median three-dimensional location error changed from 0.1821 to 0.0210 km, while median velocity error at the 203 illuminated nodes changed from 0.3000 to 0.0080 km/s. The absolute residual RMS changed from 0.35209 to 0.00635 s. The precise residual RMS changed from 0.09192 to 0.00031 s. There were 12 accepted velocity updates, 12 accepted location updates, and 0 sources on a boundary. LSQR met its normal-residual tolerance in 12 of 12 solves; the worst relative normal residual was 9.15e-09.

The fixed recovery criteria require positive finite velocities, no boundary sources, nonincreasing accepted frozen-weight objectives, at least tenfold reductions of both absolute and precise residual RMS, lower median location and illuminated-node velocity errors, and nonzero updates to both locations and velocity. Illumination means accumulated interpolated path length greater than 10^{-9} km on final rays; it does not establish resolution or recovery at every node. The arrivals are analytic and unrounded. This test examines a coupled homogeneous limiting case; it does not establish heterogeneous model recovery. Ray diagnostics compare production traces with half-spacing, 1000-iteration traces at the saved true, final standard, and final DD models and event locations. Source derivatives are compared with central differences of refined travel times using 0.01 km endpoint perturbations. A diagnostic exceedance uses the existing single-ray tolerance of 0.003 s/km; it is not a demonstrated inversion-error bound.

For the truth model, 32 rays had median and maximum absolute refinement changes of 0.054 and 0.892 ms. Production/refined convergence counts were 15/20 out of 32; maximum projected gradients were 0.00868/0.000668 s/km and maximum raw gradients were 0.00868/0.000668 s/km. Thus 17/12 production/refined traces failed the optimizer stopping criterion. Across 24 eligible source-coordinate derivatives, maximum production/refined errors were 0.000584/0.000599 s/km, with 0/0 components exceeding the diagnostic tolerance; 0 components were skipped at boundaries.

For the standard model, 32 rays had median and maximum absolute refinement changes of 0.155 and 0.684 ms. Production/refined convergence counts were 1/3 out of 32; maximum projected gradients were 0.0118/0.00645 s/km and maximum raw gradients were 0.0118/0.00645 s/km. Thus 31/29 production/refined traces failed the optimizer stopping criterion. Across 24 eligible source-coordinate derivatives, maximum production/refined errors were 0.000746/0.000328 s/km, with 0/0 components exceeding the diagnostic tolerance; 0 components were skipped at boundaries.

For the dd model, 32 rays had median and maximum absolute refinement changes of 0.100 and 0.861 ms. Production/refined convergence counts were 1/8 out of 32; maximum projected gradients were 0.0115/0.014 s/km and maximum raw gradients were 0.0115/0.014 s/km. Thus 31/24 production/refined traces failed the optimizer stopping criterion. Across 24 eligible source-coordinate derivatives,

maximum production/refined errors were 0.000589/0.000179 s/km, with 0/0 components exceeding the diagnostic tolerance; 0 components were skipped at boundaries.

Raw and projected gradients differ at active domain boundaries. Small changes in travel time do not establish derivative convergence, and unconverged refined traces are not exact references. These deterministic samples cannot establish global first-arrival optimality or the accuracy of every production ray.




Figure 3: Saved numerical ray through the specified sandwich medium, compared with the straight connection between identical endpoints. The shaded region marks the low-velocity trough. Bending reduces calculated travel time. This is a validation figure, not a manuscript-panel reproduction.

The exact homogeneous travel-time solution, derivative checks, reciprocal ray times, station-error cancellation, and known least-squares solution test different parts of the implementation (Appendix A). They do not establish uniqueness of the nonlinear inverse problem. A larger, deterministic ray sample tests the production settings against half-spacing, 1000-iteration traces. The source finite-difference checks are reported for both the true and inferred media. Small travel-time differences alone do not establish stationarity or accurate source derivatives. Raw gradients can also be nonzero at active constraints, so projected gradients and iteration limits are recorded separately. The homogeneous joint test generates arrivals analytically as distance divided by 6 km/s, independently of the ray tracer, and enables both source and velocity unknowns.

The cross-strike-only medium also admits an independent Snell-law reference for the validation ray, which crosses the layers monotonically. If R is the endpoint separation in the transverse plane, its conserved transverse slowness p satisfies

$$R = \int_{x_a}^{x_b} \frac{p\, v(x)}{\sqrt{1 - p^2 v(x)^2}}\, dx, \qquad T = \int_{x_a}^{x_b} \frac{1}{v(x)\sqrt{1 - p^2 v(x)^2}}\, dx. \tag{4}$$

This scalar reference tests both travel time and the source slowness vector without optimizing path vertices. At the 600-iteration setting, a source-perturbation sweep found finite-difference errors below 1.53×10^{-4} s/km at 0.1, 0.03, and 0.01 km, while the 0.001 km difference amplified ray-discretization and optimization errors to 0.00350 s/km. More iterations did not ensure finite-difference convergence. The consistency check therefore uses 0.01 km, matching the larger ray diagnostic sample; the derivative tolerance remains 0.003 s/km. The saved sweep in `data/development/` records this limitation. It is not evidence of derivative convergence in every inferred three-dimensional medium.

DD reduces the full-domain mean velocity error from the initial 0.456 to 0.305 km/s; standard tomography ends at 0.574 km/s. The DD value remains larger than the published reference of 0.178

Method	Median $\lvert\Delta x\rvert$	Median $\lvert\Delta y\rvert$	Median $\lvert\Delta z\rvert$	R_{abs}	R_{cat}	R_{prec}
standard	0.5232	0.4525	1.5176	0.3565	0.4621	0.4619
dd	0.3690	0.3266	0.5217	0.3526	0.4399	0.4390
loc1d	2.0050	0.5019	0.8753	0.6342	0.5095	0.5081
loc3d	0.0103	0.0221	0.1955	0.1771	0.0578	0.0154

Table 2: Independent synthetic results. Coordinate errors are in km and arrival residual RMS values are in s. Residual RMS values are unweighted, so they remain comparable when inversion weights change. Fixed-model relocations are denoted loc1d (homogeneous) and loc3d (true synthetic grid). These fixed-model comparisons retain absolute arrivals and do not copy all gauge conventions of hypoDD. The axes are Cartesian, not the latitude/longitude axes of published Table 1.

Method	Domain	Median	Mean	Standard deviation
Initial	Full domain	0.500	0.456	0.318
Initial	Interior	0.500	0.544	0.393
standard	Full domain	0.443	0.574	0.506
standard	Interior	0.258	0.313	0.245
standard	Published reference	0.164	0.245	0.249
dd	Full domain	0.252	0.305	0.262
dd	Interior	0.153	0.214	0.181
dd	Published reference	0.136	0.178	0.164

Table 3: Absolute velocity errors in km/s. Full-domain statistics use saved regular samples on all five depth slices. The interior restricts x to $[-15, 20]$, y to $[-40, 20]$, and depths to 3–11 km. Published values are reference scales only: the article does not specify the averaging mask, and its geometry and starting model differ.

Model	Depth (km)	Trough	West block	West minus trough
standard	3	4.440	5.561	1.121
standard	7	4.592	5.948	1.356
dd	3	4.202	5.779	1.577
dd	7	4.238	5.901	1.663
Specified true model	3, 7	4.000	6.000	2.000

Table 4: Spatial recovery of the specified trough and western block. All velocities and contrasts are in km/s. Arithmetic means use saved samples with $y \in [-20, 20]$ km, $x \in [1, 5]$ km for the trough, and $x \in [-15, 0]$ km for the western block, including window endpoints. These independently defined windows quantify recovery of the known 2 km/s contrast; they are not published recovered-model averages.

km/s, under the differing sampling and starting conditions described above. Median three-dimensional location error improves from 1.029 to 0.822 km with DD, although its median depth error, 0.522 km, is slightly worse than the initial 0.477 km. Thus improvement depends on both the quantity and the averaging domain.

The recovered DD western-block–trough contrasts are 1.577 and 1.663 km/s at 3 and 7 km depth, respectively, compared with the specified 2 km/s. DD preserves more contrast than standard tomography but still smooths the trough. Its unweighted precise-difference RMS remains 0.439 s, compared with 0.462 s for standard tomography and 0.0154 s with the true velocity fixed. The DD residual is far above the approximately 0.01 s upper bound on differential rounding error: the nearly exact synthetic differences are not fitted to their precision. Grid representation, regularization, source–velocity trade-offs, and robust downweighting are possible contributors; these experiments do not isolate their effects. Improved model-error averages therefore do not imply a close fit to every event pair. Among the three tested starting profiles, the homogeneous start gives the smallest final location and velocity errors; this is a sensitivity result for the independent experiment, not identification of the paper's start. The LOM and LON runs end with median three-dimensional errors of 1.432 and 1.147 km, respectively, both worse than the initial 1.029 km.

All sampled travel-time refinement changes are below 0.9 ms, smaller than the 10 ms rounding interval and 40 ms random-pick scale. None of the sampled source derivatives exceeds the 0.003 s/km diagnostic tolerance. However, most rays in the inferred models do not meet the stricter optimizer stopping criterion, including many refined traces. These checks show close agreement between the numerical settings on sampled paths. They do not certify absolute accuracy, all derivatives, or the first-arrival branch selected by every ray.

Each Figure 3 counterpart shows the five generated grid-depth slices at 0, 3, 7, 11, and 16 km. The published captions do not enumerate plotted depths, so this count is not a verified reconstruction of the published figure layout.

5 Discrepancies and reproduction limits

The largest structural difference from the target experiment is its sampling geometry. The synthetic coordinates in the original paper come from actual seismicity and station locations, not from a distribution completely specified in the text. Replacing that geometry changes the null space, ray coverage, and source–velocity trade-off. A homogeneous initial velocity also changes the nonlinear trajectory relative to the referenced layered model. We cannot attribute a mismatch with a published error statistic solely to the inversion algorithm under these changes.

Our forward grid represents the trough edges more finely than the inversion grid. This is a deliberate test of model representation error. The 0.3 s station biases are larger than the random pick errors, and the inverse problem has no explicit station-static parameters. It must accommodate these errors through source shifts, near-station velocity, or residual misfit. Robust weighting may suppress genuine model mismatch as if it were an outlier. A lower weighted objective therefore does not by itself establish a more accurate velocity model; velocity and location errors against the independent truth are reported separately.

During development, rejecting an entire trial update when one source left the domain caused the fixed-model inversion to stall. Projecting individual trial source locations onto the domain corrected that numerical obstruction. Independent review also found that boundary directions within the ray optimizer could block tangential motion, and that collapsed ray segments could corrupt endpoint derivatives. Both cases now have numerical regression checks. A further check found roundoff gradients in coordinate directions where velocity was exactly constant. Ray optimization and column scaling amplified these into spurious source-depth sensitivities. Evaluating each interpolation-gradient component with paired nodal differences preserves this symmetry exactly without changing the mathematical interpolant. Constant-

Figure 3a: Standard tomography
Partial reproduction: independently generated geometry and explicit numerical substitutions




Figure 4: Partial reconstruction of manuscript Figure 3a: standard tomographic velocity from independent observations. Dots are inferred sources selected by adjacent depth midpoints. Colors show 4–6 km/s; values outside this interval saturate and their numerical range is reported in the text. Unconstrained boundaries must not be read as resolved structure.




Figure 5: Partial reconstruction of manuscript Figure 3b: DD velocity from independent observations. The initial model, true medium, noise, and inversion grid are identical to those of the standard run. Colors and source-selection conventions follow the preceding figure.

Figure 3c: True minus DD velocity
Partial reproduction: independently generated geometry and explicit numerical substitutions




Figure 6: Partial reconstruction of manuscript Figure 3c: true minus DD velocity, following the subtraction order in the manuscript's prose. Blue marks recovered velocities that are too fast, and red marks recovered velocities that are too slow. The fixed color range is ±1 km/s. Saved samples retain unsaturated values.




Figure 7: Partial reconstruction of manuscript Figure 3d: true minus standard tomographic velocity, with the sign and color convention of the preceding figure. The spatial patterns depend on the independent geometry and cannot be compared point by point with the original inversion.




Figure 8: Independent convergence and location-error diagnostics. Top: RMS residuals before each nonlinear update. Stage transitions change the objective, so the unweighted RMS can rise while a weighted objective falls. Bottom: source-coordinate error distributions against saved true coordinates. These are additional method tests, not field-data panels.

axis and surface-ray regression checks cover the correction. The earlier numerical histories are retained in `data/development/`; they are not used for any final numerical panel. Domain projection does not make sources pinned to boundaries physically reliable. In the depth-independent true medium, a source at the surface has zero first-order depth sensitivity to the surface receivers. A linearized fixed-model relocation therefore cannot move such a source back into the interior. The boundary counts record this remaining geometric limitation; the symmetry correction removes numerical drift rather than supplying missing depth sensitivity.

For the initial Hayward Figures 4–11 attempts, the missing dependencies were observational. The field velocity model is the result to be inferred, not an input that can be reconstructed by assuming the reported contrast. We recovered the written grid, projection conventions, sample counts and section definitions, then traced each panel's required data. Those initial attempts stopped at excluded arrivals and event coordinates. In particular, resolution and uncertainty require the actual linearized ray matrix, damping convention and covariance model. A synthetic resolution matrix would not reproduce Figure 11, so none is substituted here.

6 Rebuild and artifact provenance

Run the following command from the project root:

```
./rebuild.sh
```

It configures and builds C++, runs validation, generates observations, performs four main inversion comparisons, two starting-profile sensitivities, fixed-model and joint recovery tests, and true/final-model ray checks, computes metrics, generates all figures and audit tables, and compiles this report twice with pdfLaTeX. Its outcome is recorded in `data/rebuild_result.txt` only when all commands have succeeded. Machine-readable numerical checks and iteration histories remain in `data/`; raw console logs are local diagnostics excluded from the shared repository. All caches and build intermediates are confined to the project directory.

For the baseline delivery, the complete clean procedure finished successfully on 9 October 2026 in 801 seconds, with all 32 implementation consistency checks passing and both PDF compilation passes completing. The synthetic baseline numerical results come from that run. `reference/delivery_environment.json` records the compiler, Python packages, TeX version, and parallel settings. Interpretation and plot-label revisions were compiled afterward from those same numerical outputs. Those synthetic inversions have not been regenerated with the subsequent ray-kernel corrections. The later Hayward runs use the revised kernel and have separate provenance; the 801-second record verifies only the synthetic baseline.

Requirements are a C++17 compiler, CMake, OpenMP, Python with NumPy and Matplotlib, and pdfLaTeX with standard packages. No tomography, optimization, or authors' software dependency is required. The saved input CSV files permit rerunning the inverse problem without regenerating random numbers. Standard-library random distributions may differ across platforms, so cross-platform bitwise equality of newly generated observations is not assumed. The experiment configuration, source code, plotting scripts, generated data and figure audit are retained with this LaTeX source.

7 Resumed work: authoritative inputs and ray corrections

On 9 October 2026 the user required real earthquake and station observations and metadata from authoritative sources, beginning with USGS and EarthScope. Saved records must retain SEED naming codes and resource identifiers. USGS supplied a Loma Prieta mainshock QuakeML product containing 315 P picks; EarthScope supplied 364 historical NC station records. Provider identifiers, station

epochs, retrieval URLs, raw-file hashes, and identifier attachments are saved in `data/authoritative/discovery/`. These are archive-access checks, not the manuscript's selected event and station set.

The user subsequently authorized a documented authoritative subset with its differences reported, resolving the exact-cohort selection question. Independent NCEDC queries select 1,768 Hayward earthquakes from 24,301 broad-query records. Their preferred origins have 42,236 NC P picks passing the initial quality and identity criteria, across 464 channels at 387 stations. The manuscript's 1,489 events and 52 stations are not forced by sampling. Current catalog revisions, including the geoid depth convention, remain explicit comparison differences.

At the user's request, metadata investigation preceded exclusion. Targeted EarthScope and NCEDC StationXML queries start NC.CMW1 in 2000, but the authoritative NCSN master history explicitly identifies its earlier channels and operation dates. All 142 previously unmatched picks now have unique historical matches; the full join contains 42,236 unique matches. Operation epochs do not establish calibrated response epochs. The examined written studies give no station list establishing inclusion or exclusion of any of the four investigated boreholes. Their manuscript membership remains unknown, and no study-based exclusion is applied. Identified records are preserved for a later additional-data comparison.

NC.CCH1, NC.CGP1 and NC.CSU1 retain provider-declared sensor elevations of 107, 219 and 375 m. StationXML defines channel elevation at the sensor; burial depth is not subtracted again merely because station and channel fields agree. Historical BSL wellhead tables conflict by one burial depth, and their ellipsoid labels differ from NCEDC's stated shared geoid convention. These inconsistencies are recorded without inferring a numerical datum conversion. NC.CMW1 uses its historical ELEV value of 214 m as receiver elevation under an explicit sensor interpretation. Agreement with later StationXML is consistency evidence, not an independent survey. Historical burial depth remains unknown and is not backfilled. The evidence is in `data/authoritative/stations/investigation/`.

The declared quality and model-domain filters retain 1,092 events and 37 physical stations, with 14,262 absolute P picks. All four investigated stations contribute observations. Figure 9 shows this independently selected geometry. Exact manuscript membership and waveform-correlation observations remain unmatched. The catalog DD calculation has produced numerical outputs, as has absolute-data tomography. Their velocity comparison is reported below. Fixed-model relocation and final ray diagnostics remain pending; the completed outputs do not establish nonlinear convergence.

Metadata review also found avoidable loss of differential links. Requiring the same metadata epoch split otherwise identical receiver locations: all 62 used NSLCs have one reported position across their used epochs. The initial strict selection and its identified inputs are preserved as a sensitivity subset. The primary pairing instead retains each pick's native epoch and permits links at the same NSLC and exact position. Such pairs are flagged because shared geometry does not establish unchanged timing calibration. An undirected neighbor cap of ten remains an independent pairing choice; its difference-count bound cannot recover the paper's reported count from the reported absolute-pick pool.

The code now preserves provider identities and channel epochs through numerical inputs and outputs, and computes rays only for observed source–receiver pairs. An absolute-data Jacobian exporter and a block-elimination resolution diagnostic have also been implemented. The resolution algebra agrees with a direct small joint-matrix calculation to floating-point precision. It has not yet been applied to an inferred field model. Its uncertainty calculation assumes that reported symmetric pick uncertainties represent independent one-standard- deviation errors; the archive does not supply confidence levels establishing this calibration. Correlated model errors and regularization bias are excluded.

Independent component work isolated two numerical deficiencies. First, two-point integration of a segment crossing velocity-grid boundaries left a 1.19×10^{-4} s/km discrepancy between its variational endpoint gradient and finite differences of computed time. Eight-point integration reduces that example's discrepancy to 5.6×10^{-12} s/km and its time error from 33.4 microseconds to numerical roundoff. A separate three-dimensional segment check gives 8.9×10^{-11} s/km. This improves integration accuracy;




Figure 9: Independent Figure 4 counterpart from identified authoritative inputs: 1,092 selected earthquakes, 37 receiver positions, and 77 horizontal model nodes. The source and receiver selection differs from the published 1,489-event, 52-station cohort. This is a partial geometry reproduction, not an inferred velocity or relocation result. All plotted identities and input hashes are saved.

it is not proof of exact gradients for every finite quadrature problem.

Second, the homogeneous transverse ray Hessian is a fixed-endpoint path Laplacian, whose condition number grows quadratically with path-node count. Replacing the scalar initial L-BFGS metric with an inverse weighted path Laplacian reduces this stiffness without changing the Fermat objective or admissible paths. Positive segment weights are midpoint slowness divided by segment length. An analytic circular-ray test in $v(z) = 5 + 0.2z$ km/s over 40 km failed to converge within 1,000 baseline updates at spacings 1, 0.5, and 0.25 km. The revised metric converges in 32, 23, and 25 updates with unchanged stopping criteria. Halving spacing reduces the travel-time error by approximately four. A metric-only control preserves this improvement, separating it from the quadrature correction.

All 48 current component checks pass. Forward retracing of frozen baseline models gives the following convergence counts out of 32 sampled rays:

Frozen model	Production: baseline / revised	Refined: baseline / revised
Standard	1 / 5	3 / 22
Double difference	1 / 9	8 / 23

Nonconvergence remains substantial, and some source finite-difference errors increase despite improved stationarity. These checks do not demonstrate that the 0.439-s baseline DD differential residual is corrected. The synthetic inverse solutions have not been regenerated with the revised kernel. Analytic comparisons, source hashes, all 48 checks, and frozen-model ray tables are in `data/development/ray_optimizer_audit/`. Run the standalone audit with

```
python3 diagnostics/run_ray_optimizer_audit.py
```

7.1 A stationary ray can arrive later

An isolated analytic test identifies a remaining branch-selection failure. The invented medium has $v = 4$ km/s for $-1 \le z \le 0$ km and $v = 4 + 0.2z$ km/s below zero. Both endpoints lie at $z = -0.5$ km. At horizontal separations of 20 and 40 km, the production tracer returns the straight path in the constant-velocity layer and declares convergence without an optimizer update. Its projected gradients are below 3×10^{-16} s/km. Explicit diving rays obtained from Snell's law arrive 0.0810 and 1.0106 s earlier, respectively. Fine polygonal evaluations of those diving paths agree with their analytic travel times within 0.9 microseconds. Thus the discrepancy concerns the selected ray branch, not the stopping tolerance or a small quadrature error.

Local stationarity therefore does not establish a first arrival, and a derivative consistent with the selected direct branch need not describe the first-arrival travel time. The field starting profile also has a constant above-geoid layer. The current Hayward inversions retain this tracer; the analytic counterexample does not quantify the error of any particular field ray or attribute the inferred source displacements to this mechanism. The test definition, numerical results, and source hashes are saved in `data/development/first_arrival_audit/`. No observational data enter this diagnostic, and it does not modify the field calculations.

Figure 10 tests an experimental first-order fast-marching method (FMM) initializer followed by the unchanged continuous ray bending; it preserves the historical first-order comparison. The original auxiliary-grid audit fails 10 of 61 acceptance checks across 72 cases. Independent refinement of the auxiliary grid and continuous path spacing to 0.0625 km reduces the travel-time errors at 20 and 40 km to less than 1.8 microseconds. However, 4 of 37 checks still fail in the 44-case refinement experiment. At separations 14.68 and 15 km, immediately beyond the 14.6604-km branch crossing, it retains the late direct ray. The time errors are 0.1765 and 3.1896 ms, and the corresponding source-gradient errors are 0.0673 and 0.0699 s/km. Every selected ray in these analytic experiments meets the local stopping rule. Refining local optimization therefore does not, by itself, resolve branch selection. The prototype

remains outside the production solver; all failures and the unchanged acceptance thresholds are retained in `data/development/fmm_initialization_refinement/`.




Figure 10: Original first-order FMM diagnostic, not a manuscript counterpart. The medium has $v = 4$ km/s for $-1 \le z \le 0$ km and $v = 4 + 0.2z$ below; both endpoints have $z = -0.5$ km. Left: the stationary straight ray and a faster FMM-seeded, continuously bent path at 40-km separation. Right: travel-time error relative to the analytic earliest branch, with the indicated auxiliary-grid/path spacings. The dotted line marks the branch crossing; the dashed line is the unchanged 0.025-ms tolerance. The symmetric logarithmic scale is linear within ± 0.001 ms. Both spacings miss the earlier branch near the crossing.

A controlled comparison separates the stencil order from source seeding. The optional second-order one-sided derivative is $(3T - 4T_1 + T_2)/(2h)$ when both upstream values are accepted and $T_2 \le T_1$; otherwise that direction retains first order. Here h is the auxiliary spacing, and T_1, T_2 are the nearest and next upstream times. The two opposite directions are compared on each axis. A separate seed rule uses the union of corners of every grid cell whose closure contains the receiver. It agrees with the original single-cell rule off grid, but restores seed symmetry when the receiver lies exactly on a grid plane, edge, or node. In 224 component comparisons the original rule retains 28 reflection failures; the closed-cell rule has none, with off-grid fields and paths unchanged.

Stencil	Receiver seeds	Failed at 0.25 km	Failed at 0.0625 km	Missed offsets (km)
First order	Original cell	10/61	4/37	14.68, 15
First order	Closed-cell union	10/61	4/37	14.68, 15
Second order	Original cell	9/61	2/37	14.68
Second order	Closed-cell union	9/61	0/37	None missed

Both auxiliary-grid and path spacings take the indicated finest value in each column; missed offsets refer to that finest refined calculation, among the tested offsets. The coarse suite contains 72 comparisons and 61 gates; the refined suite contains 44 comparisons and 37 gates, omitting the already tested homogeneous fixtures. The retained time and source-gradient tolerances remain 2.5×10^{-5} s and 3×10^{-6} s/km. With second order and the closed-cell rule, the finest refined time and source-gradient errors are at most 1.80×10^{-6} s and 4.43×10^{-7} s/km. All 44 refined paths extract successfully, meet the local stopping rule, retain exact endpoints, and stay in the domain. The coarse suites separately retain three first-order and four second-order strict endpoint discrepancies, at most 8.90×10^{-16} km; these are recorded outside the original accuracy-gate counts.

The repeated first-order control reproduces the historical numerical rows and all 61 gates exactly. Frozen source snapshots, output hashes, and the complete controlled comparison are in `data/development/fmm_order_comparison.json`. Passing this finite analytic suite is not a general first-arrival certificate or a demonstration of global second-order accuracy.

A subsequent three-dimensional check uses the positive affine velocity $v(\mathbf{r}) = 6 + \mathbf{g} \cdot \mathbf{r}$ km/s, with $\|\mathbf{g}\| = 0.12$ s^{-1} in a $[-15, 15]^3$ km box. One oblique off-grid geometry, its reflection, and its coordinate permutation are each traced in both directions. The exact circular rays remain at least 3.73 km inside the box. At 0.5-km auxiliary spacing and 0.25-km path spacing, all 72 checks pass across the six directional comparisons. The FMM-seeded, continuously bent paths have maximum time and source-gradient errors of 4.31×10^{-6} s and 4.58×10^{-7} s/km. Both this candidate and the selected minimum over seeds are tested separately, so a correct direct seed cannot conceal an inaccurate FMM candidate. The evidence is in `data/development/fmm_rotated_affine_baseline/`. These smooth affine cases do not establish branch completeness in arbitrary three-dimensional media. Field validation remains necessary, and the production inversions have not changed.

7.2 Independent reference for the initial one-dimensional model

An independent Snell-law calculation enumerates rays in a positive, nondecreasing, continuous piecewise-linear velocity profile. It uses the saved initial Hayward grid within its finite depth interval, $-1 \le z \le 20$ km; the profile is not extrapolated below the model. The candidates include monotone direct rays, all admissible downward turns, tangent paths along accessible constant-velocity intervals, and tangent paths along the finite bottom boundary. Rational coefficients from the binary64 inputs and outward-rounded interval bounds cover the turning parameters. Unresolved roots, tangencies, or boundary membership withhold the first-arrival result. Travel-time ordering still uses ordinary floating-point arithmetic; near ties are retained, and no interval-certified minimum-time gap is claimed.

The reference passes 250 original component checks and 106 added regressions. The added cases exposed and corrected cancellation in nearly flat-layer travel times and circular-path samples, and an incorrectly admitted root immediately beyond the finite bottom. The original 39 failed regression checks remain archived. Stable logarithmic and half-angle formulas remove the cancellation; ambiguous boundary membership now remains unresolved. These corrections change the reference times for the saved 32-ray Hayward sample by at most 8.9×10^{-15} s, with branches, source derivatives, and native identity mappings unchanged.

All 32 initial sampled rays have complete enumeration and distance closure within 7.3×10^{-12} km. Each has one candidate: 20 direct and 12 turning rays, with no near ties at the recorded 10^{-9}-s scale. The comparison preserves all 210 observation-to-ray identity associations.

Numerical time minus reference (μs)	Minimum	Median	Maximum
Production spacing	3.82	38.87	143.95
Refined spacing	1.32	11.79	37.48

For the 24 saved source-derivative components, the maximum absolute differences are 3.44×10^{-5} and 1.26×10^{-5} s/km at the production and refined spacings. These differences can contain path discretization and local optimization errors; they do not identify one mechanism. This initial sample does not show millisecond branch errors, but it does not validate every field ray or any updated three-dimensional model. The reference, checks, preserved failures, and identified comparisons are in `data/development/first_arrival_1d_enumeration/`.

7.3 Completed DD outputs and a targeted path comparison

The catalog DD run has completed its 18 configured updates; the other methods and full diagnostic workflow remain pending. Its outputs retain all 1,092 event identities and 58,980 observation mappings. Absolute/catalog-difference RMS residuals decrease from 0.3916/0.0934 s initially to 0.1606/0.0616 s finally. All 18 linear solves satisfy their stopping rule, but the final accepted maximum source-coordinate change is 2.7614 km; nonlinear convergence is not established. Thirteen sources reach $z = -1$ km and two reach $y = 30$ km. Nineteen have negative geoid depth, which does not establish position relative to local terrain. The DD output audit is in `data/development/hayward_final_output_review.json`.

The preserved first-order fixed-model diagnostic examines every saved DD absolute ray whose source and receiver depths both lie within $[-1, 0]$ km. This selection contains 186 rays and is not representative of all field paths. With auxiliary and continuous path spacings of 0.5 km, the prototype finds two admissible candidates more than 1 ms faster than the original paths: the reductions are 3.6916 and 2.2385 ms. A third is 0.0725 ms faster. A lower feasible action establishes a late original path in this fixed model; agreement does not certify a first arrival, and this experiment does not quantify source or velocity bias.

This original comparison retains its failures. One FMM extraction has no descending step (solver event index 1043, receiver index 284); its original direct-seed candidate is retained with an explicit failure flag. One selected ray misses the existing local stopping rule. All selected paths stay within the model domain, but 21 original direct paths fail the strict exact-endpoint check at arithmetic round-off, with maximum discrepancy 3.56×10^{-15} km. Without relaxing a threshold, the diagnostic saves its outputs and exits with failure status. Observation identities and native epochs are preserved in `data/development/fmm_hayward_audit/`. The unchanged production inversions remain separate; this comparison does not establish the cause of source displacement.

A controlled follow-up repeats the same 186 identified rays with the second-order stencil, closed-cell seeding option, and separately tested extraction correction, still at 0.5-km auxiliary and path spacings. All extractions succeed, but one selected ray still misses local stopping and 22 selected direct paths retain strict endpoint failures below 3.56×10^{-15} km. The failed checks remain recorded. Selected times change by at most 0.675 microseconds relative to the historical comparison; two admissible improvements over the straight seed remain above 1 ms, at 3.6916 and 2.2390 ms. All 30 physical receiver groups retain eight seed vertices, so the closed-cell option does not enlarge these seed sets. This comparison combines stencil and extraction changes; it does not isolate one correction. The complete evidence is in `data/development/fmm_hayward_order2_closed/`.

The remaining local-stopping failure has a measurable boundary bottleneck. An isolated audit retains the saved DD ray joining event index 775 to receiver epoch 184, channel `NC.CMO..EHZ`; its source lies on the artificial model top at $z = -1$ km. Instrumented and original functions give identical results in all 12 budget/spacing comparisons. Extending the update budget reaches the unchanged combined stopping rule after 1,249/1,155 updates at 0.5-km spacing and 3,009/2,703 updates at 0.25-km spacing (straight/FMM initialization). This rule requires an accepted action decrease below 10^{-11} s and projected gradient below 10^{-5} s/km; these paths do not meet the separate 10^{-7}-s/km gradient threshold.

More than 97% of accepted updates, including all of the final 200 updates in each extended run, follow the same fallback: projecting a full-space quasi-Newton direction at active faces makes it non-descending, so the code clears its curvature memory and uses the unpreconditioned projected negative gradient. No segments merge or line searches fail. Thus the boundary handling repeatedly discards the conditioning improvement measured in the interior tests. Extending the coarse budget changes travel times by only 2–4 ns, while source derivatives change by $2.1\text{–}3.4 \times 10^{-6}$ s/km. Path refinement changes derivatives by up to 1.1×10^{-5} s/km. Meeting local stopping alone therefore does not establish derivative accuracy or a first arrival. The original thresholds, failed runs, identities, and direction histories remain in `data/development/ray_stopping_i775_budget_path/`; the production inversions are unchanged.

An isolated direction control solves the principal weighted-Laplacian system on free coordinates, retaining the stiffness of edges adjoining fixed coordinates. Replacing only the original non-descent fallback still fails three of four runs: subsequent retained quasi-Newton directions exhaust the unchanged line-search limit. Using the reduced direction at every update reaches the original stopping rule in all four cases, after 12/12 updates at 0.5 km and 20/21 at 0.25 km (straight/FMM). All 65 directions have negative slope, and all 65 accepted updates strictly decrease the action with one line-search evaluation. Three runs meet the strict projected-gradient criterion; the coarse FMM run meets the combined criterion.

The reduced method does not release coordinates within a direction solve. Wrong-sign quadratic multipliers remain in 10 of 12 coarse and 20 of 21 fine FMM directions, so this method is not the exact tangent-cone quadratic minimizer. Independent matrix, feasibility, and affine-ray controls pass 560 fallback-variant and 564 pure-metric checks. Artifacts are saved under `data/development/`: `ray_stopping_i775_reduced_fallback/`, `ray_stopping_i775_reduced_all/`, and `reduced_ray_metric_checks/`. These results isolate a direction-choice effect for one identified ray. They establish neither general optimizer accuracy nor a globally first arrival, and they have not changed the production inversions.

7.4 Integrated ray correction and field preflights

The separate `first-arrival-integration` branch now combines the receiver-seeded first-arrival initializer with feasible constrained ray directions. When coordinate projection would change an L-BFGS direction, the method instead solves the positive-definite path-metric quadratic problem over directions allowed by the active model faces. It checks feasibility, multiplier signs, and descent before line search. This changes the numerical search direction; the travel-time action, domain, endpoints, quadrature, Armijo condition, and stopping rule remain fixed.

For the previously slow identified Hayward boundary ray, the integrated initializer and constrained direction meet the unchanged combined stopping rule in four updates at both tested discretizations. The 186 retained field entry checks have no initialization or local-stopping failure. Two selected actions decrease relative to the preceding implementation, by at most 5.63×10^{-9} s. Analytic layered and rotated-affine controls also pass at their assessed spacings. These checks support the implementation and isolate the boundary-direction defect; they do not certify all first arrivals or nonlinear field recovery. Full-cohort forward and inverse runs are separate requirements, and no completed corrected inverse result is available at this checkpoint.

The independently retrieved Loma Prieta geometry is also prepared for a replacement synthetic experiment: 724 catalog sources, 52 receiver positions, 37,648 source–receiver rays, and all 152,834 event pairs within the stated 20 km separation. The generator retains complete identified ray status and rejects failed rays before writing synthetic observations. The prepared full experiment has 15,932,384 observations, including both catalog and precise differences. LOM and LON starting profiles, source perturbations, and time-rounding differences are explicit assumptions in the preparation record. Geometry preparation and component checks are complete; the identified geometry diagnostic has finished with the failures described below. None of these prepared inputs has yet replaced the synthetic inversion figures earlier in this report.

7.5 Identified failures in the full-geometry checks

The larger geometry checks expose limitations that the preceding component suite did not resolve. These failures concern numerical paths in specified velocity models; they are not evidence for excluding any station or event. All identified inputs and failed results remain in the experiment.

The completed third Hayward preflight initializes all 14,262 absolute rays, but 139 fail the local stopping rule at 0.5 km field and path spacing. These failures span 128 events and 80 channel epochs.

Of the failed rays, 121 exhaust 1,000 updates; the remaining 18 stop earlier. All 32 rays in the separate refinement sample and all 24 endpoint perturbations meet their local stopping rules. Passing this sample therefore does not establish convergence on the full observation support. The terminal driver records a failed preflight and does not start an inversion. Its complete identified tables, configuration, source snapshot, and failure review are preserved in `data/development/hayward_first_arrival_003_review/`.

Continuous layered-ray enumeration has also completed for all 14,262 initial rays. Every reference is complete and valid, with no near ties; 9,069 first arrivals are direct and 5,193 turn below the source. Production times exceed the reference by a median 37.803 microseconds and a maximum 2.232 ms. Only 603 rays satisfy both existing analytic-control limits. The largest time and source-vector discrepancies belong to a locally converged ray: event `NC/30204761` to `NC.JJR..EHZ`, absolute row 14099. Its source-vector error norm is 0.033571 s/km. This discrepancy is independent of the 139 failed stopping flags. All reference candidates, original flags, identities, and frozen inputs are in `data/development/hayward_first_arrival_003_1d_support/`.

Event `quakeml:NC.anss.org/Event/NC/390228` and channel `NC.CCY..EHZ` define Hayward absolute observation row 9291. At 0.5 km field and initial path spacing, the selected FMM-started path rejects its line search after 208 accepted updates. Its time is 3.645674501 s and projected gradient is 2.7990×10^{-3} s/km. Increasing the iteration budget from 1,000 to 4,000 leaves this result unchanged. Observation hooks reproduce all eight uninstrumented candidate calculations exactly, including the paths, derivatives, iteration counts, and original stopping flags.

Two adjacent knots approach the 4 km depth interface within 2.18×10^{-13} km, leaving an approximately 1.01 km, nearly horizontal segment. Inserting a free midpoint preserves the exact curve and action. The new midpoint's depth derivative is -0.001645449 s/km; moving it 0.001 km deeper lowers time by 1.28394×10^{-6} s. The velocity increase below the interface reduces slowness, whereas a small depth tent has zero first-order length cost in the horizontal limit. Thus the retained coarse edge omits a feasible descent variation. The calculation uses the saved coordinates without moving them onto the interface.

Across all 38 segment insertions and 190 tested perturbations, the exact depth-only action changes by at most 3.04×10^{-18} s on insertion, and the production quadrature agrees with the exact action within 1.06×10^{-15} s. This excludes an action-integration discrepancy at the tested scale. At 0.25 km field and initial path spacing, the FMM-started path meets the unchanged combined stopping rule in 41 updates, with time 3.645534035 s and gradient 5.46×10^{-6} s/km. This resolves the local stopping failure for this refined calculation. Independent continuous layered-ray enumeration finds a unique first arrival at 3.645479994 s, turning at depth 4.000928869 km. The coarse selected path is therefore late by 194.506 microseconds; the refined FMM path remains late by 54.041 microseconds, with maximum source-derivative error 1.51×10^{-5} s/km. Its local flag does not establish continuous-model accuracy. The records are under `data/development/initial_ray_first_arrival_003_9291_audit001/` and `initial_ray_first_arrival_003_9291_midpoint001/` in the same parent; the independent reference is in `initial_ray_first_arrival_003_9291_reference001/`.

A further test retains the 0.5 km initialization field and refines only the path. At 0.125 km spacing, the selected FMM-started path stops after 20 updates, with time error 13.075 microseconds and source-vector error norm 2.66×10^{-6} s/km. At 0.0625 km, these errors fall to 3.267 microseconds and 7.88×10^{-7} s/km. Both selected results meet the existing analytic-control limits of 25 microseconds and 3×10^{-6} s/km. The 0.125 km straight-started candidate exceeds the source-derivative limit despite local convergence. All four candidates are retained in `initial_ray_first_arrival_003_9291_fine_path001/` under the same development-data directory. This single-ray result motivates further checks; it does not validate a full-cohort discretization.

Path refinement resolves both stopping and accuracy failures on the 139-ray subset (Table 5). Continuous layered-ray enumeration finds a unique first arrival for every ray. At 0.125 km spacing, all time

errors satisfy the 25 microsecond control limit, but 16 source-vector errors exceed 3×10^{-6} s/km. Three source errors remain above that limit at 0.0625 km. All 139 rays meet both limits at 0.03125 km, with maximum time error 0.818 microseconds and source-vector error 1.83×10^{-6} s/km. These calculations retain the frozen executable and original 1,000-update budget. They validate this subset at the initial model, not the full observation support or a later three-dimensional model. The four identified variants and their references are saved under `data/development/hayward_initial_failure_refinement_001/` through `hayward_initial_failure_refinement_004_reference/`.

Path (km)	Field (km)	Local passes	Both limits	Max time (μs)	Max source (s/km)
0.25	0.25	134	114	54.041	9.80×10^{-5}
0.125	0.5	139	123	13.075	2.87×10^{-5}
0.0625	0.5	139	136	3.267	4.92×10^{-6}
0.03125	0.5	139	139	0.818	1.83×10^{-6}

Table 5: Refinement of all 139 rays that failed the third Hayward preflight. Counts are out of 139. Accuracy uses absolute travel-time error and the Euclidean norm of the three-component source-derivative error relative to the continuous initial-model reference. The limits remain 25 microseconds and 3×10^{-6} s/km. The first row refines both discretizations; the remaining rows retain the same 0.5 km initialization field.

The locally converged outlier follows a later stationary branch (Figure 11). For this identified geometry, the continuous initial profile admits three turning rays. The first arrival turns at depth 10.896102 km with time 10.775535399 s. A deeper branch turns at 14.139000 km and arrives 2.185310 ms later. Refining the path to 0.03125 km while retaining the 0.5 km initialization field brings the numerical action within 0.353 microseconds of the deeper branch, but does not select the first arrival. Both straight and field starts remain on this branch, and increasing the update budget to 4,000 does not change their already-converged results. Exact integration of the saved polygons agrees with production action within 1.44×10^{-14} s; finite differences of those fixed polygons agree with the returned source partials within 3.06×10^{-9} s/km. These controls isolate branch selection from action integration and fixed-path differentiation for this case.

At the same 0.03125 km path spacing, refining the field to 0.25 km still selects the deeper branch. A 0.125 km field seeds the earlier branch: the bent ray stops after six updates with time error 0.333 microseconds and source-vector error 5.98×10^{-9} s/km. Its strict projected gradient is 9.49×10^{-8} s/km. The straight-started control remains on the deeper branch. These results motivate a full-cohort check at the finer settings; they do not establish accuracy for other initial rays or updated three-dimensional models. Full resource identities, epochs, paths, source snapshots, and independent comparisons are retained under `data/development/initial_ray_first_arrival_003_14099_audit001/` and the sibling field variants with suffixes `field025001` and `field0125001`.

Tightening only the stopping rule does not resolve every source sensitivity. For the near-interface ray from event `NC/321644` to `NC.CBR..EHZ`, row 8147, a separate control removes the combined small-action-change stopping condition while retaining the strict 10^{-7} s/km gradient criterion. The eight stricter calculations use both starts, path spacings 0.125 and 0.0625 km, and budgets of 1,000 and 4,000 updates. Six exhaust their budget; two reject the line search after 597 updates. None satisfies strict stationarity or the source-vector accuracy limit, although every time error remains within 25 microseconds. Further optimization therefore does not establish sensitivity accuracy at these path spacings. The unresolved stationarity also prevents attributing these errors solely to the finite polygon representation. The full comparison is in `data/development/initial_ray_first_arrival_003_8147_strict_comparison001/`.

The Loma Prieta check also separates local convergence from arrival-branch selection. The terminal record retains all 104 requested samples. For 38 rays, an independently integrated path that grazes the maximum-speed plane $x = 0$ is admissible. Twenty-three production and refined actions exceed




Figure 11: Arrival-branch failure for event `NC/30204761` and channel `NC.JJR..EHZ`, initial absolute observation row 14099. (a) Continuous layered reference curves and saved numerical paths projected onto the endpoint vertical plane; crosses mark the fixed endpoints. Open symbols use 0.03125 km path spacing and the indicated fast-marching field spacing. The horizontal line marks the 12 km velocity-profile interface. (b) Path refinement at fixed 0.5 km field spacing approaches the deeper stationary branch; the dashed line gives its 2.185310 ms arrival delay. (c) Field refinement at fixed 0.03125 km path spacing recovers the earlier arrival. Values in (b,c) select the smaller action from straight and field starts with the unchanged 1,000-update budget. Every selected candidate reports local convergence. This is an independently generated numerical diagnostic, not a reproduced manuscript panel.

that candidate. Three production paths report local convergence while remaining 25.693, 20.765, and 37.159 ms slower: event `NC/20091683` to `NC.HLT..EHZ` and `NC.HQR..EHZ`, and event `NC/256583` to `NC.JSC..EHZ`, respectively. At refined spacing their excesses fall to 0.166, 0.187, and 0.133 ms, but none meets the local stopping rule. The explicit candidate is an upper bound on the first arrival, not a complete enumeration of branches. Its absence for another ray does not justify excluding that ray. Full native identities, epochs, selector flags, and captured rows are retained in `data/development/loma_geometry_preflight/all_guides_002/`.

A subsequent continuous reference enumerates direct, turning, and maximum-speed guide branches for all 104 sampled Loma geometries. Every reference passes its completeness and distance-closure checks, with no near ties: 73 first arrivals are direct, 23 follow a critical guide, six turn, and two are direct paths in a constant-speed region. The 23 critical guides confirm the earlier feasible candidates as first arrivals for those rays. Production and refined root-mean-square time errors are 4.879 ms and 67.366 microseconds; their maxima are 37.159 ms and 186.569 microseconds. Even among locally converged production rays, the maximum error is 37.159 ms. The reference applies to the specified laterally varying synthetic truth model within its finite box and does not validate the production path solver. All identities and comparisons are saved in `data/development/loma_geometry_preflight/layered_reference_001/`.

The completed Loma Prieta diagnostic retains all 104 requested rays. Initializers succeed for every ray at both spacings, but 31 rays fail the local stopping rule at each spacing. The largest time change on refinement is 37.026 ms. Endpoint finite-difference comparisons reach approximately 1.26×10^{-3} s/km, but only 13 of 24 positive perturbations and 12 of 24 negative perturbations converge locally. These discrepancies cannot be interpreted as validated derivative-error estimates. No full synthetic observation set or Loma Prieta inversion follows this check. The identified terminal tables are in `data/loma_lom_001/`; the review and execution provenance are in `data/development/loma_geometry_preflight/001/`.

A separate exact control identifies a defect in using an ordinary gradient at an interface. For

$v(x) = 6$ km/s on $x \le 0$ and $v(x) = 6 - 2x$ km/s on $0 \le x \le 1$ km, a straight 2 km path on $x = 0$ attains the lower bound $T = 1/3$ s. The current trace returns this action but reports nonconvergence, using a midpoint normal gradient of 1/18 s/km from the right-hand cell. The true action has unequal one-sided derivatives there. The returned source-normal partial is 1/36 s/km, whereas the optimized continuous arrival has zero source-normal derivative at this control. Relabeling its stopping flag would therefore leave the sensitivity problem unresolved. The reproducible control is in `data/development/interface_stationarity/`. This component defect is distinct from the off-interface, slowly optimized Loma paths already examined. No convergence-only correction has been applied, and these checks have not produced a completed corrected inversion.

7.6 Full initial Hayward accuracy after separate refinements

The fourth Hayward calculation applies the 0.03125 km path spacing and 0.125 km initialization-field spacing to all 14,262 initial observed rays. Every initializer succeeds and every local stopping flag is true. All rays also satisfy the unchanged continuous-reference limits of 25 microseconds in time and 3×10^{-6} s/km in source-vector norm. The largest errors are 0.959 microseconds and 1.782×10^{-6} s/km; their medians are 0.333 microseconds and 2.422×10^{-8} s/km. In the preceding calculation, only 603 rays satisfied both accuracy limits. Figure 12 compares the complete distributions without removing failed rays.

The reference cache is reused only after exact equality of the complete initial geometry, model, observation tables, and numerical reference source has been verified. The original native identities, branch certificates, and local flags are retained in `data/development/hayward_first_arrival_004_1d_support/`. This passes the full initial-model comparison. The separate sampled refinement calculation remains active at this checkpoint; no corrected inversion is accepted or launched. The initial depth-dependent reference does not establish accuracy for later three-dimensional velocity models.




Figure 12: Initial Hayward errors relative to independently enumerated continuous first arrivals. Both curves retain the same 14,262 native observations and references. The left panel shows absolute travel-time error; the right shows the Euclidean norm of all three source-derivative errors. Dashed lines mark the unchanged accuracy limits. Refining path spacing and the initialization field together brings every fourth-run ray within both limits. This diagnostic does not validate a recovered velocity model.

7.7 Exact branch seeds and continuous synthetic truth

Supplying the exact earliest branch separates initialization error from path representation and local optimization. Four identified Loma rays, diagnostic rows 5, 49, 57, and 88, have continuous first arrivals along the critical guide at $x = 0$. Circular arcs and the straight guide are sampled at five spacings from 0.5 to 0.03125 km. One calculation uses the unchanged tracer, including its initial uniform resampling. A second copied tracer retains the supplied nodes initially; all interior nodes remain free during optimization. The action, derivatives, line search, and 1,000-update budget remain unchanged.

Only one of twenty optimized uniform-seed paths meets the local stopping rule; none of twenty retained-seed paths does. At 0.03125 km, all four uniform-seed results nevertheless have time errors below 4.750 microseconds and source-vector errors below 4.35×10^{-7} s/km. The finest retained row-5 path accepts no update, despite time error 4.696 microseconds and source-vector error about 2×10^{-15} s/km. Its ordinary gradient norm is 0.0751 s/km, and collective normal displacements of its interface nodes have unequal one-sided action slopes. The correct arrival branch therefore does not resolve the nonsmooth optimization problem. These controls also show that initial uniform resampling can damage a coarse guide seed; its additional row-5 time error falls from 473.730 microseconds at 0.5 km to 0.05368 microseconds at 0.03125 km. Exact polygon integration agrees with production action within 2.25×10^{-13} s. All paths, original flags, identities, and isolated source changes are retained in `data/development/loma_geometry_preflight/reference_seeds_001/`.

A separate directional calculation resolves two contributions to the large row-5 gradient. For an interface-aligned segment, the perturbed normal coordinate interpolates both endpoint directions. Its first variation must integrate the appropriate one-sided slowness derivative on each side of any directional zero; choosing one incident cell for the entire segment is insufficient. Homogeneous, fast-plateau, and reversed-jump cusp controls check this construction. The saved finest retained polygon also admits a small tangential descent. Independent 80- and 100-digit exact-segment evaluations agree on action reductions of 4.532×10^{-12} s and 2.028×10^{-10} s for two saved feasible directions. Thus the interface convention dominates the large ordinary gradient, but the polygon retains actual optimization error. Smaller normal-direction signs near 10^{-17} s are inconclusive. Neither finitely many probes nor accurate travel time justifies changing the original failed stopping flag. The identified paths, directions, and reviewed classification are saved in `data/development/interface_directional/001/`.

An interval supporting-vector calculation bounds the continuous first variation for every displacement of the retained interior knots: $F(d) \geq -B\|d\|$, with $B \leq 6.825849 \times 10^{-6}$ s/km; the arithmetic interval-radius norm is below 2.444×10^{-37} s/km. Reciprocal-affine segment primitives and rational arithmetic bounds replace fixed quadrature. A feasible convex-guide support removes the incident-cell normal artifact, while the saved tangent still has strictly negative first variation. Proofs, controls, and independent reviews are retained in `data/development/interface_stationarity_bound/` and `data/development/native_interface_support/`.

Two isolated steps test the complete first variation, including its convex-cusp term, before interval Armijo acceptance. The negative supporting-vector direction and its tangent projection first pass after 19 halvings. Action decreases by about 4.673×10^{-12} s, but new support-norm bounds increase to about 1.842×10^{-5} s/km. All 40 trials, identified rational polygons, and original false flag remain in `data/development/native_interface_step/001/`. Exact endpoint-moment analysis shows that the selected support already minimizes the Euclidean norm for the original and full-step polygons; changing normal support cannot remove their smooth residual. The tangent-step sign criterion is inconclusive. The proof and 156 controls are in `data/development/interface_moment_set/`. No production stopping rule, optimizer, or first-arrival certificate follows from these isolated results.

Continuous layered enumeration was then extended to all 37,648 pairs in the prepared 724-source, 52-receiver geometry. The first attempt completed every candidate enumeration, but eleven direct candidates failed the unchanged 10^{-8} km distance-closure requirement. Their first-arrival values remain blank. All eleven use source 357, event `NC/20091139`, at $x = -0.011677621261526383$ km, just inside

the 6 km/s constant-speed region. Solving for $pv_{\max}$ close to one loses the small critical offset in binary64 arithmetic. A separate correction solves for $\delta = 1/p - v_{\max} \geq 0$ and forms the phase speed by exact rational addition before evaluating the segment integrals. Twenty-one isolated cases, including all eleven failed geometries and ten invented near-interface controls, close within 7.11×10^{-15} km. Independent 80-digit integrals agree in time within 3.91×10^{-15} s and in source-vector norm within 2.87×10^{-17} s/km. The failed first attempt is preserved under `data/development/loma_layered_full_001/`.

The corrected second enumeration passes the full geometry and independent native-identity review: all 37,648 references are valid, with no near ties. Their first arrivals comprise 25,724 direct paths, 5,951 turning paths, 5,474 critical guides, and 499 direct paths in constant-speed regions. The maximum closure residual across all 66,302 candidates is 3.13×10^{-9} km. The eleven formerly invalid references are the only validity changes. Previously valid branch kinds remain unchanged; their largest time change is 1.55 ns. No source or receiver is removed. Both compressed identified streams, the executed reference source, and the separate completion, physical-consistency, and independent identity reviews are retained in `data/development/loma_layered_full_002/`.

An explicit analytic-truth input mode permits a separately reviewed, complete continuous reference to supply synthetic travel times. The C++ consumer verifies the model, ordered source–receiver support, every native metadata field, finite values, unique valid references, and physical speed bounds. It preserves the existing random-draw, quantization, and differential observation order. Its invented contract suite passes 55 checks, and six default-mode output files remain byte-identical to the frozen preceding executable. Analytic mode writes reference status separately and leaves polygon-ray status fields blank. This changes the declared synthetic truth calculation; it does not repair or accept the inverse forward operator.

The actual LOM-profile simulation now writes all 37,648 absolute observations, 7,947,368 catalog differences, and 7,947,368 precise differences, totaling 15,932,384 rows. A separate streaming audit checks every arrival and observation, including order, both native event identities, receiver epoch, quantization, station bias, and saved-noise arithmetic. All checks pass. The largest saved catalog-subtraction discrepancy is below 1.01×10^{-10} s, within the explicit bounds from 12-significant-digit serialization. This rounding bound does not relax a scientific accuracy criterion. The raw run remains preserved, and `data/published/loma_lom_analytic_001/` contains a lossless ten-part archive of 504,089,284 compressed bytes. Every member was read back and verified.

The independently audited LON-profile simulation contains the same number of rows. Full-file comparison establishes byte-identical synthetic observations, truth arrivals, and perturbed event geometry. Its starting grid differs at all 288 velocity nodes; the numerical configuration changes only the declared starting-profile selector. The distinct source snapshot, configurations, execution records, and reviews are retained. Identical simulated observations do not establish equivalence of inversions from the two starting models.

The inverse calculation starts from perturbed sources and a depth-dependent grid, requiring references distinct from the x-dependent synthetic truth. All 37,648 LOM initial references pass their independent reviews, with unique first arrivals and defined source derivatives; 59 pairs use the finite-bottom branch. A separate LON enumeration preserves one invalid reference among the same 37,648 pairs. Pair 8156, event `NC/255127` at `NC.JSC..EHZ`, has a turning candidate just below the 11 km knot and distance residual 1.0784×10^{-8} km, above the unchanged 10^{-8} km requirement. Its selected time and derivatives remain blank. No tolerance is relaxed. Both full identified reference streams and their reviews are retained in `data/development/loma_initial_reference_001/` and `data/development/loma_lon_initial_reference_001/`.

The LON failure arose when reconstructing turning speed from the isolated root. Rounding $u = \sqrt{u_0^2 + r^2}$ to binary64 loses part of the small offset above the layer-knot speed u_0. A separately reviewed variant retains a rational square-root enclosure midpoint while evaluating each turning candidate. It leaves the isolated roots, physical formulas, and acceptance limits unchanged. All 37,648 references

now pass in the separate second attempt. Every one of the 37,752 candidate signatures is retained, with no change to a previously selected branch. Pair 8156 is the only validity change; its original failed values remain blank in the first attempt. The maximum candidate-time change is 1.701 nanoseconds, and the maximum source-vector change is 1.321×10^{-13} s/km. Maximum distance closure residual decreases to 9.652×10^{-10} km. The full identified comparison and independent readback are retained in `data/development/loma_lon_initial_reference_002/`. This corrects the reference arithmetic; it does not validate a production ray or inversion.

Separate frozen LOM and LON initial-forward preflights are now active. The LON preflight uses the corrected complete reference and the same numerical settings as LOM. Seven bounded controls verify unchanged matrix-only outputs between the earlier LOM executable and the guarded executable used for LON, including retention of false local-convergence flags. These controls and the launch eligibility review are saved in `data/development/loma_lon_initial_preflight_adoption_001/`. Each preflight exports full source derivatives and exact endpoint/domain checks from the rays used to count matrix entries. A separate sampled refinement retains all 104 declared samples and 24 valid source finite differences while limiting cached fine receiver fields. It performs no inverse solve. Neither simulation completion nor a matrix storage estimate establishes inverse convergence.

7.8 Rejecting unsupported forward evaluations during inversion

The preceding inversion code recorded local ray failures but could still use those rays in a Jacobian, a line-search objective, or final output. An initial preflight therefore did not protect later model updates. A new guard on the integration branch applies the unchanged local, finite-value, domain, endpoint, initialization, and derivative-dimension checks before each such use. A failed evaluation stops the inverse run and saves its model, identified ray statuses and available paths separately from the last accepted state. Failure snapshots retain 17 significant digits. Unevaluated derivatives remain blank, including caught worker exceptions.

Invented component runs exercise failures at the current model, a trial step, and the final model. They reproduce the parent code's use of unconverged rays; all seven outputs of a successful fixed-model location inversion remain byte-identical. The matrix-only diagnostic retains its original behavior. Independent review and the optimized application build pass. These checks, saved in `data/development/inversion_ray_gate_checks/`, establish failure handling rather than global first-arrival accuracy or nonlinear recovery. No live executable was replaced, and no corrected inverse result is claimed.

7.9 Historical waveform metadata and identified correlations

The existing Hayward catalog pair at differential row 15108 links events `NC/238035` and `NC/30083101` at `NC.JPR.02.EHZ`. EarthScope routing identifies NCEDC as the archive for the two requested windows. The preserved miniSEED records each contain 500 contiguous samples at 100 Hz over a half-open five-second interval. Full response metadata cover the distinct 1991 and 1995 epochs; their three-stage response contents agree. Event, origin, pick, association, channel, and epoch identities accompany the raw files and derived records. Missing provider response resource IDs remain explicitly missing.

The written precursor method uses a tapered 2.56 s window and requires squared coherence above 0.9 at at least half the frequency bins between 2 and 10 Hz (Waldhauser and Ellsworth, 2002, section 2.2, paragraph 16). For the primary pilot calculation, the unspecified choices are declared: a window beginning 0.5 s before the catalog P pick, demeaning, a Hann taper, and five-bin spectral averaging. No additional filter, response removal, or resampling is applied. Only two of twenty frequency bins pass, where ten are required. All nine declared window/smoothing variants fail, with zero to four passing bins. Consequently, no waveform difference is adopted. A line through the two coherent phase bins has two free parameters; its near-zero residual does not measure timing precision. The source, spectra, controls, and failed result are in `data/development/waveform_correlation_pilot_001/`.

An offline audit retains all 14,262 picks and 44,718 catalog differences. Historical operation records identify `NC.CMW1`, but do not provide calibration: the saved EarthScope and NCEDC StationXML responses start in 2000, whereas 117 retained picks and 199 differences involve its 1995–1998 epochs. No later response is backdated. Among 398 picks whose channel names do not end in Z, provider orientation metadata identify 227 as vertical, 54 as nonvertical, and 117 as unknown. Channel names alone therefore cannot define the vertical waveform selection. A further 89 picks and 542 differences precede the archive's documented March 1984 start; this date is a coverage lead, not proof that all earlier data are absent. These audit categories remove no catalog observations and establish no study exclusion. Full native rows and the original unsubmitted requests are saved in `data/authoritative/waveforms/cohort_plan_001/`.

Normalized waveform correlation supplies differential times without calibration corrections, as clarified by the user. Missing historical responses remain metadata gaps rather than retrieval prerequisites. EQcorrscan documents local-window normalization, pick-relative extraction, and `dt.cc` identifier and sign conventions.[1] Differential time is travel time for event 1 minus that for event 2. The failed spectral pilot remains preserved as a separate calculation.

NCEDC reports covering spans for 464 pick windows; 13,798 have no reported span, which does not establish historical absence. Five sample batches provide 459 complete windows of 500 native samples at 100 Hz and five identified zero-sample outcomes, all at `NC.JPR.02.EHZ`. The unchanged first response is reused; no omitted window is re-requested, and partial or overlapping support is not accepted. The complete identified inventory, raw replies, and sample audit are in `data/authoritative/waveforms/cohort_cc_002/` and `data/authoritative/waveforms/cohort_samples_002/`. The first decoding failure remains in `data/authoritative/waveforms/cohort_samples_001/`. No catalog observation or station is removed.

The declared method correlates 256 native samples over the complete ±0.25 s search. The template starts nearest to 0.5 s before its pick, with half-sample ties resolved earlier. Direct local centering and scaling evaluates signed Pearson correlation; 41 controls include large-amplitude cancellation cases. No response removal, extra filter, resampling, or fractional interpolation is used. Nonpositive, tied, and boundary maxima are rejected. After discrete alignment, the declared Hann-window and five-bin spectral screen requires at least ten of twenty 2–10 Hz bins to have squared coherence strictly above 0.9. Actual native sample offsets determine the sign:

$$\Delta t_{ij}^{\mathrm{cc}} = \Delta t_{ij}^{\mathrm{catalog}} - \mathrm{delay}_{j-i}.$$

Native-input controls check both epochs, origins, sample hashes, and timing.

Of the original 44,718 pairs, 1,195 have eligible samples on both sides. The calculation accepts 98; 1,086 fail coherence, ten peak at the boundary, and one has a nonpositive peak. The other 43,523 remain unattempted. The 98 accepted event pairs each have one accepted channel and span 103 events: 75 are at `NC.JPR.02.EHZ`, 19 at `NC.CBR.10.EHZ`, three at `NC.JSB.10.EHZ`, and one at `NC.JSF.10.EHZ`. The complete identified roster and all attempted lag/coherence arrays remain in `data/development/native_waveform_cc_001/`. These new subset observations are not identified copies of the manuscript's 17,955 waveform differences and have not entered an inversion. Neither squared correlation nor the 10 ms sample spacing is a formal timing uncertainty. The checked `dt.cc` export in `data/development/native_dt_cc_001/` retains a complete 44,718-row identity sidecar, event integer map, and receiver epochs. It preserves binary64 values, rejects ambiguous mappings, and leaves solver weighting and the running catalog-only calculation unchanged.

[1] EQcorrscan documentation: normalized correlation; correlation-derived differential times.

8 Hayward velocity comparison before final diagnostics

Figure 13 displays the saved standard and catalog-DD velocity models for the authoritative subset of 1,092 events and 37 physical stations. Both use 14,262 absolute picks; the DD inversion also uses 44,718 catalog differences. The calculation contains no waveform-derived differences. These 21 section panels are partial counterparts to Figure 10, with the cohort, pairing, initial model, and numerical differences retained.

The standard inversion accepted 12 updates and then rejected all line-search steps at iteration 12. Its absolute and differential residual RMS values are 0.07869 and 0.07162 s. At that iteration, 572 of 14,262 ray calculations failed the local stopping rule. The DD inversion reached its 18 configured updates; its final accepted maximum source-coordinate change was 2.7614 km. Thus neither termination establishes nonlinear convergence. The histories describe pre-update rays; independent final-model ray diagnostics remain pending. The plotted results retain the original straight-seed ray algorithm. The validated initializer and constrained-direction corrections have not yet produced a completed replacement inversion.

The color limits remain fixed across sections and methods. Standard velocity falls below the 3.5 km/s display limit at three nodes. Across the 83,895 sampled section points, 1,136 DD-minus-standard values are below -0.8 km/s and 1,265 exceed 0.8 km/s; the full difference range is $[-1.16214, 1.37544]$ km/s. Triangular colorbar extensions expose saturation. These differences compare two regularized estimates, not error relative to independent velocity truth.

Regenerate these panels from the saved models with

```
python3 plot_hayward.py --velocity-only
python3 audit.py
python3 analyze.py
```

The source and output hashes, event identities, stopping histories, and section ranges are linked by `data/hayward/figure10_velocity_manifest.json`. The separate full-figure completion record is not written by this command.

Figure 10 counterpart: independent subset; points within 5 km of each section




Figure 13: Partial Figure 10 counterpart from the saved original field runs: catalog-DD P velocity, absolute-data P velocity, and DD minus absolute-data velocity at seven written section locations. Hypocenters are within 5 km of each section. The full computational depth interval includes the layer above the geoid needed to contain elevated receivers. Colorbar extensions mark values outside the common display limits. These numerical outputs retain unresolved ray and inverse-problem convergence limitations.

A Numerical validation record

This appendix retains the baseline 32-check record supporting the baseline figures.

Check	Error	Tolerance
homogeneous_travel_time	4.44e-15	1e-10
homogeneous_source_derivative	5.19e-17	1e-09
homogeneous_velocity_derivative	4.44e-16	1e-09
partition_of_unity	1.42e-14	1e-09
ray_reciprocity	6.58e-07	2e-05
bending_not_slower_than_straight	0	1e-10
ray_refinement_seconds	4.14e-05	0.015
bent_source_finite_difference	9.22e-05	0.003
velocity_finite_difference	3.71e-13	1e-07
identical_event_dd_cancellation	0	1e-12
station_bias_cancellation	0	1e-12
lsqr_known_solution	7.11e-16	1e-09
matrix_adjoint	0	1e-12
interpolation_constant_x_symmetry	0	0
interpolation_constant_y_symmetry	0	0
interpolation_constant_z_symmetry	0	0
surface_ray_plane_symmetry	0	0
surface_ray_source_symmetry	0	0
snell_reference_travel_time	6.9e-05	0.001
snell_reference_source_derivative	8e-06	0.0001
trilinear_gradient_finite_difference	3.52e-12	2e-10
boundary_ray_feasible_descent	0	1e-10
boundary_ray_projected_gradient	1.84e-06	1e-05
boundary_ray_domain	0	0
collapsed_vertices_preserve_endpoints	0	0
collapsed_ray_minimum_segment	0	0
collapsed_ray_gradient_regression	4.53e-05	0.001
scaled_damped_lsqr_solution	2.48e-16	1e-10
scaled_damped_lsqr_normal_residual	5.55e-17	1e-08
scaled_damped_lsqr_converged	0	0
zero_rhs_lsqr	0	0
truncated_lsqr_not_converged	0	0

Error units follow each test: seconds for time comparisons, seconds per km for source derivatives, seconds per (km/s) for velocity derivatives, and algebraic units for the sparse-system checks. These tolerances were chosen for numerical consistency tests, not to enforce agreement with the paper.

B Figure-by-figure record

The complete machine-readable record is in `data/panel_audit.csv` and `data/panel_audit.json`; its readable counterpart is `FIGURE_AUDIT.md`. "Independently reproduced" means the quantity is reconstructed from the written specification using our model output. "Partially reproduced" means an independent numerical counterpart exists but one or more specified experimental inputs or algorithms differ. "Not reproduced" means no corresponding result is claimed.

Panel	Quantity	Status and limiting dependency
1	Horizontal true P-velocity slice	independently reproduced. Physical velocity profile reconstructed from prose; plotting extent and depth chosen from model domain.
2	Synthetic-test event/station geometry and inversion grid	partially reproduced. The saved baseline uses random geometry. A documented authoritative Loma Prieta subset is authorized; it has not yet replaced this experiment. No digitization used.
3a	Standard tomographic P velocity; all generated depth slices	partially reproduced. Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
3b	DD tomographic P velocity; all generated depth slices	partially reproduced. Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
3c	True minus DD P velocity; all generated depth slices	partially reproduced. Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
3d	True minus standard P velocity; all generated depth slices	partially reproduced. Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
4	Hayward network, catalog seismicity, and inversion grid	partially reproduced. Generated geometry uses authoritative observations and an independent selection; the original 1489-event/52-station cohort remains unidentified. CMW1 historical elevation interpretation and conflicting borehole wellhead tables are documented. This input-geometry panel does not establish an inferred velocity model.
5a	Hayward event map: NCSN catalog	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate.

Panel	Quantity	Status and limiting dependency
5b	Hayward event map: standard tomography	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.
5c	Hayward event map: DD location	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
5d	Hayward event map: DD tomography	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.
6a	catalog: geographical map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
6b	catalog: rotated Cartesian map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
6c	catalog: across-strike depth section AA-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
6d	catalog: along-strike depth section BB-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
7a	standard tomography: geographical map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
7b	standard tomography: rotated Cartesian map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
7c	standard tomography: across-strike depth section AA-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
7d	standard tomography: along-strike depth section BB-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
8a	DD location: geographical map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
8b	DD location: rotated Cartesian map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
8c	DD location: across-strike depth section AA-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
8d	DD location: along-strike depth section BB-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
9a	DD tomography: geographical map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
9b	DD tomography: rotated Cartesian map	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
9c	DD tomography: across-strike depth section AA-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
9d	DD tomography: along-strike depth section BB-prime	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Panel	Quantity	Status and limiting dependency
10a (y=-30)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=-20)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=-10)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=0)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=10)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=20)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10a (y=30)	Hayward across-strike DD velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=-30)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=-20)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=-10)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=0)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=10)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=20)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10b (y=30)	Hayward across-strike standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=-30)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=-20)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=-10)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=0)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=10)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=20)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
10c (y=30)	Hayward across-strike DD minus standard velocity	partially reproduced. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Panel	Quantity	Status and limiting dependency
11a (y=-30)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=-20)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=-10)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=0)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=10)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=20)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11a (y=30)	Hayward resolution-matrix diagonal	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=-30)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=-20)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=-10)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=0)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=10)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=20)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Panel	Quantity	Status and limiting dependency
11b (y=30)	Hayward interpolated nodal conditional velocity noise from independent picks	not reproduced. Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

B.1 Specifications, commands, and attempted dependencies

The following records specify the source description, parameters, generation command, and plot command for each panel. Seven sections of a common Figure 10 or 11 panel share the explicitly listed settings. Unavailable commands are reported as unavailable rather than replaced by placeholders that would fabricate field observations.

Panel 1

Quantity: Horizontal true P-velocity slice
Written specification: Synthetic Test, p.1877; caption 1
Parameters: Written sandwich profile, x=-35..35 km, y=-60..40 km; constant with depth

Status: independently reproduced
Attempt and limitation: Physical velocity profile reconstructed from prose; plotting extent and depth chosen from model domain.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic
```

Create plot:

```
python3 plot.py --figure 1
```

Panel 2

Quantity: Synthetic-test event/station geometry and inversion grid
Written specification: Caption 2, p.1879
Parameters: Published 8x6x5 inversion grid; 120 independent sources and 32 independent stations, seed 2003
Status: partially reproduced
Attempt and limitation: The saved baseline uses random geometry. A documented authoritative Loma Prieta subset is authorized; it has not yet replaced this experiment. No digitization used.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic
```

Create plot:

```
python3 plot.py --figure 2
```

Panel 3a

Quantity: Standard tomographic P velocity; all generated depth slices
Written specification: Synthetic Test pp.1877-1879; captions 3 pp.1880-1881
Parameters: Depths 0,3,7,11,16 km; grid from caption 2; noise 0.04 s and station bias +/-0.3 s; configs/synthetic.cfg
Status: partially reproduced
Attempt and limitation: Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic && ./build/dd invert configs/synthetic.cfg data/synthetic
    standard
```

Create plot:

```
python3 plot.py --figure 3a
```

Panel 3b

Quantity: DD tomographic P velocity; all generated depth slices
Written specification: Synthetic Test pp.1877-1879; captions 3 pp.1880-1881
Parameters: Depths 0,3,7,11,16 km; grid from caption 2; noise 0.04 s and station bias +/-0.3 s; configs/synthetic.cfg
Status: partially reproduced
Attempt and limitation: Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic && ./build/dd invert configs/synthetic.cfg data/synthetic
    dd
```

Create plot:

```
python3 plot.py --figure 3b
```

Panel 3c

Quantity: True minus DD P velocity; all generated depth slices
Written specification: Synthetic Test pp.1877-1879; captions 3 pp.1880-1881
Parameters: Depths 0,3,7,11,16 km; grid from caption 2; noise 0.04 s and station bias +/-0.3 s; configs/synthetic.cfg
Status: partially reproduced
Attempt and limitation: Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic && ./build/dd invert configs/synthetic.cfg data/synthetic
    dd
```

Create plot:

```
python3 plot.py --figure 3c
```

Panel 3d

Quantity: True minus standard P velocity; all generated depth slices
Written specification: Synthetic Test pp.1877-1879; captions 3 pp.1880-1881
Parameters: Depths 0,3,7,11,16 km; grid from caption 2; noise 0.04 s and station bias +/-0.3 s; configs/synthetic.cfg
Status: partially reproduced
Attempt and limitation: Independent geometry, homogeneous initial velocity, and explicit replacements for unpublished numerical settings; numerical comparisons cannot isolate implementation from experiment differences.
Generate data:

```
./build/dd generate configs/synthetic.cfg data/synthetic && ./build/dd invert configs/synthetic.cfg data/synthetic
    standard
```

Create plot:

```
python3 plot.py --figure 3d
```

Panel 4

Quantity: Hayward network, catalog seismicity, and inversion grid
Written specification: Caption 4, p.1882
Parameters: 1092 independently selected current catalog earthquakes, 37 physical stations, 77 horizontal grid nodes; identified source and receiver inputs in data/hayward
Status: partially reproduced
Attempt and limitation: Generated geometry uses authoritative observations and an independent selection; the original 1489-event/52-station cohort remains unidentified. CMW1 historical elevation interpretation and conflicting borehole wellhead tables are documented. This input-geometry panel does not establish an inferred velocity model.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py --geometry-only
```

Panel 5a

Quantity: Hayward event map: NCSN catalog
Written specification: Event Locations pp.1881-1883; caption 5
Parameters: 1489 earthquakes, 52 stations; 20257 absolute picks, 767127 catalog differences, 17955 precise differences
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 5b

Quantity: Hayward event map: standard tomography
Written specification: Event Locations pp.1881-1883; caption 5
Parameters: 1489 earthquakes, 52 stations; 20257 absolute picks, 767127 catalog differences, 17955 precise differences
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 5c

Quantity: Hayward event map: DD location
Written specification: Event Locations pp.1881-1883; caption 5
Parameters: 1489 earthquakes, 52 stations; 20257 absolute picks, 767127 catalog differences, 17955 precise differences
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward loc1d; ./
    build/dd raycheck configs/hayward.cfg data/hayward loc1d
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 5d

Quantity: Hayward event map: DD tomography
Written specification: Event Locations pp.1881-1883; caption 5
Parameters: 1489 earthquakes, 52 stations; 20257 absolute picks, 767127 catalog differences, 17955 precise differences
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 6a

Quantity: catalog: geographical map
Written specification: Event Locations pp.1882-1884; captions 6-9

Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 6b

Quantity: catalog: rotated Cartesian map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 6c

Quantity: catalog: across-strike depth section AA-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate.

Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 6d

Quantity: catalog: along-strike depth section BB-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Catalog coordinates are provider preferred-origin estimates; this panel contains no inverse estimate. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 7a

Quantity: standard tomography: geographical map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 7b

Quantity: standard tomography: rotated Cartesian map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 7c

Quantity: standard tomography: across-strike depth section AA-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 7d

Quantity: standard tomography: along-strike depth section BB-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 8a

Quantity: DD location: geographical map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward loc1d; ./
    build/dd raycheck configs/hayward.cfg data/hayward loc1d
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 8b

Quantity: DD location: rotated Cartesian map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward loc1d; ./
    build/dd raycheck configs/hayward.cfg data/hayward loc1d
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 8c

Quantity: DD location: across-strike depth section AA-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward loc1d; ./
    build/dd raycheck configs/hayward.cfg data/hayward loc1d
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 8d

Quantity: DD location: along-strike depth section BB-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Fixed-model relocation uses catalog differences with absolute-data anchoring; it is an independent counterpart of the published DD-location experiment. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward loc1d; ./
    build/dd raycheck configs/hayward.cfg data/hayward loc1d
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 9a

Quantity: DD tomography: geographical map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or

digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 9b

Quantity: DD tomography: rotated Cartesian map
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 9c

Quantity: DD tomography: across-strike depth section AA-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within

the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 9d

Quantity: DD tomography: along-strike depth section BB-prime
Written specification: Event Locations pp.1882-1884; captions 6-9
Parameters: Center 37.72 N, 122.06 W; 35-degree anticlockwise rotation; map selection 37.83..37.91 N, -122.26..-122.18 E
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Zoom membership is fixed by the provider catalog coordinates within the written geographic window. AA/BB section-box boundaries and fault traces were not recovered or digitized; both depth views include every selected zoom event, with common coordinate limits across methods.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 10a: all seven sections at y=-30,-20,-10,0,10,20,30 km

Quantity: Hayward across-strike DD velocity
Written specification: Velocity Structure p.1884; caption 10 p.1887
Parameters: Published Hayward grid; smoothing weight 10; all seven y sections listed above; hypocenters within 5 km
Status: partially reproduced
Attempt and limitation: Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against

independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard; ./build/dd invert configs/hayward.cfg data/
    hayward dd; ./build/dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py --velocity-only
```

Panel 10b: all seven sections at y=-30,-20,-10,0,10,20,30 km

Quantity: Hayward across-strike standard velocity
Written specification: Velocity Structure p.1884; caption 10 p.1887
Parameters: Published Hayward grid; smoothing weight 10; all seven y sections listed above; hypocenters within 5 km
Status: partially reproduced
Attempt and limitation: Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard; ./build/dd invert configs/hayward.cfg data/
    hayward dd; ./build/dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py --velocity-only
```

Panel 10c: all seven sections at y=-30,-20,-10,0,10,20,30 km

Quantity: Hayward across-strike DD minus standard velocity
Written specification: Velocity Structure p.1884; caption 10 p.1887
Parameters: Published Hayward grid; smoothing weight 10; all seven y sections listed above; hypocenters within 5 km
Status: partially reproduced
Attempt and limitation: Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. Seven written y sections are generated from the saved standard and catalog-DD models, with hypocenters within 5 km. Sections include the full saved depth range, including the above-geoid layer added to contain receivers and any source estimates at its top. Absolute-data tomography uses the independently adapted continuous starting profile and numerical settings. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs. Velocity-only checkpoint: final model ray diagnostics and fixed-model relocation remain pending. The original standard solve stopped after a rejected line search, with 572 of 14262 rays failing its last pre-update local stopping check. The DD run exhausted its 18 configured updates; its last accepted maximum source-coordinate change was 2.7614 km. These panels visualize saved calculations without establishing nonlinear convergence or model accuracy.
Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward standard;
    ./build/dd raycheck configs/hayward.cfg data/hayward standard; ./build/dd invert configs/hayward.cfg data/
    hayward dd; ./build/dd raycheck configs/hayward.cfg data/hayward dd
```

Create plot:

```
python3 plot_hayward.py --velocity-only
```

Panel 11a: all seven sections at y=-30,-20,-10,0,10,20,30 km

Quantity: Hayward resolution-matrix diagonal
Written specification: Velocity Structure pp.1884,1886; caption 11 p.1888
Parameters: Absolute catalog-P Jacobian at saved DD sources/velocity; source parameters eliminated; independent log-slowness smoothing and column-scaled damping; y=-30
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the

original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd; ./build/dd linearize configs/hayward.cfg data/hayward dd;
    OPENBLAS_NUM_THREADS=1 python3 resolution_hayward.py
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

Panel 11b: all seven sections at y=-30,-20,-10,0,10,20,30 km

Quantity: Hayward interpolated nodal conditional velocity noise from independent picks
Written specification: Velocity Structure pp.1884,1886; caption 11 p.1888
Parameters: Absolute catalog-P Jacobian at saved DD sources/velocity; source parameters eliminated; independent log-slowness smoothing and column-scaled damping; y=-30
Status: not reproduced
Attempt and limitation: Completed field figure metrics are not available yet; this panel has not been promoted to a completed counterpart. Independent subset: 1092 events, 37 physical stations, 14262 absolute P picks and 44718 catalog differences; the manuscript reports 1489, 52, 20257 and 767127, respectively. Its exact cohort remains unidentified, and its 17955 waveform differences are absent here. The generated resolution is the absolute-data velocity block at the catalog-DD estimate, with source parameters included in the joint algebra. It is not a full DD resolution matrix or the original simul2000 calculation. Noise propagation assumes reported symmetric pick uncertainties are independent 1-sigma errors; provider confidence levels are absent. Sections interpolate nodal diagnostics; they do not propagate covariance to interpolated velocity. Unconstrained source elimination omits the effect of active source-coordinate bounds. It omits correlated station/model errors, ray and nonlinear errors, selection effects and regularization bias. Small conditional noise does not establish small total velocity error. Joint catalog-DD tomography uses catalog differences with absolute-data anchoring and preserves both native channel epochs. Catalog displacement is not location error against independent truth. Source-derivative and ray-refinement checks sample the saved model; they do not establish all-ray or nonlinear inverse convergence. Residual reduction alone does not establish recovery or unchanged timing calibration across metadata epochs.

Generate data:

```
python3 prepare_hayward.py --selection reproduction; ./build/dd invert configs/hayward.cfg data/hayward dd; ./build
    /dd raycheck configs/hayward.cfg data/hayward dd; ./build/dd linearize configs/hayward.cfg data/hayward dd;
    OPENBLAS_NUM_THREADS=1 python3 resolution_hayward.py
```

Create plot:

```
python3 plot_hayward.py # complete field pipeline: ./rebuild_hayward.sh
```

C References and guidance

Zhang, H., and C. H. Thurber (2003). Double-Difference Tomography: The Method and Its Application to the Hayward Fault, California. *BSSA*, 93, 1875–1889. doi:10.1785/0120020190.

Oppenheimer, D. H., F. W. Klein, J. P. Eaton, and F. W. Lester (1993). The Northern California Seismic Network bulletin, January–December 1992. USGS Open-File Report 93-578, Table 4. doi:10.3133/ofr93578.

Waldhauser, F., and W. L. Ellsworth (2002). Fault structure and mechanics of the Hayward Fault, California, from double-difference earthquake locations. *Journal of Geophysical Research*, 107(B3), ESE 3-1–ESE 3-15. doi:10.1029/2000JB000084.

The two user-permitted guidance files are repository guidelines and academic style guide. They govern research-code simplicity and prose. They were not used as permission to inspect any original model implementation or output.