


Chapter 5

Divide and Conquer



*Slides by Kevin Wayne.
Copyright © 2005 Pearson-Addison Wesley.
All rights reserved.*

Divide-and-Conquer

Divide-and-conquer.

- Break up problem into several parts.
- Solve each part recursively.
- Combine solutions to sub-problems into overall solution.

Most common usage.

- Break up problem of size n into *two* equal parts of size $\frac{1}{2}n$.
- Solve two parts recursively.
- Combine two solutions into overall solution in *linear time*.

Consequence.

- Brute force: n^2 .
- Divide-and-conquer: $n \log n$.

*Divide et impera.
Veni, vidi, vici.
- Julius Caesar*

5.3 Counting Inversions

Counting Inversions


Music site tries to match your song preferences with others.

- You rank n songs.
- Music site consults database to find people with **similar** tastes.

Similarity metric: number of inversions between two rankings.

- My rank: $1, 2, \dots, n$.
- Your rank: a_1, a_2, \dots, a_n .
- Songs i and j **inverted** if $i < j$, but $a_i > a_j$.

Songs					
	\mathcal{A}	\mathcal{B}	\mathcal{C}	\mathcal{D}	\mathcal{E}
Me	1	2	3	4	5
You	1	3	4	2	5



Inversions

3-2, 4-2

Brute force: check all $\Theta(n^2)$ pairs i and j .

Applications

Applications.

- *Voting theory*
- *Collaborative filtering for people with similar preferences (suggestions in web)*
- *Measuring the "sortedness" of an array.*
- *Sensitivity analysis of Google's ranking function.*
- *Rank aggregation for meta-searching on the Web.*
- *Nonparametric statistics (e.g., Kendall's Tau distance).*

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- *Divide*: separate list into two pieces.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- *Divide:* separate list into two pieces.
- *Conquer:* recursively count inversions in each half.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Conquer: $2T(n/2)$

5 blue-blue inversions

8 green-green inversions

5-4, 5-2, 4-2, 8-2, 10-2

6-3, 9-3, 9-7, 12-3, 12-7, 12-11, 11-3, 11-7

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- *Divide*: separate list into two pieces.
- *Conquer*: recursively count inversions in each half.
- *Combine*: count inversions where a_i and a_j are in different halves, and return sum of *three* quantities.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Conquer: $2T(n/2)$

5 *blue-blue* inversions

8 *green-green* inversions

9 *blue-green* inversions

5-3, 4-3, 8-6, 8-3, 8-7, 10-6, 10-9, 10-3, 10-7


Combine: ???

$$\text{Total} = 5 + 8 + 9 = 22.$$

Counting Inversions: Combine

Combine: count blue-green inversions

- Assume each half is *sorted*.
- Count inversions where a_i and a_j are in different halves.
- Merge* two sorted halves into sorted whole.



to maintain sorted invariant

3	7	10	14	18	19
---	---	----	----	----	----

2	11	16	17	23	25
6	3	2	2	0	0

13 blue-green inversions: $6 + 3 + 2 + 2 + 0 + 0$

Count: $O(n)$

2	3	7	10	11	14	16	17	18	19	23	25
---	---	---	----	----	----	----	----	----	----	----	----

Merge: $O(n)$

$$T(n) \leq T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + O(n) \Rightarrow T(n) = O(n \log n)$$

Counting inversions: divide-and-conquer

- Divide: separate list into two halves A and B .
- Conquer: recursively count inversions in each list.
- Combine: count inversions (a, b) with $a \in A$ and $b \in B$.
- Return sum of three counts.

input

1	5	4	8	10	2	6	9	3	7
---	---	---	---	----	---	---	---	---	---

count inversions in left half A

1	5	4	8	10
---	---	---	---	----

5-4

count inversions in right half B

2	6	9	3	7
---	---	---	---	---

6-3 9-3 9-7

count inversions (a, b) with $a \in A$ and $b \in B$

1	5	4	8	10
---	---	---	---	----

2	6	9	3	7
---	---	---	---	---

4-2 4-3 5-2 5-3 8-2 8-3 8-6 8-7 10-2 10-3 10-6 10-7 10-9

output $1 + 3 + 13 = 17$

Counting inversions: how to combine two subproblems?

Q. How to count inversions (a, b) with $a \in A$ and $b \in B$?

A. Easy if A and B are sorted!

Warmup algorithm.

- Sort A and B .
- For each element $b \in B$,
 - binary search in A to find how elements in A are greater than b .

list A

7	10	18	3	14
---	----	----	---	----

list B

17	23	2	11	16
----	----	---	----	----

sort A

3	7	10	14	18
---	---	----	----	----

sort B

2	11	16	17	23
---	----	----	----	----

binary search to count inversions (a, b) with $a \in A$ and $b \in B$

3	7	10	14	18
---	---	----	----	----

2	11	16	17	23
---	----	----	----	----

5 2 1 1 0

Counting inversions: how to combine two subproblems?

Count inversions (a, b) with $a \in A$ and $b \in B$, assuming A and B are sorted.

- Scan A and B from left to right.
- Compare a_i and b_j .
- If $a_i < b_j$, then a_i is not inverted with any element left in B .
- If $a_i > b_j$, then b_j is inverted with every element left in A .
- Append smaller element to sorted list C .



count inversions (a, b) with $a \in A$ and $b \in B$



merge to form sorted list C



Counting Inversions: Implementation

Pre-condition. [Merge-and-Count] \mathcal{A} and \mathcal{B} are sorted.

Post-condition. [Sort-and-Count] \mathcal{L} is sorted.

```
Sort-and-Count(L) {  
    if list L has one element  
        return 0 and the list L  
  
    Divide the list into two halves A and B  
    ( $r_A$ , A)  $\leftarrow$  Sort-and-Count(A)  
    ( $r_B$ , B)  $\leftarrow$  Sort-and-Count(B)  
    ( $r$ , L)  $\leftarrow$  Merge-and-Count(A, B)  
  
    return  $r = r_A + r_B + r$  and the sorted list L  
}
```