

IGOR Final Report

IGOR Final Report

Executive Summary

This research initiative aimed to optimize the PURE cell-free protein synthesis system to maximize both protein yield and reaction rate. The IGOR process began by synthesizing existing literature, which confirmed that system performance is a complex interplay between the concentrations of translation machinery and a robust energy regeneration system. The process identified a key gap in understanding the trade-off between the initial reaction rate and the final protein yield. Through a competitive debate process, the most compelling hypothesis emerged, proposing that this trade-off defines a Pareto frontier. This concept was further evolved to integrate specific data-driven insights about ionic instabilities, leading to a comprehensive final hypothesis: the PURE system's performance is governed by a Rate-Yield Pareto frontier whose shape and stability are critically dependent on ionic conditions, particularly the Magnesium-to-Potassium ratio. An experimental plan has been designed to first map this frontier under stable conditions and then probe the predicted regions of instability.

Top-Ranked Hypotheses

1: The PURE System Pareto Frontier: Mapping the Rate-Yield Trade-off and Identifying Regions of Ionic Instability

Hypothesis: Hypothesis: The trade-off between protein synthesis rate and final yield in the PURE system defines a Pareto frontier, where improving one objective necessitates a compromise in the other. We posit that the shape and stability of this frontier are critically dependent on ionic conditions. Specifically, the bifurcation in performance observed at certain Magnesium-to-Potassium ratios represents a region of high sensitivity or instability on this frontier, likely near the 'knee' where the trade-off is most acute. Other regions of the frontier, such as the rate-dominant and yield-dominant extremes, are expected to be more robust to ionic variations.

Rationale: This hypothesis evolves the winning ‘Pareto Frontier’ concept by integrating the specific, data-driven insight from the ‘Bifurcation’ hypothesis. It provides a unified framework where the narrow phenomenon of bifurcation is explained as a predictable feature of the broader multi-objective landscape. Instead of simply identifying two distinct optimal regimes, this evolved hypothesis aims to map the entire operational landscape and its stability, providing a much more powerful engineering tool for optimizing the PURE system under different constraints.

Testable Question: Can we map a Pareto frontier by systematically varying the ratio of catalytic machinery ([PMix] + [Ribosome]) to the energy regeneration substrate ([Creatine Phosphate])? Furthermore, by performing this mapping at different background Magnesium-to-Potassium ratios (e.g., a ‘stable’ 1:20 ratio vs. the ‘unstable’ 1:12.5 ratio), can we demonstrate that the frontier’s shape changes and the variance in outcomes increases significantly in the unstable ionic regime, particularly at the frontier’s knee? **Final Rationale:** This hypothesis was top-ranked because it provides the most comprehensive and predictive model for PURE system optimization. It evolved by successfully integrating the superior conceptual framework of a ‘Pareto Frontier’ (from Explorer 1), which won the debates for its novelty and engineering utility, with the specific, data-driven insight of ionic instability causing performance bifurcations (from Quant 1). This synthesis creates a more robust and verifiable model that elegantly explains observed outliers not as random errors, but as predictable features of a broader, multi-objective landscape.

Key Supporting Evidence: - Rationale from debate Explorer 1 vs. Quant 1: The bifurcation phenomenon can be understood as a special case within a broader trade-off landscape. - Rationale from debate Explorer 1 vs. Architect 1: The ‘Pareto frontier’ is a more novel and powerful conceptual framing for the rate-yield trade-off. - Finding: A primary factor limiting protein synthesis is the depletion of the energy source and accumulation of byproducts. (Shimizu et al., 2001) - Data Observation: Outlier experiments (PMix_(AR-931)_QC_PURE_Timecourse_D8, PMix-AR-898_B12) show divergent outcomes at nearly identical Mg:K ratios.

2: PURE System Dynamics as Competing Control Loops: Feed-Forward Synthesis vs. Feedback Energy Regeneration

Hypothesis: Hypothesis: The PURE system’s rate-versus-yield trade-off can be modeled as a competition between two interconnected control loops: a ‘feed-forward’ synthesis loop that consumes resources (NTPs) to produce protein, and a ‘feedback’ energy regeneration loop that senses resource depletion (NDPs) and replenishes the NTP pool.

The system's performance profile (high-rate/low-yield vs. low-rate/high-yield) is determined by the kinetic balance and relative 'gain' of these two loops.

Rationale: This hypothesis evolves the original 'Rate vs. Yield Stoichiometry Trade-off' by introducing a more novel and mechanistic framework from control systems engineering. It addresses the weakness identified in the debates—a less powerful conceptual framing than the 'Pareto Frontier'—by providing a clear, causal dynamic model. A high-gain feed-forward loop (high concentrations of Ribosomes, PMix) results in a high initial rate but rapidly overwhelms the feedback loop's capacity, causing a system crash from energy depletion. Conversely, a high-gain feedback loop (high concentration of Creatine Phosphate) sustains the energy supply, enabling a higher final yield at the cost of a potentially lower initial rate.

Testable Question: Can we demonstrate that the dominant performance objective shifts from maximizing sigmoid_rate to maximizing sigmoid_steady_state as the stoichiometric ratio of the feedback loop's key substrate ([Creatine Phosphate]) to the feed-forward loop's key catalysts ([Ribosome] + [PMix]) is increased? Does a model based on these competing loops accurately predict the reaction's inflection point and termination time? **Final Rationale:** This hypothesis was highly ranked due to its significant evolution during the research cycle. Its predecessor, Architect 1, was scientifically sound but lost in debate due to a less compelling conceptual framework. This evolved version addressed that weakness by reframing the rate-yield trade-off using a novel control theory analogy (competing feed-forward/feedback loops). This provided a more sophisticated, mechanistic explanation for the system dynamics, making it conceptually distinct and powerful. **Key Supporting Evidence:** - Rationale from debate Explorer 1 vs. Architect 1: The original hypothesis, while scientifically sound, lacked a novel conceptual framing. - Finding: Energy depletion is a key factor that terminates the reaction. (Shimizu et al., 2001) - Finding: Increasing translation factor concentrations boosts performance. (Li et al., 2014)

Proposed Experimental Plan

Objective

To experimentally validate the hypothesis 'The PURE System Pareto Frontier' by first mapping the rate-yield trade-off under stable ionic conditions and then demonstrating that specific ionic ratios introduce instability, particularly at the frontier's most acute trade-off point.

Phase 1: Mapping the Pareto Frontier under Stable Ionic Conditions

Goal: To establish the baseline shape of the Pareto frontier between reaction rate and total yield.

1. Independent Variables: - **Catalytic Machinery Ratio:** This will be a combined variable of [PMix] (mg/mL) and [Ribosome] (uM). We will test 4 levels, maintaining a constant ratio between them, spanning the lower to upper bounds (e.g., [1.0 mg/mL PMix, 1.0 uM Ribo], [1.6, 1.8], [2.2, 3.2], [2.8, 4.5]). - **Energy Substrate ([Creatine phosphate] (mM)):** We will test 4 levels across the experimental range: 0, 25, 50, 100 mM.

2. Controlled Variables: - **Ionic Condition (Stable):** [Magnesium acetate] (mM) will be set to 7 mM and [Potassium glutamate] (mM) will be set to 140 mM (a 1:20 ratio, distant from the observed unstable point). - **DNA Template:** [DNA] (nM) will be held constant at a non-limiting concentration, e.g., 5 nM. - **Other components (ATP, GTP, etc.):** Set to the mean of their respective bounds as per the research goal.

3. Experimental Design: - A 4x4 full factorial design will be executed, resulting in 16 unique conditions. - Each condition will be run in triplicate (n=3) to assess variability. - A standard control reaction (e.g., conditions from experiment PMix-MFG-128-PURE-Activity-QC_F2) will be included on each plate for normalization.

4. Key Metrics: - Real-time fluorescence monitoring of protein synthesis. - Calculation of sigmoid_rate (1/h) and sigmoid_steady_state (ng/uL) for each run.

5. Expected Outcome: - A scatter plot of sigmoid_rate vs. sigmoid_steady_state that clearly delineates a curve representing the Pareto frontier, showing that conditions promoting high rates result in lower yields, and vice-versa.

Phase 2: Probing Instability on the Frontier

Goal: To test if the previously identified 'unstable' ionic ratio increases performance variability at key points along the established frontier.

1. Selection of Conditions: - From the Phase 1 frontier, select 3 representative points: 1. **High-Rate Extreme:** The condition with the highest rate. 2. **Knee Point:** The condition representing the best compromise (e.g., highest product of normalized rate and yield). 3. **High-Yield Extreme:** The condition with the highest yield.

2. Independent Variable: - Ionic Condition (Unstable): [Magnesium acetate] (mM) will be set to 8 mM and [Potassium glutamate] (mM) will be set to 100 mM (the 1:12.5 ratio identified from outliers).

3. Experimental Design: - Run the 3 selected machinery/energy conditions with the 'unstable' ionic background. - Increase the number of replicates to at least n=6 for each condition to robustly measure variance.

4. Key Metrics: - Same as Phase 1. - Calculate the Coefficient of Variation (CV) for both rate and yield for each condition.

5. Expected Outcome: - A significant increase in the CV for both rate and yield under the 'unstable' ionic condition compared to the 'stable' condition from Phase 1. This effect is hypothesized to be most pronounced at the 'Knee Point'. This would confirm that the bifurcation is a region of instability on the broader Pareto frontier.

References

- Shimizu, Y., Inoue, A., Tomari, Y., Suzuki, T., Yokogawa, T., Nishikawa, K., & Ueda, T. (2001). Cell-free translation reconstituted with purified components. *Nature Biotechnology*, 19(8), 751-755.
- Li, J., Gu, L., Aach, J., & Church, G. M. (2014). Improved Cell-Free RNA and Protein Synthesis System. *PLOS ONE*, 9(9), e106232.
- Lavickova, B., & Maerkl, S. J. (2019). A Simple, Robust, and Low-Cost Method To Produce the PURE Cell-Free System. *ACS Synthetic Biology*, 8(2), 455-462.
- Kurz, L., et al. (2025). ATP Regeneration from Pyruvate in the PURE System. *ACS Synthetic Biology*.
- Matsuura, T., Hosoda, K., & Shimizu, Y. (2018). Robustness of a Reconstituted Escherichia coli Protein Translation System Analyzed by Computational Modeling. *ACS Synthetic Biology*, 7(8), 1964-1972.

Dead Ends

- Hypothesis: 'Bifurcation of PURE System Performance at a Critical Magnesium-to-Potassium Ratio' (Quant 1). Reason for Rejection: While this hypothesis correctly identified a critical data anomaly, it was deemed too narrow in scope during the debates. The phenomenon was more powerfully explained as a special case of 'ionic instability' within the broader, more fundamental framework of a 'Pareto Frontier,' making it a symptom rather than the root cause.
- Hypothesis: 'Rate vs. Yield Stoichiometry Trade-off' (Architect 1 -

original version). Reason for Rejection: This hypothesis was scientifically sound but was defeated in debate by the 'Pareto Frontier' concept, which was judged to be a more novel, powerful, and actionable conceptual framework for the same underlying principle. The original hypothesis was effectively superseded by its more sophisticated competitor before being evolved.