

# A note on a proof of Zorn's Lemma from Axiom of Choice without transfinite induction

Koji Nuida

December 8, 2011

## Abstract

In this note, we give a proof of Zorn's Lemma from Axiom of Choice without transfinite induction (whose essential idea is the same as the proof of Theorem 4.19 in the book [H. Rubin and J. E. Rubin, "Equivalents of the Axiom of Choice, II", Second Edition, Studies in Logic and the Foundations of Mathematics vol.116, North-Holland, 1985], as I noticed after writing the note ...). For the sake of comparison, we also include as an appendix a proof of the claim using transfinite induction.

Throughout this note, let  $(X, \leq)$  denote an arbitrary non-empty partially ordered set in which every chain (linearly or totally ordered subset)  $C$  has an upper bound (with respect to  $\leq$ ), that is, an element  $x \in X$  satisfying that  $c \leq x$  for every  $c \in C$ . Then Zorn's Lemma states that such an  $X$  always has a maximal element (with respect to  $\leq$ ), that is, an element  $x \in X$  satisfying that there exist no elements  $y \in X$  with  $x < y$ . In this note, we give a proof of Zorn's Lemma from Axiom of Choice (in Zermelo–Fraenkel set theory), without transfinite induction which would be used in a “natural” proof of the claim.

Assume, for the contrary, that an  $X$  satisfying the hypothesis has no maximal elements. We will derive a contradiction from this assumption.

We summarize some basic definitions and terminologies:

- We say that a chain  $C$  in a subset  $Y$  of  $X$  is *bounded in  $Y$*  if there exists an element  $x \in Y \setminus C$  satisfying that  $c < x$  for every  $c \in C$ , and *unbounded in  $Y$*  otherwise.
- A chain  $C$  in  $X$  is called *well-ordered* if any non-empty subset of  $C$  has a unique minimal element.
- For a chain  $C$  and an element  $c \in C$ , the subset  $\{d \in C \mid d < c\}$  is called the *initial segment* of  $C$  relative to  $c$  and is denoted by  $s_C(c)$ .

Let  $\mathcal{C}$  denote the set of all non-empty chains in  $X$ . For each  $C \in \mathcal{C}$ , define  $U_C = \{x \in X \setminus C \mid c < x \text{ for every } c \in C\}$ . We notice the following property:

**Lemma 1.** *We have  $U_C \neq \emptyset$  for every  $C \in \mathcal{C}$ .*

*Proof.* By the hypothesis of Zorn's Lemma,  $C$  has an upper bound  $x \in X$ . If  $y \in U_{\{x\}}$ , then we have  $c \leq x < y$  (hence  $c < y$ ) for every  $c \in C$ , therefore  $y \notin C$  and  $y \in U_C$ . Hence we have  $U_{\{x\}} \subset U_C$ , while  $U_{\{x\}} \neq \emptyset$  by the assumption that  $X$  has no maximal elements, therefore we have  $U_C \neq \emptyset$ .  $\square$

We would like to construct (by using Axiom of Choice) a choice function  $f$  of the family  $\{U_C\}_{C \in \mathcal{C}}$  which satisfies some more appropriate properties described below.

For an arbitrary pair of  $C_1, C_2 \in \mathcal{C}$ , we define a symmetric relation  $C_1 \sim_{\text{pre}} C_2$  to mean that the chain  $C_1 \cap C_2$  is unbounded in both  $C_1$  and  $C_2$ . Then we define an equivalence relation  $\sim$  on  $\mathcal{C}$  to be the transitive closure of  $\sim_{\text{pre}}$ , that is, we have  $C_1 \sim C_2$  if and only if there exists a finite sequence  $C_1 = C'_0, C'_1, \dots, C'_n = C_2$  ( $n \geq 0$ ) of elements of  $\mathcal{C}$  with  $C'_i \sim_{\text{pre}} C'_{i+1}$  for every  $0 \leq i \leq n-1$ . Note that if  $C_1 \sim C_2$  and  $C_1$  has a maximum element  $c$ , then  $c \in C_2$  and  $c$  is also a maximum element of  $C_2$ . Now we present the following lemma:

**Lemma 2.** *If  $C_1, C_2 \in \mathcal{C}$  and  $C_1 \sim C_2$ , then  $U_{C_1} = U_{C_2}$ .*

*Proof.* It suffices to show that  $U_{C_1} \subset U_{C_2}$  if  $C_1 \sim_{\text{pre}} C_2$ . Let  $x \in U_{C_1}$ . If  $x \in C_2$ , then the upper bound  $x$  ( $\in C_2 \setminus (C_1 \cap C_2)$ ) of  $C_1$  is also an upper bound of  $C_1 \cap C_2$ , therefore  $C_1 \cap C_2$  is bounded in  $C_2$ , contradicting the hypothesis  $C_1 \sim_{\text{pre}} C_2$ . Hence  $x \notin C_2$ . If  $c \in C_2$  and  $c \not\leq x$ , then, as  $C_1 \cap C_2$  is unbounded in  $C_2$ , there exists a  $d \in C_1 \cap C_2$  with  $c \leq d \not\leq x$ . Now we have  $d \in C_1$  and  $d \not\leq x$ , contradicting the property  $x \in U_{C_1}$ . Hence we have  $c < x$  for every  $c \in C_2$ , therefore  $x \in U_{C_2}$ .  $\square$

We define a subset  $U_{\mathcal{E}}$  of  $X$  for each  $\sim$ -equivalence class  $\mathcal{E}$  by  $U_{\mathcal{E}} = U_C$ , where  $C \in \mathcal{E}$ . This is well-defined by virtue of Lemma 2. Now we obtain (by using Axiom of Choice) a choice function  $f$  of the family  $\{U_{\mathcal{E}}\}_{\mathcal{E} \in X/\sim}$ . Moreover, for simplicity, we write  $f(C) = f([C])$  for any  $C \in \mathcal{C}$ , where  $[C]$  denotes the  $\sim$ -equivalence class of  $C$ . Hence  $f(C) \in U_C$  for any  $C \in \mathcal{C}$ . Moreover, if  $C_1, C_2 \in \mathcal{C}$  and  $C_1 \sim C_2$ , then we have  $f(C_1) = f(C_2)$ .

From now, we fix an element  $x_0 \in X$ . We give the following definition:

- We say that a  $C \in \mathcal{C}$  is *f-consecutive* if  $C$  is well-ordered,  $x_0$  is the minimum element of  $C$ , and we have  $f(s_C(c)) = c$  for any  $c \in C \setminus \{x_0\}$ .

We present some properties of *f-consecutive* chains:

**Lemma 3.** *Let  $C \in \mathcal{C}$  be f-consecutive, and suppose that  $\emptyset \neq C' \subset C$  and  $C'$  is bounded in  $C$ . Then we have  $f(C') = d$ , where  $d$  is the unique minimal element of the set of all upper bounds  $x \in C \setminus C'$  of  $C'$ .*

*Proof.* Note that  $C' \in \mathcal{C}$ . Let  $d$  be as specified in the statement, which indeed exists by virtue of the fact that  $C$  is well-ordered and  $C'$  is bounded in  $C$ . We show that  $s_C(d) \sim C'$ . First, we have  $C' \subset s_C(d)$ , therefore  $s_C(d) \cap C' = C'$  is unbounded in  $C'$ . On the other hand, if  $x \in s_C(d) \setminus C'$ , then  $x$  is not an upper bound of  $C'$  by the choice of  $d$ . Therefore  $C'$  is unbounded in  $s_C(d)$ . Hence we

have  $s_C(d) \sim C'$ . Now we have  $f(C') = f(s_C(d))$  by the property of  $f$ , while we have  $f(s_C(d)) = d$ , as  $C$  is  $f$ -consecutive. This implies that  $f(C') = d$ .  $\square$

**Lemma 4.** *Let  $C_1, C_2 \in \mathcal{C}$  be two  $f$ -consecutive chains. Then exactly one of the following three conditions is satisfied:*

1.  $C_1 = C_2$ ;
2.  $C_1$  is an initial segment of  $C_2$ ;
3.  $C_2$  is an initial segment of  $C_1$ .

*Proof.* It is obvious that two of the three conditions specified in the statement do not occur simultaneously.

First we show that for each  $i \in \{1, 2\}$ , the conditions  $x \in C_1 \cap C_2$  and  $y \in s_{C_i}(x)$  imply  $y \in C_1 \cap C_2$ . Assume, for the contrary, that a counterexample  $y$  exists for some  $x$ , and let  $y$  be the minimal counterexample relative to the fixed  $x$  (which exists, as  $C_i$  is well-ordered). Then we have  $s_{C_i}(y) \neq \emptyset$ , as  $x_0 \in C_1 \cap C_2$ . By the choice of  $y$ , we have  $s_{C_i}(y) \subset C_{3-i}$  and  $y \notin C_{3-i}$ . We have  $f(s_{C_i}(y)) = y$ , as  $C_i$  is  $f$ -consecutive. On the other hand, as  $x \in C_1 \cap C_2$  and  $y < x$ , the subset  $s_{C_i}(y)$  of  $C_{3-i}$  is bounded in  $C_{3-i}$ , therefore we have  $f(s_{C_i}(y)) \in C_{3-i}$  by Lemma 3. But this contradicts the property  $y \notin C_{3-i}$ . Hence the claim of this paragraph holds. Moreover, as  $C_1$  and  $C_2$  are well-ordered, the above fact implies that  $C_1 \cap C_2$  is either equal to or an initial segment of  $C_i$  for each  $i \in \{1, 2\}$  (if  $C_1 \cap C_2 \neq C_i$ , consider the minimum element of  $C_i \setminus (C_1 \cap C_2)$ ).

Now assume, for the contrary, that  $C_1 \cap C_2 \neq C_1$  and  $C_1 \cap C_2 \neq C_2$ . Then  $C_1 \cap C_2$  is an initial segment of both  $C_1$  and  $C_2$ , therefore we have  $f(C_1 \cap C_2) \in C_1 \cap C_2$ , as  $C_1$  and  $C_2$  are  $f$ -consecutive. But this contradicts the definition of  $f$ . Hence we have  $C_1 \cap C_2 = C_i$  for some  $i \in \{1, 2\}$ , and now the above result implies that  $C_i = C_{3-i}$  or  $C_i$  is an initial segment of  $C_{3-i}$ .  $\square$

Let  $C_0$  denote the union of all  $f$ -consecutive chains (note that at least one  $f$ -consecutive chain, say  $\{x_0\}$ , exists). Then Lemma 4 implies that  $C_0 \in \mathcal{C}$  (for two elements  $x, y \in C_0$ , there exist  $f$ -consecutive chains  $C$  containing  $x$  and  $C'$  containing  $y$ , and  $x \in C \subset C'$  or  $y \in C' \subset C$  by the lemma). Now we present the following lemma:

**Lemma 5.** *Let  $c \in C_0$ , and let  $C$  be an  $f$ -consecutive chain containing  $c$ . Then  $s_{C_0}(c) = s_C(c)$ .*

*Proof.* First note that  $s_C(c) \subset s_{C_0}(c)$ , as  $C \subset C_0$  by the definition of  $C_0$ . We show that  $d \in C$  for an arbitrary element  $d \in s_{C_0}(c)$ . By the definition of  $C_0$ , there exists an  $f$ -consecutive chain  $C'$  such that  $d \in C'$ . Now we have  $d \in C' \subset C$  if  $C' = C$  or  $C'$  is an initial segment of  $C$ . On the other hand, if  $C$  is an initial segment of  $C'$ , then the properties  $c \in C$ ,  $d \in C'$  and  $d < c$  imply  $d \in C$ . Hence we have  $d \in C$  by Lemma 4.  $\square$

From now, we will show that  $C_0$  itself is  $f$ -consecutive. It is obvious that  $x_0$  is the minimum element of  $C_0$ .

**Lemma 6.**  $C_0$  is well-ordered.

*Proof.* Let  $A$  be an arbitrary non-empty subset of  $C_0$ . Choose an element  $a \in A$ , and let  $C$  be an  $f$ -consecutive chain with  $a \in C$  (which exists by the definition of  $C_0$ ). As  $C$  is well-ordered,  $A \cap C$  has the minimum element, say  $a_0$ . Now if  $a' \in A$  and  $a' < a_0$ , then we have  $a' \in s_{C_0}(a_0) = s_C(a_0)$  by Lemma 5, therefore  $a' \in A \cap C$  and  $a' < a_0$ , contradicting the choice of  $a_0$ . Hence we have  $a' \geq a_0$  for every  $a' \in A$ , therefore  $A$  itself has the minimum element  $a_0$ .  $\square$

**Lemma 7.** We have  $f(s_{C_0}(c)) = c$  for any  $c \in C_0 \setminus \{x_0\}$ .

*Proof.* Let  $C$  be an  $f$ -consecutive chain with  $c \in C$  (which exists by the definition of  $C_0$ ). Then we have  $s_{C_0}(c) = s_C(c)$  by Lemma 5. As  $C$  is  $f$ -consecutive, now we have  $f(s_{C_0}(c)) = f(s_C(c)) = c$ .  $\square$

By Lemma 6 and Lemma 7, it follows that  $C_0$  is  $f$ -consecutive. Now it is straightforward to verify that  $C_0 \cup \{f(C_0)\}$  is also an  $f$ -consecutive chain, but it is not a subset of  $C_0$ , contradicting the definition of  $C_0$ . Hence  $X$  should have a maximal element, concluding the proof of Zorn's Lemma.

## Appendix: A proof using transfinite induction

In this appendix, for the sake of comparison, we describe a proof of Zorn's Lemma from Axiom of Choice using transfinite induction. First we clarify the statement of the principle for "definition by transfinite recursion" (see e.g., Theorem 9.3 in Chapter I of [K. Kunen, "SET THEORY, An Introduction to Independence Proofs", Elsevier, 1980]):

**Theorem 1.** Let  $\varphi(x, y)$  be a formula (in Zermelo–Fraenkel set theory) with free variables  $x, y$  satisfying  $\forall x \exists! y \varphi(x, y)$ . Then there exists a formula  $\Phi(x, y)$  with free variables  $x, y$  satisfying the following two conditions;

1.  $\forall x ((x \in \mathbf{ON} \rightarrow \exists! y \Phi(x, y)) \wedge (\neg x \in \mathbf{ON} \rightarrow \neg \exists y \varphi(x, y)))$ ;
2.  $\forall x (x \in \mathbf{ON} \rightarrow \forall y, z (y = \Phi \upharpoonright_x \wedge \varphi(y, z) \rightarrow \Phi(x, z)))$ ,

where " $x \in \mathbf{ON}$ " is an abbreviation of " $x$  is an ordinal number" and " $\Phi \upharpoonright_x$ " is an abbreviation of the set  $\{\langle a, b \rangle \mid a \in x \wedge \Phi(a, b)\}$  (with  $\langle a, b \rangle$  denoting the ordered pair of  $a$  and  $b$ ).

Intuitively, the theorem means that, if we would like to define a "function"  $\Phi$  with domain consisting of all ordinal numbers (the whole of which is never a set) in such a way that the value of  $\Phi$  at each ordinal number  $\alpha$  is determined by a given rule from the values of  $\Phi$  at ordinal numbers less than  $\alpha$ , then there indeed exists such a "function"  $\Phi$ . Note that this is a theorem of ZF set theory and does not depend on Axiom of Choice.

Now we give a proof of Zorn's Lemma from Axiom of Choice using Theorem 1 (as well as transfinite induction). Let  $X \neq \emptyset$  be a partially ordered set appeared

in the statement of Zorn's Lemma. Assume, for the contrary, that  $X$  has no maximal elements. Then, for each non-empty subset  $C$  of  $X$  which is isomorphic to an ordinal number (hence a chain), it follows from Axiom of Choice that there exists a distinguished upper bound  $b_C$  of  $C$  with  $b_C \in X \setminus C$ .

To apply Theorem 1, first we define a formula  $\varphi(x, y)$  in the following manner, where we fix an element  $a \in X$  throughout the proof:

- If  $x = 0$  ( $= \emptyset$ ), then let  $\varphi(x, y)$  mean that  $y = a$ .
- If  $x$  is a function from an ordinal number  $\alpha > 0$  to  $X$  which is an isomorphism (between partially ordered sets) onto the image  $\text{Im}(x)$  of  $x$ , then let  $\varphi(x, y)$  mean that  $y = b_{\text{Im}(x)}$  (note that  $\text{Im}(x)$  is isomorphic to the non-empty ordinal number  $\alpha$ , therefore  $b_{\text{Im}(x)}$  is indeed defined).
- Otherwise, let  $\varphi(x, y)$  mean that  $y = 0$ .

This formula  $\varphi(x, y)$  satisfies the hypothesis of Theorem 1, therefore a formula  $\Phi(x, y)$  as in the theorem exists. Now we have the following lemma:

**Lemma 8.** *Let  $x$  be an ordinal number, and let  $x'$  be the unique element satisfying  $\Phi(x, x')$ .*

1. *We have  $x' \in X$ .*
2. *If  $y < x$  and  $\Phi(y, y')$ , then  $y' < x'$  in  $X$ .*

*Proof.* We prove the claim by transfinite induction on  $x$ . First, if  $x = 0$ , then it follows from the definition of the formula  $\varphi$  that  $x' = a$ , therefore the specified conditions are satisfied. Secondly, suppose that  $x > 0$ . Then, by the hypothesis of the transfinite induction, the set  $\Phi \upharpoonright_x$  in the statement of Theorem 1 is an isomorphism from  $x$  to a subset of  $X$ , say,  $C$  (note that  $x$  is totally ordered). Now by the definitions of  $\Phi$  and  $\varphi$ , it follows that  $x' = b_C$ , therefore the specified conditions are satisfied for  $x$  (the second condition follows from the property that  $b_C \in X \setminus C$  is an upper bound of  $C$ ). Hence the claim holds.  $\square$

By the second property shown in Lemma 8, for each element  $v \in X$ , there exists at most one ordinal number  $x$  satisfying  $\Phi(x, v)$ . Let  $X'$  denote the subset of  $X$  defined in such a way that  $v \in X'$  if and only if  $v \in X$  and  $\Phi(x, v)$  for some (or equivalently, a unique) ordinal number  $x$ . By the Axiom Schema of Replacement applied to the set  $X'$  and the formula  $\Phi'(x, y) := \Phi(y, x)$ , there exists a set  $Y$  for which we have  $y \in Y$  if  $y$  is an ordinal number and the unique element  $y'$  satisfying  $\Phi(y, y')$  belongs to  $X'$ . Now by the first property shown in Lemma 8, the set  $Y$  contains every ordinal number. However, this contradicts Burali-Forti Paradox (which states that there exist no sets containing all ordinal numbers). Hence  $X$  should have a maximal element, concluding the proof of Zorn's Lemma.