

# Divergent Syntheses of Indoles and Quinolines Involving N1–C2–C3 Bond Formation through Two Distinct Pd Catalyses

Su San Jang, Young Ho Kim, and So Won Youn\*



Cite This: *Org. Lett.* 2020, 22, 9151–9157



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

**ABSTRACT:** Pd-catalyzed annulative couplings of 2-alkenylanilines with aldehydes using alcohols as both the solvent and hydrogen source have been developed. These domino processes allow divergent syntheses of two significant *N*-heterocycles, indoles and quinolines, from the same substrate by tuning reaction parameters, which seems to invoke two distinct mechanisms. The nature of the ligand and alcoholic solvent had a profound influence on the selectivity and efficiency of these protocols. Particularly noteworthy is that indole formation was achieved by overcoming two significant challenges, regioselective hydropalladation of alkenes and subsequent reactions between the resulting  $\text{Csp}^3$ –Pd species and less reactive imines.




Transition-metal-catalyzed domino annulations enable several transformations in one operation with high selectivity and efficiency, leading to synthesis of a diverse range of carbo- and heterocycles from relatively simple starting materials.<sup>1</sup> In these reactions, the subsequent reactions occur as a consequence of the functionality achieved in the prior step. Thus, switching the reacting sequence may allow access to diverse scaffolds. In recent years, catalytic reactions to transform common substrates or intermediates into distinct molecular scaffolds have increasingly been explored. In this regard, use of different ligands often allows modulation of chemo- and/or regioselectivity, resulting in selective formation of divergent products.<sup>2</sup>

Pd-catalyzed cyclization of nitrogen-containing unsaturated compounds is a straightforward method for constructing *N*-heterocycles such as indoles and quinolines.<sup>1a–i</sup> When aldehydes are added as another reactant in this process, two reaction pathways could occur: (1) aminopalladation followed by nucleophilic addition of the resulting Pd species to aldehydes (Scheme 1a, path a) or (2) initial imine formation followed by annulative coupling (path b).


For path a, Han and Lu reported a cationic Pd-catalyzed domino aminopalladation, nucleophilic addition with aldehydes, and protonation to afford 2-aryl-3-hydroxymethylindoles involving the alkenylpalladium ( $\text{Csp}^2$ –Pd) species (left upper quadrant in the box of Scheme 1a).<sup>3a</sup> Recently, Huang successfully demonstrated such reactions with alkenylanilines by controlling  $\beta$ -hydride elimination of the resulting alkylpalladium ( $\text{Csp}^3$ –Pd) species,<sup>4</sup> which is highly challenging (left lower quadrant).<sup>3b</sup> In this protocol, only electron-donating alkyl groups on the N atom could promote the reaction, while free  $\text{NH}_2$  gave only the corresponding imines resulting from the reaction with aldehydes.

## Scheme 1. Divergent Domino Reactions for Selective Formation of *N*-Heterocycles

(a) Pd-Catalyzed Domino Annulative Coupling Between Alkynyl- or Alkenylanilines and Aldehydes



(b) This Work (via path b): Divergent Synthesis of Indoles and Quinolines via Two Distinct Mechanisms



In contrast, a different reaction order in Pd catalysis (path b) was observed by Yamamoto's group. The initial imine formation triggers hydropalladation of an alkyne moiety followed by annulative addition of the resulting  $\text{Csp}^2$ –Pd species to imine, leading to 3-alkenylindoles by forming the N1–C2–C3 bonds of the pyrrole nucleus of indoles (right

Received: September 1, 2020

Published: October 13, 2020



upper quadrant).<sup>5</sup> Related cationic Pd-catalyzed tandem processes involving initial hydropalladation and sequential intramolecular addition of alkyne-tethered carbonyl compounds using ethanol as a hydrogen source were also reported by Han and Lu.<sup>6</sup> Inspired by these prior studies,<sup>3b,5,6</sup> we envisioned that initially formed imines from 2-alkenylanilines and aldehydes might undergo regioselective hydropalladation of alkenes followed by annulative coupling of the resulting Csp<sup>3</sup>–Pd species with an imine to give indoles (right lower quadrant).

Here, we disclose the realization of this proposal. Pd-catalyzed annulative couplings of 2-alkenylanilines with aldehydes led to divergent syntheses of indoles and quinolines, which are both significant heterocycles due to their ubiquity and broad applications (Scheme 1b).<sup>7,8</sup> In sharp contrast to prior work using 2-alkynylanilines,<sup>9,10</sup> alcohols were used as both the hydrogen source<sup>11</sup> and solvent rather than a nucleophile, in the reactions presented herein.


Considering several reports on the effect of *N*-protecting groups in aminopalladation,<sup>3,12</sup> we first examined various *N*-protecting groups under Huang's conditions<sup>3b</sup> to identify one with a delicately balanced nucleophilicity for aminopalladation and imine/iminium ion formation.<sup>13</sup> Among them, *N*-Boc, *N*-alkyl, and NH<sub>2</sub> free derivatives afforded the desired products, and NH<sub>2</sub> free anilines were the optimal substrate for this reaction, promoting rapid imine formation rather than aminopalladation.

Subsequently, we conducted an extensive investigation of this process using **1a** and PhCHO. Ideally, suitable ligands could stabilize the speculated Csp<sup>3</sup>–Pd species and allow switching of the regioselectivity of the proposed reaction. After careful examination of the reaction parameters, we found that a reagent blend consisting of [Pd(allyl)Cl]<sub>2</sub> and Xantphos in EtOH afforded **3a** along with a smaller amount of **4a**.<sup>13</sup> When the solvent was changed to F-containing alcohol, the ratio of **4a** to **3a** increased (entries 5 and 6). Selectivity for **4a** was markedly improved by switching the ligand from Xantphos to DPEphos in F-containing alcohol solvents (entries 13–16). From the evaluation of bidentate phosphine ligands (Table 1, bottom), we observed preliminary relationships among ligand bite angle, reactivity, and selectivity.<sup>14</sup> With the exception of dppf, bisphosphine ligands with bite angles of ~110° showed good efficiency for indole formation, and the use of Xantphos (108°) provided the best result, while DPEphos had an optimal bite angle (104°) for quinoline formation.

Although Conditions A and B selectively provided **3a** and **4a**, respectively, the selectivity for **3a** over **4a** and the yields of both were only moderate. In addition, although the ratio of **4a** to **3a** was generally good under Conditions B, a non-negligible amount of an indoline side product (e.g., **5a**)<sup>13</sup> was always observed. Therefore, a variety of additives were examined to determine more effective conditions, but no beneficial effect was obtained in any case.

Under Conditions A and B, the reaction of (*Z*)-**1a** also afforded **3a** and **4a**, respectively, in comparable selectivity but lower yields compared to (*E*)-**1a** (entry 7 vs 8, entry 19 vs 20). During the course of these studies, we observed that substituents residing on the aromatic moiety of anilines had a very intriguing effect on the reactivity and selectivity of indole formation. When **1b** (R = Me) was used as a substrate, both yield and ratio of products were significantly improved to afford **3b** as the sole product in 90% yield (entry 9), while this was unsuitable for quinoline formation (entry 21). While the

Table 1. Optimization Studies

						
Entry	<b>1</b>	Pd Catalyst (mol%)	Ligand (mol%)	Solvent	<b>3</b> (%) <sup>a</sup>	<b>4</b> (%) <sup>a</sup>
1	<b>1a</b>	Pd(OAc) <sub>2</sub> (2.5)	Xantphos (5.5)	EtOH	-	-
2	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2.5)	Xantphos (5.5)	EtOH	(38)	(20)
3	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (2.5)	Xantphos (5.5)	EtOH	(44)	(22)
4	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (2.5)	Xantphos (5.5)	<i>i</i> PrOH	10	7
5	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (2.5)	Xantphos (5.5)	TFE	(23)	(31)
6	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (2.5)	Xantphos (5.5)	HFIP	(23)	(32)
7	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (5)	Xantphos (10)	EtOH	(55)	(25)
8 <sup>b</sup>	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (5)	Xantphos (10)	EtOH	(24)	(11)
9 <sup>c</sup>	<b>1b</b>	[Pd(allyl)Cl] <sub>2</sub> (5)	Xantphos (10)	EtOH	(90)	-
10	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (5)	-	EtOH	-	4
11	<b>1a</b>	-	Xantphos (10)	EtOH	-	-
12	<b>1a</b>	-	-	EtOH	-	-
13	<b>1a</b>	Pd(OAc) <sub>2</sub> (2.5)	DPEphos (5.5)	TFE	4	(54)
14	<b>1a</b>	[Pd(allyl)Cl] <sub>2</sub> (2.5)	DPEphos (5.5)	TFE	7	(42)
15	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2.5)	DPEphos (5.5)	TFE	6	(55)
16	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2.5)	DPEphos (5.5)	HFIP	-	29
17	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2.5)	DPEphos (5.5)	EtOH	trace	13
18	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2.5)	DPEphos (5.5)	<i>i</i> PrOH	-	-
19 <sup>d</sup>	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2)	DPEphos (4)	TFE	3	(56)
20 <sup>b,d</sup>	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2)	DPEphos (4)	TFE	-	(31)
21 <sup>d</sup>	<b>1b</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2)	DPEphos (4)	TFE	25	-
22 <sup>d</sup>	<b>1a</b>	PdCl <sub>2</sub> (PhCN) <sub>2</sub> (2)	-	TFE	-	13
23 <sup>d</sup>	<b>1a</b>	-	DPEphos (4)	TFE	-	-
24 <sup>d</sup>	<b>1a</b>	-	-	TFE	-	2

<sup>a</sup> Conditions A : 5 mol% [Pd(allyl)Cl]<sub>2</sub>, 10 mol% ligand in EtOH (0.1 M) at 120 °C for 24 h under Ar  
<sup>b</sup> Conditions B : 2 mol% PdCl<sub>2</sub>(PhCN)<sub>2</sub>, 4 mol% ligand in CF<sub>3</sub>CH<sub>2</sub>OH (0.1 M) at 100 °C for 24 h under Ar

Ligand (bite angle)	dppf (n = 0, 86°)	dppf (n = 1, 91°)	dppf (n = 2, 94°)	DPEphos (104°)	Xantphos (108°)	<i>N</i> -Xantphos (114°)	<i>t</i> Bu-Xantphos (140°)
Conditions A <sup>a</sup>	0–4/(0–11)	(44)/(28)	0/trace	(55)/(25)	21/14	0/0 <sup>f</sup>	0/7 <sup>f</sup>
Conditions B <sup>a</sup>	0/0	0/0	3/(56)	0/24	0/0 <sup>f</sup>	0/0 <sup>f</sup>	0/7 <sup>f</sup>

<sup>a</sup>Determined by <sup>1</sup>H NMR. Values in parentheses indicate isolated yields. <sup>b</sup>Using (*Z*)-**1a** instead of (*E*)-**1a**. <sup>c</sup>For 4 h. <sup>d</sup>At 100 °C. <sup>e</sup>Yields of **3a/4a**. <sup>f</sup>Using 2.5 mol % Pd catalyst and 5.5 mol % ligand at 120 °C.


reason for this substituent effect in indole synthesis remains unclear at this stage, we surmised that the steric effect of *ortho*-Me substituent might render the imine's C atom being in close proximity to a speculated Csp<sup>3</sup>–Pd species (arising from hydropalladation of alkene) to cause a more favorable subsequent reaction (for a detailed explanation, see the Supporting Information).<sup>13</sup>

It should be noted that both cationic Pd(II) species<sup>3a,15</sup> and typical Lewis acids were unable to promote this reaction for formation of **3a**, whereas all tested Lewis acids afforded **4a** in low to moderate yield.<sup>13</sup> These findings suggest that indole formation under Conditions A is unlikely to involve Lewis acid catalyzed electrophilic cyclization via imine activation, which might be involved in the synthesis of quinolines under Conditions B.<sup>16</sup> Control experiments indicated that all the

reaction components (Pd, ligand, alcohol) were necessary for both reactions to occur (entries 10–12 and 22–24).

With viable catalytic systems for each indole and quinoline synthesis, we explored the substrate scope of these processes. First, we examined the reaction of various aldehydes with **1b** under Conditions A (Scheme 2a). In general, a variety of

**Scheme 2. Substrate Scope: Indoles<sup>a</sup>**




<sup>a</sup>Isolated yields. Quinoline byproducts (e.g., **4a**) were obtained in  $\leq 5\%$  yields unless otherwise noted. <sup>b</sup>Quinolines were obtained in 5–15% <sup>1</sup>H NMR yields. <sup>c</sup>Quinolines were obtained in 15–25% <sup>1</sup>H NMR yields. <sup>d</sup>Inseparable mixture of **3f** and dechlorinated product (i.e., **3b**) was obtained with a 10:1 ratio in 74% combined yield. <sup>e</sup>2.1 equiv of **1b** and 1 equiv of terephthalaldehyde were used.

arylaldehydes were well tolerated with little electronic and steric dependence, leading to diversely substituted indoles (**3a–s**) in high yields with good to excellent selectivities. Heteroaryl- (**3t–v**), ester- (**3w**), alkenyl- (**3x**), and secondary alkyl-substituted (**3y**) aldehydes proved to be suitable, while the reaction failed with primary alkylaldehydes. The reaction with terephthalaldehyde proceeded uneventfully to afford bisindole **3z** in 71% yield.

Next, we explored the substituent effect ( $R^1$ ) at the alkene moiety of **1** (Scheme 2b-1). Various (hetero)aryl-substituted alkenes smoothly underwent this reaction to afford **3aa–an** in good to high yields. Unfortunately, alkyl-substituted alkenes resulted in low conversion. The effects of substituents ( $R^2$ ) residing on the aromatic moiety of **1** were also investigated (Scheme 2b-2). As mentioned earlier, a methyl substituent at the *ortho* position had a significant influence on chemical reactivity, forming **3at–au** with excellent yields and selectivities. In contrast, other substituents (e.g., OMe, OH, F) located at the same position had deleterious effects, producing low yields (7–24%) of the desired products.

As shown in Scheme 3, diversely substituted **1** and **2** could also be successfully employed to give the corresponding 2,3-

**Scheme 3. Substrate Scope: Quinolines<sup>a</sup>**




<sup>a</sup>Isolated yields. In most cases, indole (e.g., **3a**) and indoline (e.g., **5a**) byproducts were obtained in 0– $\leq 5\%$  and 10–25% <sup>1</sup>H NMR yields, respectively, unless otherwise noted. <sup>b</sup>Indolines were obtained in 5–15% <sup>1</sup>H NMR yields. <sup>c</sup>Quinolines were obtained in 15–25% <sup>1</sup>H NMR yields. <sup>d</sup>Inseparable mixture of **3f** and dechlorinated product (i.e., **3b**) was obtained with a 10:1 ratio in 74% combined yield. <sup>e</sup>2.1 equiv of **1b** and 1 equiv of terephthalaldehyde were used.

diaryl-substituted quinolines **4**, albeit in moderate yields. We note that various functional groups were commonly well tolerated in both indole- and quinoline-forming reactions except for a nitro group. No matter where the NO<sub>2</sub> group resided in **1** and **2**, no reaction occurred or only imine intermediates were observed.

A series of control experiments were performed to gain mechanistic insight into these reactions.<sup>13</sup> Imine intermediates such as **6a** were observed during the reaction progress. Therefore, probable intermediates (**6a–b**) were synthesized and subjected to Conditions A and B. Each reaction resulted in formation of **3b** and **4a** in 88% and 40% yield, respectively

(Scheme 4a); the latter result is distinct from a related Pd catalysis reported by Jiang.<sup>17</sup>

#### Scheme 4. Mechanistic Studies



Next, deuterium-labeling experiments were carried out. The reaction of 1b or 1a with PhCDO afforded 3b or 4a with no deuterium incorporation (Scheme 4b), excluding the involvement of initial oxidative addition of an imine C–D bond to Pd<sup>17</sup> followed by carbopalladation or hydropalladation to an alkene moiety. When the reaction of 1b with 2a was carried out in  $\text{CH}_3\text{CH}_2\text{OD}$  under Conditions A, indole 3b with deuterium at the benzylic position of the C3 substituent (88% D) was formed, whereas no deuterium incorporation was observed in  $\text{CH}_3\text{CD}_2\text{OH}$  (Scheme 4c, upper).<sup>11e–g</sup> Similarly, quinoline 4a deuterated at the C4 position (42% D) was obtained using  $\text{CF}_3\text{CH}_2\text{OD}$  under Conditions B, while deuterium was not incorporated into 4a in  $\text{CF}_3\text{CD}_2\text{OH}$  (Scheme 4c, lower). These findings indicate that the incoming hydrogen atoms for the benzylic position of the C3 substituent of 3b and for the C4 position of 4a are derived from O–H groups<sup>11c–i</sup> instead of the  $\alpha\text{C–H}$  bond of alcohols.<sup>6,11j–p</sup> Moreover, as mentioned, quinoline synthesis was accompanied by some byproducts, such as *N*-benzylanilines (7) generated by reduction of 6<sup>18</sup> and their cyclized products, indolines (e.g., 5a).<sup>13</sup> Interestingly, 7 deuterated at the benzylic position was also obtained in the reactions using ROD-type solvents. Competition experiments with deuterium-labeled 2a demonstrated a kinetic isotope effect of 3.5 (Scheme 4d), suggesting that C–H bond cleavage of imine<sup>17</sup> may be involved in the rate-determining step.


Reactions employing  $\beta,\beta$ -disubstituted (1al–am) and terminal alkenes (1an) as a substrate were undertaken to demonstrate that Lewis acid catalyzed Prins-type reactions<sup>10,11q,r</sup> are a possible mechanistic pathway in the indole-forming process. The former failed to give the desired indoles,

while the latter produced 3-methylindole 3aw along with 4w, albeit in low yields (Scheme 4e). This constitutes evidence against an electrophilic pathway involving a carbocationic intermediate and ethanol as a proton source in the indole-forming reaction under Conditions A.

Although more detailed investigations are needed to clarify the mechanism, our experimental findings suggest that these two reactions proceed through different mechanistic pathways.<sup>13</sup> For quinoline (4) synthesis, the reaction seems to take place through  $6\pi$ -electrocyclization of imine intermediates (6) promoted by Lewis acidic Pd salt.<sup>16</sup> In sharp contrast, indole (3) formation presumably commences with regioselective insertion of the alkene moiety of 6 into the Pd–H bond.<sup>19</sup> Then, subsequent reactions between the resulting  $\text{Csp}^3$ –Pd species and imine moiety may occur through either oxidative addition/reductive elimination<sup>17</sup> or carbopalladation/ $\beta$ -hydride elimination,<sup>5,6</sup> followed by isomerization to afford 3.

To highlight the synthetic utility of this transformation, dearomatization of indoles, a powerful tool for construction of spiroindolenine and polycyclic indoline skeletons,<sup>20</sup> was undertaken (Scheme 5).  $\text{I}_2$ -mediated cyclization<sup>21</sup> and

#### Scheme 5. Synthetic Applications



fluoroetherification<sup>22</sup> of 3ag proceeded smoothly to afford spiroindolenine 8a and fused indoline 8b, respectively. Subsequent acetylation or benzylation gave indolenines 8c–d containing a fluorinated quaternary carbon center through ring opening. On the other hand, 3-(indolylmethyl)carbazole derivatives are known as highly efficient hosts for phosphorescent OLEDs<sup>23a</sup> and as colorimetric/fluorometric detectors of DNA.<sup>23b</sup> Reaction of carbazole-tethered 1ao with PhCHO was readily accomplished to afford 3ax in 88% yield.

In summary, we developed Pd-catalyzed domino annulative processes, one of which involves the significant challenges associated with controlling both regioselective hydropalladation and reaction of alkylpalladium species with less reactive imines. We demonstrated that exquisite tuning of reaction parameters such as ligand and solvent could modulate the reaction propensity and pathway for formation of different molecular frameworks from the same substrate. In these protocols, alcohols were used as the hydrogen source as well as solvent, delivering hydrogen atoms from O–H groups rather than the  $\alpha\text{C–H}$  of alcohols. To the best of our knowledge, this represents the first example of Pd-catalyzed domino couplings between alkenes and imines generated in situ from amines and



aldehydes, leading to construction of two different N-heterocycles.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c02898>.

Full experimental details and characterization data (PDF)

## Accession Codes

CCDC 2012360 and 2012365–2012366 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## ■ AUTHOR INFORMATION

### Corresponding Author

**So Won Youn** — Center for New Directions in Organic Synthesis, Department of Chemistry and Research Institute for Natural Sciences, Hanyang University, Seoul 04763, Korea;  
✉ [orcid.org/0000-0002-2134-6487](mailto:orcid.org/0000-0002-2134-6487); Email: [sowony73@hanyang.ac.kr](mailto:sowony73@hanyang.ac.kr)

### Authors

**Su San Jang** — Center for New Directions in Organic Synthesis, Department of Chemistry and Research Institute for Natural Sciences, Hanyang University, Seoul 04763, Korea  
**Young Ho Kim** — Center for New Directions in Organic Synthesis, Department of Chemistry and Research Institute for Natural Sciences, Hanyang University, Seoul 04763, Korea

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.0c02898>

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by both the Basic Science Research Program and Nano-Material Technology Department Program through the National Research Foundation of Korea (NRF) funded by the Korea government (MSIT) (Nos. 2012M3A7B4049644, 2018R1A2A2A05018392, and 2014-011165). We thank Dr. Ji-Eun Lee (Central Instrument Facility, Gyeongsang National University) for X-ray crystallographic analysis.

## ■ REFERENCES

(1) Selected reviews: (a) Christodoulou, M. S.; Beccalli, E. M.; Foschi, F.; Giofré, S. Pd-Catalyzed Domino Reactions Involving Alkenes To Access Substituted Indole Derivatives. *Synthesis* **2020**, 52, 2731. (b) Han, X.; Lu, X. Palladium(II)-Catalyzed Redox-Neutral Cyclizations of Alkynes Containing Alkenyl or Electrophilic Functional Groups: A Convenient Synthesis of Carbocycles and Heterocycles. *Synlett* **2018**, 29, 2461. (c) Platon, M.; Amardeil, R.; Djakovitch, L.; Hierso, J.-C. Progress in palladium-based catalytic systems for the sustainable synthesis of annulated heterocycles: a focus on indole backbones. *Chem. Soc. Rev.* **2012**, 41, 3929. (d) Cacchi, S.; Fabrizi, G. Update 1 of: Synthesis and Functionalization of Indoles Through Palladium-Catalyzed Reactions. *Chem. Rev.*

**2011**, 111, PR215. (e) Beccalli, E. M.; Broggini, G.; Martinelli, M.; Sottocornola, S. C–C, C–O, C–N Bond Formation on  $sp^2$  Carbon by Pd(II)-Catalyzed Reactions Involving Oxidant Agents. *Chem. Rev.* **2007**, 107, 5318. (f) Zeni, G.; Larock, R. C. Synthesis of Heterocycles via Palladium  $\pi$ -Olefin and  $\pi$ -Alkyne Chemistry. *Chem. Rev.* **2004**, 104, 2285. (g) Grigg, R.; Sridharan, V. Palladium catalysed cascade cyclisation-anion capture, relay switches and molecular queues. *J. Organomet. Chem.* **1999**, 576, 65. (h) Negishi, E.; Copéret, C.; Ma, S.; Liou, S. Y.; Liu, F. Cyclic Carbopalladation. A Versatile Synthetic Methodology for the Construction of Cyclic Organic Compounds. *Chem. Rev.* **1996**, 96, 365. (i) Barluenga, J.; Rodríguez, F.; Fañanás, F. J. Recent Advances in the Synthesis of Indole and Quinoline Derivatives through Cascade Reactions. *Chem. - Asian J.* **2009**, 4, 1036. (j) Youn, S. W. Rhodium-Catalyzed Tandem Transformations with Organoboron Reagents: Sequential Multiple C–C Bond Formations. *Eur. J. Org. Chem.* **2009**, 2009, 2597. (k) Montgomery, J. Nickel-Catalyzed Reductive Cyclizations and Couplings. *Angew. Chem., Int. Ed.* **2004**, 43, 3890.

(2) Review: (a) Nájera, C.; Beletskaya, I. P.; Yus, M. Metal-catalyzed regiodivergent organic reactions. *Chem. Soc. Rev.* **2019**, 48, 4515. (b) Lee, Y.-C.; Kumar, K.; Waldmann, H. Ligand-Directed Divergent Synthesis of Carbo- and Heterocyclic Ring Systems. *Angew. Chem., Int. Ed.* **2018**, 57, 5212. (c) Mahatthananchai, J.; Dumas, A. M.; Bode, J. W. Catalytic Selective Synthesis. *Angew. Chem., Int. Ed.* **2012**, 51, 10954. For selected recent examples, see: (d) Meng, Q.-Y.; Schirmer, T. E.; Katou, K.; König, B. Controllable Isomerization of Alkenes by Dual Visible-Light-Cobalt Catalysis. *Angew. Chem., Int. Ed.* **2019**, 58, 5723. (e) Youn, S. W.; Ko, T. Y.; Jang, Y. H. Palladium-Catalyzed Regioselective Synthesis of 3-Arylindoles from N-Ts-Anilines and Styrenes. *Angew. Chem., Int. Ed.* **2017**, 56, 6636. (f) Hyster, T. K.; Dalton, D. M.; Rovis, T. Ligand design for Rh(III)-catalyzed C–H activation: an unsymmetrical cyclopentadienyl group enables a regioselective synthesis of dihydroisoquinolones. *Chem. Sci.* **2015**, 6, 254. (g) Fujino, D.; Yorimitsu, H.; Osuka, A. Regiocontrolled Palladium-Catalyzed Arylative Cyclizations of Alkynols. *J. Am. Chem. Soc.* **2014**, 136, 6255. (h) Tang, S.-Y.; Guo, Q.-X.; Fu, Y. Mechanistic Origin of Ligand-Controlled Regioselectivity in Pd-Catalyzed C–H Activation/Arylation of Thiophenes. *Chem. - Eur. J.* **2011**, 17, 13866. (3) (a) Han, X.; Lu, X. Pd(II)-Catalyzed Tandem Reaction of 2-Arylethynylanilines and Aldehydes: An Efficient Synthesis of Substituted 3-Hydroxymethyl Indoles. *Org. Lett.* **2010**, 12, 3336. (b) Hu, Y.; Xie, Y.; Shen, Z.; Huang, H. Palladium-Catalyzed Ring-Forming Aminoalkenylation of Alkenes with Aldehydes Initiated by Intramolecular Aminopalladation. *Angew. Chem., Int. Ed.* **2017**, 56, 2473.

(4) For reviews on the preference of  $\beta$ -hydride elimination of alkylpalladium species, see: (a) *Metal-Catalyzed Cross-Coupling Reactions*; de Meijere, A.; Diederich, F., Eds.; Wiley-VCH: New York, 2004. (b) *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. I; Negishi, E.-i., Ed.; Wiley-Interscience: New York, 2002. (c) Cárdenas, D. J. Towards Efficient and Wide-Scope Metal-Catalyzed Alkyl–Alkyl Cross-Coupling Reactions. *Angew. Chem., Int. Ed.* **1999**, 38, 3018. (d) Luh, T.-Y.; Leung, M.-K.; Wong, K.-T. Transition Metal-Catalyzed Activation of Aliphatic C–X Bonds in Carbon–Carbon Bond Formation. *Chem. Rev.* **2000**, 100, 3187. (e) Cárdenas, D. J. Advances in Functional-Group-Tolerant Metal-Catalyzed Alkyl–Alkyl Cross-Coupling Reactions. *Angew. Chem., Int. Ed.* **2003**, 42, 384.

(5) Takeda, A.; Kamijo, S.; Yamamoto, Y. Indole Synthesis via Palladium-Catalyzed Intramolecular Cyclization of Alkynes and Imines. *J. Am. Chem. Soc.* **2000**, 122, 5662.

(6) (a) Shen, K.; Han, X.; Lu, X. Cationic Pd(II)-Catalyzed Reductive Cyclization of Alkyne-Tethered Ketones or Aldehydes Using Ethanol as Hydrogen Source. *Org. Lett.* **2013**, 15, 1732. (b) Shen, K.; Han, X.; Xia, G.; Lu, X. Cationic Pd(II)-catalyzed cyclization of N-tosyl-aniline tethered alkynyl ketones initiated by hydropalladation of alkynes: a facile way to 1,2-dihydro or 1,2,3,4-tetrahydroquinoline derivatives. *Org. Chem. Front.* **2015**, 2, 145. (c) Zhang, X.; Han, X.; Hu, Z.; Lu, X. Synthesis of Substituted

Piperidines via Cationic Palladium(II)-Catalyzed Reductive Coupling of *N*-Tosyl-Tethered Alkynes. *Synthesis* **2017**, 49, 4687.

(7) Indoles: (a) Shiri, M. Indoles in Multicomponent Processes (MCPs). *Chem. Rev.* **2012**, 112, 3508. (b) Inman, M.; Moody, C. J. Indole synthesis – something old, something new. *Chem. Sci.* **2013**, 4, 29. (c) Youn, S. W.; Ko, T. Y. Metal-Catalyzed Synthesis of Substituted Indoles. *Asian J. Org. Chem.* **2018**, 7, 1467 and refs 1c, d, and 1i. Quinolines: (d) Kouznetsov, V. V.; Méndez, L. Y. V.; Gómez, C. M. M. Recent Progress in the Synthesis of Quinolines. *Curr. Org. Chem.* **2005**, 9, 141. (e) Madapa, S.; Tusi, Z.; Batra, S. Advances in the Syntheses of Quinoline and Quinoline-Annulated Ring Systems. *Curr. Org. Chem.* **2008**, 12, 1116. (f) Marco-Contelles, J.; Pérez-Mayoral, E.; Samadi, A.; Carreiras, M. C.; Soriano, E. Recent Advances in the Friedländer Reaction. *Chem. Rev.* **2009**, 109, 2652. (g) Khushnutdinov, R. I.; Bayguzina, A. R.; Dzhemilev, U. M. Metal complex catalysis in the synthesis of quinolines. *J. Organomet. Chem.* **2014**, 768, 75. (h) Bharate, J. B.; Vishwakarma, R. A.; Bharate, S. B. Metal-free domino one-pot protocols for quinoline synthesis. *RSC Adv.* **2015**, 5, 42020. (i) Naidoo, S.; Jeena, V. Synthesis of 2,4-Disubstituted Quinoline Derivatives via A<sup>3</sup>-Coupling: An EcoScale Evaluation. *Synthesis* **2017**, 49, 2621 and ref 1i.

(8) For examples on the divergent construction of either an indole or a quinoline scaffold from the same precursors, 2-alkenyl- or 2-alkynylanilines, see: (a) Parker, K. A.; Mindt, T. L. Heterocycle Annulation of Enolizable Vinyl Quinone Imides. Dihydroquinolines and Quinolines from Thermal 6 $\pi$ -Electrocyclizations and Indoles from Photochemical Cyclizations. *Org. Lett.* **2002**, 4, 4265. (b) Bowman, W. R.; Fletcher, A. J.; Pedersen, J. M.; Lovell, P. J.; Elsegood, M. R. J.; López, E. H.; McKee, V.; Potts, G. B. S. Amides as precursors of imidoyl radicals in cyclisation reactions. *Tetrahedron* **2007**, 63, 191. (c) Halim, R.; Scammells, P. J.; Flynn, B. L. Alternating Iodonium-Mediated Reaction Cascades Giving Indole- And Quinoline-Containing Polycycles. *Org. Lett.* **2008**, 10, 1967. (d) Kwon, S. H.; Seo, H.-A.; Cheon, C.-H. Total Synthesis of Luotonin A and Rutaecarpine from an Aldimine via the Designed Cyclization. *Org. Lett.* **2016**, 18, 5280.

(9) (a) Kamijo, S.; Sasaki, Y.; Yamamoto, Y. Copper-catalyzed tandem reaction between imines and alcohols leading to indoles. *Tetrahedron Lett.* **2004**, 45, 35. (b) Fu, W.; Xu, C.; Zou, G.; Hong, D.; Deng, D.; Wang, Z.; Ji, B. AuCl<sub>3</sub>-Catalyzed Tandem Reaction of *N*-(*o*-Alkynylphenyl)imines: A Modular Entry to Polycyclic Frameworks Containing an Indole Unit. *Synlett* **2009**, 2009, 763. (c) Yuan, S.; Zhang, D.-Q.; Zhang, J.-Y.; Yu, B.; Liu, H.-M. Palladium-Catalyzed Ligand-Free Double Cyclization Reactions for the Synthesis of 3-(1'-Indolyl)-phthalides. *Org. Lett.* **2020**, 22, 814 and ref 8c.

(10) (a) Wang, Y.; Peng, C.; Liu, L.; Zhao, J.; Su, L.; Zhu, Q. Sulfuric acid promoted condensation cyclization of 2-(2-(trimethylsilyl)ethynyl)anilines with arylaldehydes in alcoholic solvents: an efficient one-pot synthesis of 4-alkoxy-2-arylquinolines. *Tetrahedron Lett.* **2009**, 50, 2261. (b) Ock, S. K.; Youn, S. W. Synergistic Effect of Pd(II) and Acid Catalysts on Tandem Annulation Reaction for the Regioselective Synthesis of Ring-Fused Quinolines. *Bull. Korean Chem. Soc.* **2010**, 31, 704 and ref 8c.

(11) Reviews: (a) Grushin, V. V. Hydrido Complexes of Palladium. *Chem. Rev.* **1996**, 96, 2011. (b) Muzart, J. Pd-Catalyzed Hydrogen-Transfer Reactions from Alcohols to C = C, C = O, and C = N Bonds. *Eur. J. Org. Chem.* **2015**, 2015, 5693. O–H of alcohol as a H-atom source of M–H: (c) Marcum, J. S.; Taylor, T. R.; Meek, S. J. Enantioselective Synthesis of Functionalized Arenes by Nickel-Catalyzed Site-Selective Hydroarylation of 1,3-Dienes with Aryl Boronates. *Angew. Chem., Int. Ed.* **2020**, 59, 14070. (d) Ji, D.-W.; Hu, Y.-C.; Zheng, H.; Zhao, C.-Y.; Chen, Q.-A.; Dong, V. M. A regioselectivity hydroallylation of switch in Pd-catalyzed alkynes. *Chem. Sci.* **2019**, 10, 6311. (e) Chen, Y.-G.; Shuai, B.; Xu, X.-T.; Li, Y.-Q.; Yang, Q.-L.; Qiu, H.; Zhang, K.; Fang, P.; Mei, T.-S. Nickel-catalyzed Enantioselective Hydroarylation and Hydroalkenylation of Styrenes. *J. Am. Chem. Soc.* **2019**, 141, 3395. (f) Xiao, L.-J.; Cheng, L.; Feng, W.-M.; Li, M.-L.; Xie, J.-H.; Zhou, Q.-L. Nickel(0)-Catalyzed Hydroarylation of Styrenes and 1,3-Dienes with Organoboron

Compounds. *Angew. Chem., Int. Ed.* **2018**, 57, 461. (g) Wang, X.; Nakajima, M.; Martin, R. Ni-Catalyzed Regioselective Hydrocarboxylation of Alkynes with CO<sub>2</sub> by Using Simple Alcohols as Proton Sources. *J. Am. Chem. Soc.* **2015**, 137, 8924. (h) Clegg, W.; Eastham, G. R.; Elsegood, M. R. J.; Heaton, B. T.; Iggo, J. A.; Tooze, R. P.; Whyman, R.; Zacchini, S. Characterization and Dynamics of [Pd(L-L)H(solvent)]<sup>+</sup>, [Pd(L-L)(CH<sub>2</sub>CH<sub>3</sub>)]<sup>+</sup>, and [Pd(L-L)(C(O)Et)-(THF)]<sup>+</sup> (L-L = 1,2-(CH<sub>2</sub>PBu<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>): Key Intermediates in the Catalytic Methoxycarbonylation of Ethene to Methylpropanoate. *Organometallics* **2002**, 21, 1832. (i) Eastham, G. R.; Heaton, B. T.; Iggo, J. A.; Tooze, R. P.; Whyman, R.; Zacchini, S. Synthesis and spectroscopic characterisation of all the intermediates in the Pd-catalysed methoxycarbonylation of ethene. *Chem. Commun.* **2000**, 609.  $\alpha$ C–H of alcohol as a H-atom source of M–H: (j) Chen, J.; Han, X.; Lu, X. Palladium(II)-Catalyzed Reductive Cyclization of *N*-Tosyl-Tethered 1,7-Enynes: Enantioselective Synthesis of 1,2,3,4-Tetrahydroquinolines. *Org. Lett.* **2019**, 21, 8153. (k) Wu, W.; Chen, T.; Chen, J.; Han, X. Cationic Palladium(II)-Catalyzed Reductive Cyclization of Alkynyl Cyclohexadienones. *J. Org. Chem.* **2018**, 83, 1033. (l) Liao, L.; Sigman, M. S. Palladium-Catalyzed Hydroarylation of 1,3-Dienes with Boronic Esters via Reductive Formation of  $\pi$ -Allyl Palladium Intermediates under Oxidative Conditions. *J. Am. Chem. Soc.* **2010**, 132, 10209. (m) Gligorich, K. M.; Cummings, S. A.; Sigman, M. S. Palladium-Catalyzed Reductive Coupling of Styrenes and Organostannanes under Aerobic Conditions. *J. Am. Chem. Soc.* **2007**, 129, 14193. (n) Monguchi, D.; Beemelmans, C.; Hashizume, D.; Hamashima, Y.; Sodeoka, M. Catalytic asymmetric conjugate reduction with ethanol: A more reactive system Pd(II)–<sup>t</sup>Pr-DUPHOS complex with molecular sieves 4A. *J. Organomet. Chem.* **2008**, 693, 867. (o) Tsuchiya, Y.; Hamashima, Y.; Sodeoka, M. A New Entry to Pd–H Chemistry: Catalytic Asymmetric Conjugate Reduction of Enones with EtOH and a Highly Enantioselective Synthesis of Warfarin. *Org. Lett.* **2006**, 8, 4851. (p) Clegg, W.; Eastham, G. R.; Elsegood, M. R. J.; Heaton, B. T.; Iggo, J. A.; Tooze, R. P.; Whyman, R.; Zacchini, S. Synthesis and reactivity of palladium hydrido-solvent complexes, including a key intermediate in the catalytic methoxycarbonylation of ethene to methyl propanoate. *J. Chem. Soc., Dalton Trans.* **2002**, 3300 and ref 6. Metal-catalyzed direct hydride transfer from  $\alpha$ C–H of alcohol without involving M–H species: (q) Zheng, Y. L.; Liu, Y. Y.; Wu, Y. M.; Wang, Y. X.; Lin, Y. T.; Ye, M. C. Iron-Catalyzed Regioselective Transfer Hydrogenative Couplings of Unactivated Aldehydes with Simple Alkenes. *Angew. Chem., Int. Ed.* **2016**, 55, 6315. (r) Li, J.; Preinfalk, A.; Maulide, N. Enantioselective Redox-Neutral Coupling of Aldehydes and Alkenes by an Iron-Catalyzed “Catch–Release” Tethering Approach. *J. Am. Chem. Soc.* **2019**, 141, 143.

(12) For reviews on Pd(II)-catalyzed C–H amination via intramolecular aminopalladation of alkenes for the synthesis of indoles and related heterocycles, see: (a) McDonald, R. I.; Liu, G.; Stahl, S. S. Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, 111, 2981. (b) Minatti, A.; Muñiz, K. Intramolecular aminopalladation of alkenes as a key step to pyrrolidines and related heterocycles. *Chem. Soc. Rev.* **2007**, 36, 1142. (c) Müller, T. E.; Beller, M. Metal-Initiated Amination of Alkenes and Alkynes. *Chem. Rev.* **1998**, 98, 675. (d) Hege, L. S. Transition Metals in the Synthesis and Functionalization of Indoles. *Angew. Chem., Int. Ed. Engl.* **1988**, 27, 1113. For selected recent examples, see: (e) Muñiz, K. Advancing Palladium-Catalyzed C–N Bond Formation: Bisindoline Construction from Successive Amide Transfer to Internal Alkenes. *J. Am. Chem. Soc.* **2007**, 129, 14542. (f) Tselikhovskiy, D.; Buchwald, S. L. Synthesis of Heterocycles via Pd-Ligand Controlled Cyclization of 2-Chloro-*N*-(2-vinyl)aniline: Preparation of Carbazoles, Indoles, Dibenzazepines, and Acridines. *J. Am. Chem. Soc.* **2010**, 132, 14048. (g) Youn, S. W.; Bihn, J. H.; Kim, B. S. Pd-Catalyzed Intramolecular Oxidative C–H Amination: Synthesis of Carbazoles. *Org. Lett.* **2011**, 13, 3738. (h) Youn, S. W.; Lee, S. R. Unusual 1,2-Aryl Migration in Pd(II)-Catalyzed Aza-

Wacker-Type Cyclization of 2-Alkenylanilines. *Org. Biomol. Chem.* **2015**, *13*, 4652.

(13) For details, see the [Supporting Information](#).

(14) Reviews: (a) Dierkes, P.; van Leeuwen, P. W. N. M. The bite angle makes the difference: a practical ligand parameter for diphosphine ligands. *J. Chem. Soc., Dalton Trans.* **1999**, 1519. (b) van Leeuwen, P. W. N. M.; Kamer, P. C. J.; Reek, J. N. H.; Dierkes, P. Ligand Bite Angle Effects in Metal-catalyzed C–C Bond Formation. *Chem. Rev.* **2000**, *100*, 2741. (c) Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Reek, J. N. H. Wide Bite Angle Diphosphines: Xantphos Ligands in Transition Metal Complexes and Catalysis. *Acc. Chem. Res.* **2001**, *34*, 895. (d) Freixa, Z.; van Leeuwen, P. W. N. M. Bite angle effects in diphosphine metal catalysts: steric or electronic? *Dalton Trans* **2003**, 1890. (e) Birkholz, M.-N.; Freixa, Z.; van Leeuwen, P. W. N. M. Bite angle effects of diphosphines in C–C and C–X bond forming cross coupling reactions. *Chem. Soc. Rev.* **2009**, *38*, 1099. (f) Mansell, S. M. Catalytic applications of small bite-angle diphosphorus ligands with single-atom linkers. *Dalton Trans* **2017**, 46, 15157. Selected examples: (g) Daniels, D. S. B.; Jones, A. S.; Thompson, A. L.; Paton, R. S.; Anderson, E. A. Ligand Bite Angle-Dependent Palladium-Catalyzed Cyclization of Propargylic Carbonates to 2-Alkynyl Azacycles or Cyclic Dienamides. *Angew. Chem., Int. Ed.* **2014**, *53*, 1915. (h) Niemeyer, Z. L.; Milo, A.; Hickey, D. P.; Sigman, M. S. Parameterization of phosphine ligands reveals mechanistic pathways and predicts reaction outcomes. *Nat. Chem.* **2016**, *8*, 610. (i) Fang, X.; Li, Q.; Shi, R.; Yao, H.; Lin, A. Controllable Pd-Catalyzed Allylation of Indoles with Skipped Enynes: Divergent Synthesis of Indolenines and *N*-Allylindoles. *Org. Lett.* **2018**, *20*, 6084. (j) Chuentragool, P.; Yadagiri, D.; Morita, T.; Sarkar, S.; Parasram, M.; Wang, Y.; Gevorgyan, V. Aliphatic Radical Relay Heck Reaction at Unactivated C(sp<sup>3</sup>)-H Sites of Alcohols. *Angew. Chem., Int. Ed.* **2019**, *58*, 1794 and ref [11d](#).

(15) Mikami, K.; Hatano, M.; Akiyama, K. Active Pd(II) Complexes as Either Lewis Acid Catalysts or Transition Metal Catalysts. *Top. Organomet. Chem.* **2005**, *14*, 279 and references cited therein.

(16) (a) Qiang, L. G.; Baine, N. H. A Convenient Synthesis of Substituted Quinolines by Thermal Electrocyclic Rearrangement of *o*-Vinyl Anils under Nonacidic Conditions. *J. Org. Chem.* **1988**, *53*, 4218. (b) Zhang, X.; Xu, X.; Yu, L.; Zhao, Q. Bronsted acid-mediated reactions of aldehydes with 2-vinylaniline and biphenyl-2-amine. *Tetrahedron Lett.* **2014**, *55*, 2280.

(17) Ji, X.; Huang, H.; Li, Y.; Chen, H.; Jiang, H. Palladium-Catalyzed Sequential Formation of C–C Bonds: Efficient Assembly of 2-Substituted and 2,3-Disubstituted Quinolines. *Angew. Chem., Int. Ed.* **2012**, *51*, 7292.

(18) Examples on the oxidation of dihydroquinolines by imines via hydrogen transfer to give quinolines accompanied by the formation of amines as a side product: (a) Saito, K.; Shibata, Y.; Yamanaka, M.; Akiyama, T. Chiral Phosphoric Acid-Catalyzed Oxidative Kinetic Resolution of Indolines Based on Transfer Hydrogenation to Imines. *J. Am. Chem. Soc.* **2013**, *135*, 11740. (b) Saito, K.; Miyashita, H.; Akiyama, T. Chiral phosphoric acid catalyzed oxidative kinetic resolution of cyclic secondary amine derivatives including tetrahydroquinolines by hydrogen transfer to imines. *Chem. Commun.* **2015**, *51*, 16648 and refs [16a](#) and [17](#).

(19) Ascic, E.; Buchwald, S. L. Highly Diastereo- and Enantioselective CuH-Catalyzed Synthesis of 2,3-Disubstituted Indolines. *J. Am. Chem. Soc.* **2015**, *137*, 4666.

(20) For selected reviews, see: (a) Huang, G.; Yin, B. Recent Developments in Transition Metal-Catalyzed Dearomative Cyclizations of Indoles as Dipolarophiles for the Construction of Indolines. *Adv. Synth. Catal.* **2019**, *361*, 405. (b) Zheng, C.; You, S.-L. Catalytic asymmetric dearomatization (CADA) reaction-enabled total synthesis of indole-based natural products. *Nat. Prod. Rep.* **2019**, *36*, 1589. (c) Saya, J. M.; Ruijter, E.; Orru, R. V. A. Total Synthesis of *Aspidosperma* and *Strychnos* Alkaloids through Indole Dearomatization. *Chem. - Eur. J.* **2019**, *25*, 8916. (d) Bariwal, J.; Voskressensky, L. G.; Van der Eycken, E. V. Recent advances in spirocyclization of indole derivatives. *Chem. Soc. Rev.* **2018**, *47*, 3831. (e) Roche, S. P.;

Tendoung, J.-J. Y.; Tréguier, B. Advances in dearomatization strategies of indoles. *Tetrahedron* **2015**, *71*, 3549. (f) Zhuo, C.-X.; Zhang, W.; You, S.-L. Catalytic Asymmetric Dearomatization Reactions. *Angew. Chem., Int. Ed.* **2012**, *51*, 12662. (g) Roche, S. P.; Porco, J. A., Jr. Dearomatization Strategies in the Synthesis of Complex Natural Products. *Angew. Chem., Int. Ed.* **2011**, *50*, 4068.

(21) Miao, C.-B.; Sun, Y.-F.; Wu, H.; Sun, X.-Q.; Yang, H.-T. CuI-Catalyzed Selective 3-Alkylation of Indoles with *N*-Tosylhydrazones and the I<sub>2</sub>-Mediated Further Cyclization to Chromeno[2,3-*b*]Indoles. *Adv. Synth. Catal.* **2018**, *360*, 2440.

(22) (a) Lozano, O.; Blessley, G.; del Campo, T. M.; Thompson, A. L.; Giuffredi, G. T.; Bettati, M.; Walker, M.; Borman, R.; Gouverneur, V. Organocatalyzed Enantioselective Fluorocyclizations. *Angew. Chem., Int. Ed.* **2011**, *50*, 8105. (b) Zhao, W.; Wang, Z.; Chu, B.; Sun, J. Enantioselective Formation of All-Carbon Quaternary Stereocenters from Indoles and Tertiary Alcohols Bearing A Directing Group. *Angew. Chem., Int. Ed.* **2015**, *54*, 1910. (c) Wang, H.; Wang, Y.; Liu, S.; Mai, Y.; Zong, X.; Gao, H.; Hu, R.; Jiang, X.; Yeung, Y.-Y. Enantioselective Fluorocyclizations Mediated by Amino-Acid-Derived Phthalazine. *Adv. Synth. Catal.* **2019**, *361*, 5334.

(23) (a) Keruckiene, R.; Volyniuk, D.; Ostrauskaite, J.; Peculyte, L.; Grazulevicius, J. V.; Kostjuk, S. V.; Lazauskas, A. Derivatives of 2-phenylindole and carbazole as host materials for phosphorescent organic light emitting diodes. *Dyes Pigm.* **2017**, *137*, 58. (b) Meesala, R.; Nagarajan, R. Synthesis of new heteroaryldi(diindolyl)methanes: Colorimetric detection of DNA by di(diindolylmethyl)carbazoles. *J. Chem. Sci.* **2009**, *121*, 183.