


Lewis Acid Promoted Benzylic Cross-Couplings of Pyridines with Aryl Bromides**

Stéphanie Duez, Andreas K. Steib, Sophia M. Manolikakes, and Paul Knochel*

The functionalization of pyridines and related heterocycles is very important because of their biological properties and relevance to material science.^[1] The benzylic arylation of pyridines, in particular, is a challenging synthetic problem. Palladium-catalyzed arylations of 2-picoline involving direct C–H activation^[2] have no generality, and only few examples have been reported. Thus, azaarenes bearing electron-withdrawing groups may be arylated at 100 °C with a Pd catalyst.^[3] Several alternative procedures involving the fragmentation of a 2-(2-pyridyl)ethanol,^[4] the arylation of *N*-oxides,^[5] and *N*-iminopyridinium ylides^[6] have been described. These methods, although displaying generality, require modified *N*-heterocyclic precursors.^[4–6] In addition, whereas 2-picoline (**1a**) can be functionalized in this way, the arylation of 4-picoline (**2a**) has not been described. The difficulty in forming a new carbon–carbon bond with metalated 2-picoline (**3**; or 4-picoline) may be due to the nature of the palladium complexes^[7] (**4a–c**) resulting from the reaction with ArPdX (Scheme 1). We anticipate that all of these possible structures are reluctant to undergo a reductive elimination because of the chelation of the heterocyclic nitrogen with the Pd center. Hartwig and co-workers have already shown that palladium-catalyzed aminations are accelerated by a Lewis acid (BEt₃).^[8] Nolan and co-workers have also reported that reductive eliminations of Pd complexes are accelerated by AlCl₃.^[9]


We envisioned that the presence of an appropriate Lewis acid (LA) complexing the nitrogen atom of the heterocycle may lead to a new Pd intermediate such as **5**, which would then undergo fast reductive elimination leading to the cross-coupling product **6**. Similar behavior may be expected for the arylation of 4-picoline (Scheme 1). The beneficial effect of Lewis acids in the additions of 2-picoline to imines and enones has already been demonstrated.^[10,11] Thus, we directed our attention toward the use of bases (Met-base) bearing a Lewis acidic metal center for the metalation. Recently, we have reported a kinetically highly active LiCl-solubilized TMP base (TMP = 2,2,6,6-tetramethylpiperidyl): TMPZnCl·LiCl (**7**) displays high chemoselectivity in various directed zinca-



Scheme 1. Lewis acid (LA) promoted benzylic cross-coupling.

tions of arenes and heterocycles.^[12,13] Besides, **7** proved to be an excellent base for the generation of nitrile and ester enolates.^[13,14] We have also demonstrated that **7** is compatible with additional strong Lewis acids (MgCl₂, BF₃·OEt₂) and forms frustrated Lewis pairs.^[15] Herein, we report that Lewis acids such as ZnCl₂, MgCl₂, BF₃·OEt₂, and Sc(OTf)₃ in combination with TMPZnCl·LiCl promote efficiently the Negishi cross-coupling^[16] of various methyl-substituted *N*-heterocycles.

Thus, the zincation of 2-picoline (**1a**) with **7** (2.0 equiv)^[17] gives the zincated picoline **8a**. Its cross-coupling with 5-bromoindole (**9a**, 0.8 equiv) in the presence of 2 mol% Pd(OAc)₂ and 2–4 mol% SPhos^[18] affords the desired pyridine **6a** in 86% yield (in Scheme 2). Such cross-coupling reactions can be extended to various substituted aryl bromides (**9b–d**) leading to products **6b–d** in 66 to 95% yield (Table 1, entries 1–3). Also, 2-substituted pyridines such as **1b,c** are metalated with **7** under the same conditions and react

Scheme 2. Palladium-catalyzed direct cross-coupling of picolines **1a** and **2a**.

[*] Dr. S. Duez, M. Sc. A. K. Steib, M. Sc. S. M. Manolikakes, Prof. Dr. P. Knochel
Department Chemie, Ludwig Maximilians-Universität München
Butenandtstrasse 5–13, Haus F, 81377 München (Germany)
E-mail: paul.knochel@cup.uni-muenchen.de

[**] We thank the Fonds der Chemischen Industrie, the European Research Council (ERC), and the Deutsche Forschungsgemeinschaft (DFG) for financial support. We also thank BASF AG (Ludwigshafen), Chemetall GmbH (Frankfurt), and Heraeus (Hanau) for generous gifts of chemicals.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201103074>.

Table 1: Direct benzylic cross-coupling of 2- and 4-picoline derivatives.


Entry	Picoline ^[a]	Aryl-Br	Product	Yield [%] ^[b]
		Br-		
1	1a (3)	9b : R = 4-OMe	6b : R = 4-OMe	95
2	1a (6)	9c : R = 4-F	6c : R = 4-F	78
3	1a (6)	9d : R = 3-Cl	6d : R = 3-Cl	66
		9b	6e	99
4	1b (20)	9b	6e	99
		9b	6f	92
5 ^[e]	1c (11)	9b	6f	92
		9b	10b : R = 4-OMe	98 ^[c]
6	2a (1)	9b	10b : R = 4-OMe	98 ^[c]
7	2a (1)	9f : R = 3-Me	10c : R = 3-Me	82
8	2a (1)	9g : R = 4-NMe ₂	10d : R = 4-NMe ₂	70
9	2a (1)	9h : R = 4-OH	10e : R = 4-OH	84 ^[c]
10	2a (1)	9i : R = 4-OPiv	10f : R = 4-OPiv	81 ^[c]
		9f	10g	69 ^[d]
11	2b (1)	9f	10g	69 ^[d]
		9a	10h	69
12	2b (1)	9a	10h	69
		9b	10i	93
13	10b (3)	9b	10i	93

[a] Reaction time (h) for the arylation in brackets. [b] Yield of isolated analytically pure product. [c] Pd(OCOCF₃)₂ was used as the Pd source. [d] 2 mol % Pd(OAc)₂, 4 mol % PCy₃ was used. [e] TBDMS = *tert*-butyldimethylsilyl.

with 4-bromoanisole (**9b**) to provide the desired products (**6e,f**) in very high yields (92–99%, Table 1, entries 4 and 5).

To our knowledge, no arylation of 4-picoline (**2a**) has been reported in the literature. However, the smooth zincation of **2a** with **7** (1.5 equiv) proceeds readily, and the palladium-catalyzed cross-coupling of **8b** with various aryl bromides (**9b**, **9e–i**) furnishes the 4-substituted pyridines **10a–f** in 70 to 98% yield (Scheme 2 and Table 1, entries 6–10). 2-Chloro-4-methylpyridine (**2b**) similarly reacts and produces the arylated products **10g** and **10h** in 69% yield (Table 1, entries 11 and 12). Cross-coupling of the 4-substituted pyridine (**10b**) with 4-bromoanisole (**9b**) leads to the desired product (**10i**) in high yield (entry 13). These smooth cross-couplings may be explained by the role that ZnCl₂ plays as a Lewis acid. Interestingly, the use of TMPZnCl·MgCl₂·2LiCl

(prepared from TMPMgCl·LiCl^[19] and ZnCl₂) leads to even faster cross-couplings (at least six times faster). However, the reaction is complicated by increased amounts of diarylation^[20] making the general use of this Lewis acid unattractive. A further hint showing the importance of Lewis acids for the tentative Pd intermediate of type **5** (Scheme 1) is found in the cross-coupling reaction of picolines (**1a** or **2a**) with electron-deficient aryl bromides. Substrates like 4-bromobenzonitrile (**9j**) and ethyl 4-bromobenzoate (**9k**) gave disappointing results in the presence of either ZnCl₂ or MgCl₂ as Lewis acids. Therefore, we screened^[21] other powerful alternative Lewis acids^[22] such as ScCl₃, Sc(OTf)₃,^[23] Yb(OTf)₃,^[24] and Y(OTf)₃.^[22a] Thus, the direct cross-coupling of zincated 2-picoline (**8a**) with 4-bromobenzonitrile (**9j**) gave no product (even after additional ligands for the Pd catalyst were screened).^[25] However, in the presence of 10 mol % Sc(OTf)₃, an efficient palladium-catalyzed cross-coupling took place and afforded the coupling product **6g** in 87% yield (Scheme 3). Similarly, 4-picoline (**2a**) gave the cross-



Scheme 3. Sc(OTf)₃-catalyzed cross-coupling of 2-picoline (**1a**) and 4-picoline (**2a**) with electron-poor aryl bromides **9j** and **k**.

coupling product **10j** only in 41% yield without Sc(OTf)₃, but the addition of 10 mol % Sc(OTf)₃ increased the cross-coupling yield to 78% (Scheme 3). The effect of Sc(OTf)₃ may be best explained by an acceleration of the reductive-elimination step in the cross-coupling as a result of the complexation of Sc(OTf)₃ to the heterocyclic nitrogen (see **11a,b**, Scheme 3). It is anticipated that electron-withdrawing substituents lead to Pd intermediates of type **4** (Scheme 1) which are especially reluctant to undergo reductive elimination. We expect the effect of a Lewis acid to be crucial in these cases. Thus, the cross-couplings of picolines **1a** and **2a** with various electron-deficient aryl bromides (**9j–l**) are dramatically improved by the presence of 10 mol % Sc(OTf)₃ and the cross-coupling products **6h** and **10k–l** are obtained in 75–85% yield. In the absence of Sc(OTf)₃, the yields of the cross-coupling are between 0 and 51% (Table 2, entries 1–3).


Following the mild zincation of picolines and efficient subsequent cross-coupling, we were set to tackle regioselectivity issues in the arylation of dimethylpyridines. Thus, we

Table 2: Effect of $\text{Sc}(\text{OTf})_3$ on the benzylic cross-coupling of 2- and 4-picoline with electron-deficient electrophiles.

Entry	Picoline ^[a]	Aryl-Br	Product ^[b]
1			
2			
3			

[a] Conditions: 50 °C, 1 h. [b] Yield of isolated analytically pure product. [c] Yield of isolated product when the reaction was performed without $\text{Sc}(\text{OTf})_3$.

examined the arylation of 2,3-, 3,4-, and 2,4-lutidines (**12a–c**). With 2,3-lutidine (**12a**), zincation with **7** occurs exclusively at position 2, leading after palladium-catalyzed arylation with 4-bromo-*N,N*-dimethylaniline (**9g**) to the disubstituted pyridine (**13a**) in 85% yield (Scheme 4). Further arylations are



Scheme 4. Selective cross-couplings of lutidines **12a–12c**.^[28]

described in Table 3, entries 1–3. In the case of 3,4-lutidine (**12b**), completely regioselective metalation occurs at position 4 leading after a cross-coupling with 4-bromoanisole (**9b**) to the disubstituted pyridine **14** in 92% yield (Scheme 4). The regioselective arylation of 2,4-lutidine (**12c**) is more challenging since the direct zincation with **7** produces a 2:1 mixture of regioisomers. However, the addition of $\text{BF}_3 \cdot \text{OEt}_2$ ^[15a,26] prior to **7** directs the zincation only at position 4 since the complexation of **12c** with $\text{BF}_3 \cdot \text{OEt}_2$ at the heterocyclic nitrogen hampers the metalation by **7** at position 2 for steric factors. Thus the zincation occurs at position 4 leading after


Table 3: Selective benzylic cross-couplings of lutidines with various aryl bromides, chlorides, and a triflate.

Entry	Lutidine	Aryl-X	Product	Yield [%] ^[a]
1				90
2				91
3				88
4				92 ^[b]
5				82 ^[b]
6				77 ^[b]
7				98 ^[b]
8				92 ^[b]
9				86 ^[b]
10				88 ^[b]

[a] Yield of isolated analytically pure product. [b] $\text{BF}_3 \cdot \text{OEt}_2$ was added prior to $\text{TMPZnCl} \cdot \text{LiCl}$.

cross-coupling with ethyl 4-bromobenzoate (**9k**) to the pyridine (**15a**) in 82% yield. This reactivity is general and several typical aryl bromides, chlorides, and a triflate (**9b,i,j,n–q**) undergo regioselective arylations at position 4 to provide products **15b–g** in 77–98% yield (Table 3, entries 4–10).^[27]

Finally, we briefly examined the arylation of methyl-substituted quinolines (**16a,b**) and isoquinoline (**16c**). With **7** zincation is rapid, and the subsequent palladium-catalyzed arylation proceeds well with various aryl bromides (Scheme 5 and Table 4, entries 1–8). In the case of Negishi cross-couplings with aryl bromides bearing an acidic proton, the use of $\text{Pd}(\text{OCOCF}_3)_2$ introduced by Oshima and Yorimitsu^[4]



Scheme 5. Cross-coupling of quinolines **16a–c** with aryl bromides.

Table 4: Cross-couplings of quinolines with various aryl bromides.

Entry	Quinoline ^[a]	Aryl-X	Product	Yield [%] ^[b]
1	16a (1)	9g	17b : R = 4-NMe ₂	93
2	16a (1)	9j	17c : R = 4-CN	66
3	16a (1)	9l	17d : R = 3-CF ₃	72
4	16a (2)	9h	17e : R = 4-OH	76 ^[c]
5	16a (2)	9s : R = 4-NH ₂	17f : R = 4-NH ₂	74 ^[c]
6	16b (1)	9f	18b : R = 3-Me	96
7	16b (1)	9m	18c : R = 4-CF ₃	86
8	16b (1)	9t : R = 3-F	18d : R = 3-F	95

[a] Reaction time (h) for the arylation in brackets. [b] Yield of the isolated analytically pure product. [c] 2 equiv TMPZnCl-LiCl, 2 mol % Pd(OCOCF₃)₂, and 4 mol % SPhos were used.

was advantageous and ensured high yields and fast cross-couplings (Table 4, entries 4 and 5). The arylation of 2-methylquinoline (**16b**) is best performed with the Xantphos^[29] ligand since the formation of diarylation by-products can be avoided (Table 4, entries 6–8).

In summary, we have reported the direct palladium-catalyzed arylation of methyl-substituted N-heterocycles (pyridines, quinolines, and isoquinoline) promoted by ZnCl₂, Sc(OTf)₃, or BF₃·OEt₂. The action of these Lewis acids may be, in each case, complexation with the heterocyclic nitrogen which facilitates the reductive elimination of the Pd intermediate (ZnCl₂, Sc(OTf)₃). The addition of a Lewis acid such as BF₃·OEt₂ can also trigger the regioselectivity of the 2,4-lutidine metalation. The possibility of improving related Pd cross-couplings by the addition of Lewis acids is currently under investigation in our laboratory.

Received: May 4, 2011

Published online: June 30, 2011

Keywords: cross-coupling · metalation · pyridine · zinc

- [1] a) K. C. Nicolaou, R. Scarpelli, B. Bollbuck, B. Werschkun, M. M. A. Pereira, M. Wartmann, K.-H. Altmann, D. Zaharevitz, R. Gussio, P. Giannakakou, *Chem. Biol.* **2000**, *7*, 593; b) B. Oliva, K. Miller, N. Caggiano, A. J. O'Neill, G. D. Cuny, M. Z. Hoemann, J. R. Hauske, I. Chopra, *Antimicrob. Agents Chemother.* **2003**, *47*, 458; c) A. Bouillon, A. S. Voisin, A. Robic, J.-C. Lancelot, V. Collot, S. Rault, *J. Org. Chem.* **2003**, *68*, 10178; d) E. M. Nolan, J. Jaworski, K.-I. Okamoto, Y. Hayashi, M. Sheng, S. J. Lippard, *J. Am. Chem. Soc.* **2005**, *127*, 16812; e) A. Hayashi, M. Arai, M. Fujita, M. Kobayashi, *Biol. Pharm. Bull.* **2009**, *32*, 1261; f) J. Quiroga, J. Trilleras, B. Insuasty, R. Abonia, M. Nogueras, A. Marchal, J. Cobo, *Tetrahedron Lett.* **2010**, *51*, 1107; g) T. Laird, *Org. Process Res. Dev.* **2006**, *10*, 851; h) M. A. Yurovskaya, A. V. Karchava, *Chem. Heterocycl. Compd.* **1994**, *30*, 1331; i) P. N. W. Baxter, J.-M. Lehn, J. Fischer, M.-T. Youinou, *Angew. Chem.* **1994**, *106*, 2432; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 2284; j) J.-M. Lehn, *Science* **2002**, *295*, 2400; k) X.

- Chen, K. M. Engle, D.-H. Wang, J. Q. Yu, *Angew. Chem.* **2009**, *121*, 5196; *Angew. Chem. Int. Ed.* **2009**, *48*, 5094.
- [2] a) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174; b) For a special issue on Selective Functionalization of C-H Bonds: *Chem. Rev.* **2010**, *110*, 575; c) D. Kalyani, N. R. Deprez, L. V. Desai, M. S. Sanford, *J. Am. Chem. Soc.* **2005**, *127*, 7330; d) S. Murai, *Activation of Unreactive Bonds and Organic Synthesis*, Springer, Berlin, **1999**; e) A. R. Dick, M. S. Sanford, *Tetrahedron* **2006**, *62*, 2439; f) M. Schlosser, F. Mongin, *Chem. Soc. Rev.* **2007**, *36*, 1161; g) T. Niwa, H. Yorimitsu, K. Oshima, *Org. Lett.* **2007**, *9*, 2373.
- [3] P. M. Burton, J. A. Morris, *Org. Lett.* **2010**, *12*, 5359.
- [4] T. Niwa, H. Yorimitsu, K. Oshima, *Angew. Chem.* **2007**, *119*, 2697; *Angew. Chem. Int. Ed.* **2007**, *46*, 2643.
- [5] a) L.-C. Campeau, D. J. Schipper, K. Fagnou, *J. Am. Chem. Soc.* **2008**, *130*, 3266; b) D. J. Schipper, L.-C. Campeau, K. Fagnou, *Tetrahedron* **2009**, *65*, 3155.
- [6] J. J. Mousseau, A. Larivée, A. B. Charette, *Org. Lett.* **2008**, *10*, 1641.
- [7] For structures of palladium picolyl derivatives, see: a) M. Onishi, K. Hiraki, K. Maeda, T. Itoh, *J. Organomet. Chem.* **1980**, *188*, 245; b) K. Isobe, Y. Nakamura, S. Kawaguchi, *Bull. Chem. Soc. Jpn.* **1989**, *62*, 1802; c) B. Qian, S. Guo, J. Shao, Q. Zhu, L. Yang, C. Xia, H. Huang, *J. Am. Chem. Soc.* **2010**, *132*, 3650.
- [8] Q. Shen, J. F. Hartwig, *J. Am. Chem. Soc.* **2007**, *129*, 7734.
- [9] J. Huang, C. M. Haar, S. P. Nolan, *Organometallics* **1999**, *18*, 297.
- [10] For additions to imines, see: a) B. Qian, S. Guo, C. Xia, H. Huang, *Adv. Synth. Catal.* **2010**, *352*, 3195; b) M. Rueping, N. Tolstoluzhsky, *Org. Lett.* **2011**, *13*, 1095.
- [11] For additions to enones, see: H. Komai, T. Yoshino, S. Matsunaga, M. Kanai, *Org. Lett.* **2011**, *13*, 1706.
- [12] a) S. Wunderlich, P. Knochel, *Angew. Chem.* **2007**, *119*, 7829; *Angew. Chem. Int. Ed.* **2007**, *46*, 7685; b) S. Wunderlich, P. Knochel, *Org. Lett.* **2008**, *10*, 4705; c) S. Wunderlich, P. Knochel, *Chem. Commun.* **2008**, 6387; d) M. Mosrin, P. Knochel, *Org. Lett.* **2009**, *11*, 1837; e) M. Mosrin, G. Monzon, T. Bresser, P. Knochel, *Chem. Commun.* **2009**, 5615; f) C. J. Rohbogner, S. Wirth, P. Knochel, *Org. Lett.* **2010**, *12*, 1984; g) T. Bresser, M. Mosrin, G. Monzon, P. Knochel, *J. Org. Chem.* **2010**, *75*, 4686.
- [13] a) M. L. Hlavinka, J. R. Hagadorn, *Tetrahedron Lett.* **2006**, *47*, 5049; b) M. L. Hlavinka, J. R. Hagadorn, *Organometallics* **2007**, *26*, 4105.
- [14] S. Duez, S. Bernhardt, J. Heppekausen, F. F. Fleming, P. Knochel, *Org. Lett.* **2011**, *13*, 1690.
- [15] a) M. Jaric, B. A. Haag, A. Unsinn, K. Karaghiosoff, P. Knochel, *Angew. Chem.* **2010**, *122*, 5582; *Angew. Chem. Int. Ed.* **2010**, *49*, 5451; b) For an excellent review see: D. W. Stephan, G. Erker, *Angew. Chem.* **2010**, *122*, 50; *Angew. Chem. Int. Ed.* **2010**, *49*, 46.
- [16] a) E.-i. Negishi, *Metal-Catalyzed Cross-Coupling Reactions* (Eds.: F. Diederich, P. J. Stang), Wiley, New York, **1998**, chap. 1; b) E. Erdik, *Tetrahedron* **1992**, *48*, 9577; c) E.-i. Negishi, L. F. Valente, M. Kobayashi, *J. Am. Chem. Soc.* **1980**, *102*, 3298; d) E.-i. Negishi, *Acc. Chem. Res.* **1982**, *15*, 340.
- [17] Usually, 1.5 equiv of base **7** is used. In this case, 2.0 equiv of base is used because of the presence of the indole NH group.
- [18] a) S. D. Walker, T. E. Barder, J. R. Martinelli, S. L. Buchwald, *Angew. Chem.* **2004**, *116*, 1907; *Angew. Chem. Int. Ed.* **2004**, *43*, 1871; b) T. E. Barder, S. D. Walker, J. R. Martinelli, S. L. Buchwald, *J. Am. Chem. Soc.* **2005**, *127*, 4685; c) R. A. Altman, S. L. Buchwald, *Nat. Protoc.* **2007**, *2*, 3115; d) R. Martin, S. L. Buchwald, *Acc. Chem. Res.* **2008**, *41*, 1461.
- [19] a) A. Krasovskiy, V. Krasovskaya, P. Knochel, *Angew. Chem.* **2006**, *118*, 3024; *Angew. Chem. Int. Ed.* **2006**, *45*, 2958; b) R. Mulvey, F. Mongin, M. Uchiyama, Y. Kondo, *Angew. Chem.* **2007**, *119*, 3876; *Angew. Chem. Int. Ed.* **2007**, *46*, 3802.
- [20] The palladium-catalyzed arylation of 2-picoline (**1a**) with 3-chlorobromobenzene (**9d**) proceeds in 1 h in the presence of

TMPZnCl·MgCl₂·LiCl instead of in 6 h in the presence of TMPZnCl·LiCl (**7**) but led to less product **6d** (61 %) owing to the formation of 17 % of diarylated product.

- [21] See the Supporting Information for the screening of various Lewis acids.
- [22] a) H. Yamamoto, *Lewis Acid Reagents: A Practical Approach*, Oxford University Press, New York, **1999**; b) D. Schinzer, *Selectivities in Lewis acid promoted reactions*, Kluwer Academic, Dordrecht, **1989**; c) S. Kobayashi, *Lanthanides: Chemistry and Use in Organic Synthesis*, Springer, Berlin, **1999**; d) M. Shibasaki, S. Matsunaga, N. Kumagai, *Acid Catalysis in Modern Organic Synthesis, Vol. 2*, Wiley-VCH, Weinheim, **2008**, p. 635.
- [23] a) C. Ogawa, Y. Gu, M. Boudou, S. Kobayashi, *Acid Catalysis in Modern Organic Synthesis, Vol. 2*, Wiley-VCH, Weinheim, **2008**, p. 589; b) S. Kobayashi, *Lewis Acids in Organic Synthesis, Vol. 2*, Wiley-VCH, Weinheim, **2000**, p. 883; c) S. Kobayashi, I. Hachiya, M. Araki, H. Ishitani, *Tetrahedron Lett.* **1993**, *34*, 3755.
- [24] a) S. Kobayashi, *Transition Metals for Organic Synthesis, Vol. 1*, Wiley, Weinheim, **1998**, p. 285; b) S. Kobayashi, M. Sugiura, H. Kitagawa, W. L. Lam, *Chem. Rev.* **2002**, *102*, 2227.
- [25] Screenings of various Pd sources such as Pd(OAc)₂, [Pd(dba)₂], and Pd(OCOCF₃)₂ in combination with ligands like SPhos, Xantphos, Davephos, binap, tfp,^[30] PCy₃, and PtBu₃ as well as PEPPSI,^[31] [Pd(PPh₃)₄], and [PdCl₂(dppf)] did not give satisfactory results.
- [26] M. Jaric, B. A. Haag, S. M. Manolikakes, P. Knochel, *Org. Lett.* **2011**, DOI: 10.1021/ol200563j.
- [27] Interestingly, the less bulky base *i*Pr₂NZnCl·LiCl is not compatible with the Lewis acid BF₃·OEt₂, since it forms the stable complex *i*Pr₂NBF₃ZnCl, which does not dissociate reversibly under the reaction conditions, unlike the frustrated Lewis acid pair TMPBF₃ZnCl.^[15b]
- [28] The presence of *N*-ethylpyrrolidinone (NEP) leads to shorter reaction times (three times faster).
- [29] P. W. N. M. van Leeuwen, P. C. J. Kamer, J. N. H. Reek, P. Dierkes, *Chem. Rev.* **2000**, *100*, 2741.
- [30] N. G. Andersen, B. A. Keay, *Chem. Rev.* **2001**, *101*, 997.
- [31] a) C. J. O'Brien, E. A. B. Kantchev, C. Valente, N. Hadei, G. A. Chass, A. Lough, A. C. Hopkinson, M. G. Organ, *Chem. Eur. J.* **2006**, *12*, 4743; b) M. G. Organ, S. Avola, I. Dubovyk, N. Hadei, E. A. B. Kantchev, C. J. O'Brien, C. Valente, *Chem. Eur. J.* **2006**, *12*, 4749.