

Pd-Catalyzed Coupling of Thioamides with *N*-Tosylhydrazones/Trapping by Esters Cascade Reaction

Zhongliang Cai,[†] Zhi Yao,[†] and Liqin Jiang*



Cite This: *Org. Lett.* 2021, 23, 311–316



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: *N,N*-Disubstituted thioamides coupled with *N*-tosylhydrazones under $\text{Pd}(\text{TFA})_2/t\text{BuXPhos}$ catalyst and NaOtBu , and the intermediates from palladium carbene migratory insertion containing β -hydrogen were trapped by intramolecular esters activated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ instead of undergoing β -H elimination, providing polyfunctional thiophen-3(2*H*)-ones with sulfur-containing tetrasubstituted carbon centers in moderate to good yields. The reaction features the formation of three bonds in a single operation, odorless, safe, and easily available substrates, wide substrate scope, and excellent functional group tolerance.


Sulfur-containing heterocycles exist widely in natural products, drugs, biological compounds, and materials.¹ Odorless, nontoxic, and easily available starting materials for preparing sulfur-containing heterocycles are highly desirable. Recently, we reported the ready synthesis of *N,N*-disubstituted thioamides bearing α,α -diesters from malonate esters and thiocarbamoyl fluorides² in which the sulfur atom was derived from sulfur (S_8).³ Thioamides, which are odorless and nontoxic/low toxic, have been widely used to synthesize sulfur-containing heterocycles via intramolecular coupling reactions including C–H functionalizations.^{2,4} Metal-catalyzed intramolecular or intermolecular reactions of thioamides with hydrazones or α -diazocarbonyl compounds to transform $\text{C}=\text{S}$ to $\text{C}=\text{C}$ bonds have also been reported.⁵ However, intermolecular reactions involving thioamides in a single operation for the construction of complex and diversified heterocycles with sulfur-containing tetrasubstituted carbon centers are underdeveloped. Sulfur heterocycles in this class exist in natural products such as thiolactomycin and thiotetraamide as well as in biological compounds demonstrating anticancer,^{1d} weight loss,^{1d} antibacterial,^{6a–c} antimalarial,^{6d,e} antituberculosis,^{6f,g} and pesticidal activities.^{6h}


The Hu group pioneered the development of a series of multicomponent reactions of trapping of ylides or zwitterionic intermediates derived from nucleophiles and diazocarbonyl compounds by electrophiles such as aldehydes, ketones, imines, or α,β -unsaturated compounds to form tetrasubstituted carbon centers.⁷ The challenge of these methods lies in the 1,2-proton transfer in the active intermediates. Esters have never

been documented to realize the trapping of active intermediates as above. Due to their low electrophilicity, competitive 1,2-proton transfer might be faster. Dual-metal-catalyzed carbene sp^2 C–H functionalization/Conia–ene cascade reactions have also been reported.⁸ Diazo compounds bearing alkyls have never been used in the above reactions due to their instability and difficulty in separation and purification. *N*-Tosylhydrazones are safe and easily available precursors of diazo compounds and have attracted considerable attention.⁹ Palladium-catalyzed cross-couplings involving *N*-tosylhydrazones bearing alkyl groups normally give alkenyl products via β -H elimination of the metal alkyl intermediates generated from palladium carbene migratory insertion.^{9a,b} For example, Yamaguchi reported that thioesters reacted with *N*-tosylhydrazones under Pd-catalysis to furnish *Z*-alkenyl thioethers (Scheme 1, previous work, 1).¹⁰ Reactions of heteroatom nucleophiles (X–H) with *N*-tosylhydrazones under transition-metal free conditions could lead to X–H insertion products. For example, thiophenols/mercaptans coupled with *N*-tosylhydrazones under basic conditions to give S–H insertion products (Scheme 1, previous work, 2).¹¹ Under photoredox conditions, thiophenols/mercaptans and *N*-tosylhydrazones have been

Received: November 15, 2020

Published: January 6, 2021



Scheme 1. Reactions of Sulfur Reagents with *N*-Tosylhydrazones

reported to form carbanion intermediates, which can then be trapped by CO_2 or aldehydes but cannot be trapped by ketones (Scheme 1, previous work, 3).¹² Our question was

whether the palladium intermediates containing β -hydrogens, formed via palladium carbene migratory insertion, could be trapped by more challenging esters instead of undergoing β -hydride elimination. To the best of our knowledge, the formation of a second bond at the same carbon of the palladium carbene, which also contains β -hydrogens, via 1,2 addition or 1,2 addition/elimination with electrophiles such as esters has never been explored and accomplished.

We envision expanding the use of *N,N*-disubstituted thioamides bearing α,α -diesters **1** to intermolecular reactions with *N*-tosylhydrazones **2** (Scheme 1, this work). Under suitable catalytic systems, a base abstracts the α -H of thiocarbonyl of **1** to form thioenolate anion **I**, which coordinates with the metal of the carbene **III** derived from *N*-tosylhydrazone **2** to give the intermediate **V**, followed by carbene migratory insertion to give **VI**. The resulting intermediate **VI** or the anion derived from **VI** losing metal might then undergo cascade intramolecular 1,2-addition/elimination with an ester functionality activated by a co-catalyst, selectively delivering polyfunctional thiophen-3(2*H*)-one **3** (Scheme 1, this work). Various undesired pathways exist that could minimize or halt the formation of the desired products. For instance, β -H elimination or protonation of **VI** yields **4** or **5**. *N*-Tosylhydrazones **2** might also self-condense to form N-H insertion products **6**. Thioamides **1** might also react with **2** to transform the C=S bond into a C=C bond, furnishing enamines **7** (Scheme 1, this work, other possible products). Herein, we disclose that *N,N*-disubstituted thioamides **1** couple with *N*-tosylhydrazones **2** bearing alkyl substituents under Pd(II) catalysis, ^{*t*}BuXPhos as the ligand,

Table 1. Optimization of the Reaction Conditions for **3**^a

entry	catalyst	ligand	base	additive	T (°C)	solvent	yield ^b (%)
1 ^c	Pd(TFA) ₂		NaO ^{<i>t</i>} Bu ^d		90	MeCN	23
2 ^c	Rh ₂ (OAc) ₄		NaO ^{<i>t</i>} Bu ^d		90	MeCN	<5%
3 ^c	AgTFAfA		NaO ^{<i>t</i>} Bu ^d		90	MeCN	9
4 ^c	CuI		NaO ^{<i>t</i>} Bu ^d		90	MeCN	12
5 ^c	Pd(TFA) ₂	BINAP	NaO ^{<i>t</i>} Bu ^d		90	MeCN	14
6 ^c	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu ^d		90	MeCN	39
7 ^c	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu ^d		100	MeCN/toluene (1:1)	45
8 ^c	Pd(TFA) ₂	X-Phos	LiO ^{<i>t</i>} Bu ^d		100	MeCN/toluene (1:1)	29
9 ^c	Pd(TFA) ₂	X-Phos	NaH ^d		100	MeCN/toluene (1:1)	28
10 ^c	Pd(TFA) ₂	X-Phos	NaOSiMe ₃ ^d		100	MeCN/toluene (1:1)	48
11	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu		100	MeCN/toluene (1:1)	50
12	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu	FeCl ₃	100	MeCN/toluene (1:1)	62
13	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu	AlCl ₃	100	MeCN/toluene (1:1)	54
14	Pd(TFA) ₂	X-Phos	NaO ^{<i>t</i>} Bu	BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	65
15	Pd(TFA) ₂	^{<i>t</i>} BuXPhos	NaO ^{<i>t</i>} Bu	BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	70
16	Pd(TFA) ₂	^{<i>t</i>} BuXPhos	NaOSiMe ₃	BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	0
17		^{<i>t</i>} BuXPhos	NaO ^{<i>t</i>} Bu	BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	6
18	Pd(TFA) ₂	^{<i>t</i>} BuXPhos		BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	0
19	Pd(TFA) ₂		NaO ^{<i>t</i>} Bu	BF ₃ ·Et ₂ O	100	MeCN/toluene (1:1)	41
20	Pd(TFA) ₂	^{<i>t</i>} BuXPhos	NaO ^{<i>t</i>} Bu		100	MeCN/toluene (1:1)	55


^aReaction conditions: **1a** (0.15 mmol), **2a** (0.375 mmol), 10 mol % of catalyst, 20 mol % of ligand, 50 mol % of additive, NaO^{*t*}Bu (0.6 mmol) reacted in 3 mL of anhydrous solvent for 12 h. ^bIsolated yield. ^c**2a** (0.3 mmol, 2.0 equiv). ^dNaO^{*t*}Bu (0.3 mmol, 2.0 equiv). X-Phos: 2-(dicyclohexylphosphino)-2',4',6'-tri(isopropyl)biphenyl; ^{*t*}BuXPhos: 2-di-*tert*-butylphosphino-2',4',6'-tri(isopropyl)biphenyl.

and with $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as the co-catalyst to selectively access polyfunctional thiophen-3(2H)-ones with sulfur-containing tetrasubstituted carbon centers in moderate to good yields.

We started our studies with **1a** and **2a** as model substrates. Selected results are shown in Table 1 (details are provided in the Supporting Information (SI)). Initially, palladium catalysts including $\text{Pd}(\text{TFA})_2$, $\text{Pd}(\text{OAc})_2$, $\text{Pd}_2(\text{dba})_3$, and $\text{Pd}(\text{PPh}_3)_4$ were investigated in the reaction of 0.15 mmol of **1a**, 2.0 equiv of **2a**, and 2.0 equiv of NaO^tBu in the presence of 10 mol % of palladium catalyst in MeCN at 90 °C under N_2 (Table 1, entry 1, and SI). $\text{Pd}(\text{TFA})_2$ was optimal and afforded the desired product **3a** in 23% yield (Table 1, entry 1, and SI). Catalysts based on other metals such as Rh, Ag, and Cu were tried but led to inferior results compared to $\text{Pd}(\text{TFA})_2$ (Table 1, entries 2–4, and SI). Then we attempted to investigate the effect of the ligands under $\text{Pd}(\text{TFA})_2$ during this transformation. 20 mol % of ligand including BINAP, dppp, dppb, Xantphos, PCy_3 , PPh_3 , or X-Phos was added to the entry 1 conditions (Table 1, entries 5, 6, and SI). It was found that X-Phos was optimal (Table 1, entry 6). Next, different solvents and temperatures were screened. It was found that mixed solvents of MeCN and toluene (1:1) at 100 °C (oil bath temperature) improved the yield of **3a** to 45% (Table 1, entry 7) and N–H insertion product **6a** (21% yield) was the main byproduct. Evaluation of other bases revealed that only NaOSiMe_3 gave a slightly higher yield of **3a** than NaO^tBu (Table 1, entries 8–10, and SI), but further study demonstrated that NaOSiMe_3 was not compatible with Lewis acid additives (SI). We improved NaO^tBu to 4.0 equiv and **2a** to 2.5 equiv, leading to **3a** in 50% yield with **8a** (main byproduct) in 25% yield (Table 1, entry 11). We found that introducing an additive such as FeCl_3 , AlCl_3 , or $\text{BF}_3 \cdot \text{Et}_2\text{O}$ to activate the ester functionality (Table 1, entries 12–14) under NaO^tBu to facilitate the desired pathway and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ was superior (Table 1, entries 12–14). Finally, ligands including Xantphos, DPEphos, DavePhos, $^t\text{BuXPhos}$, S-Phos, and RuPhos were screened (Table 1, entry 15, and SI). $^t\text{BuXPhos}$ furnished the desired **3a** in 70% yield (Table 1, entry 15), which was identified as the optimized conditions for **3a**. The use of sealing tube under the otherwise same conditions afforded **3a** in 71% yield (SI). Decreasing the amounts of catalyst or ligand both led to a decrease in the yield of **3a** (SI). Deletion experiments demonstrated that $\text{Pd}(\text{TFA})_2$ and NaO^tBu were crucial for this cascade reaction (entries 17 and 18), and $^t\text{BuXPhos}$ and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ improved the reactivity and selectivity for **3a** (Table 1, entries 19 and 20).

With the optimized conditions in hand, we investigated the substrate scope of the novel cascade reactions of *N,N*-disubstituted thioamides **1** and *N*-tosylhydrazones **2** (Scheme 2). *N*-Tosylhydrazones **2** were first investigated. *N*-Tosylhydrazones **2** bearing *para* (**2b–2g**)-, *meta* (**2h–2j**)-, or *ortho* (**2k**)-electron-withdrawing substituted aryls proceeded smoothly to afford the corresponding products **3b–3k** in moderate to good yields. It is important to note that fluoro (**2b**, **2k**), chloro (**2c**, **2h**), bromo (**2d**, **2i**), trifluoromethyl (**2e**, **2j**), ester (**2f**), and even cyano (**2g**) were all well tolerated. Fluoro, chloro, and bromo could serve as handles for further elaboration, and trifluoromethyl is a privileged group in medicinal chemistry. The reactivity of **2l** bearing electron-donating *p*-methoxy-substituted aryl with **1a** was sluggish, but still moderate yields of the desired product **3l** could be obtained via slightly improved amounts of the catalyst and the ligand. *N*-Tosylhydrazone **2m** bearing heterocycle 2-thienyl was also tolerated. *N*-Tosylhydrazone **2n** bearing a longer alkyl

Scheme 2. Substrate Scope of the Cascade Reactions^a



^aReaction conditions: **1** (0.15 mmol), **2** (0.375 mmol), 10 mol % of $\text{Pd}(\text{TFA})_2$, 20 mol % of $^t\text{BuXPhos}$, 50 mol % of $\text{BF}_3 \cdot \text{Et}_2\text{O}$, NaO^tBu (0.6 mmol) reacted in 3 mL of MeCN and toluene (1:1) for 12 h. ^bIsolated yield. ^c1 mmol. ^d15 mol % of $\text{Pd}(\text{TFA})_2$, 30 mol % of $^t\text{BuXPhos}$. ^e15 mol % of $\text{Pd}(\text{OAc})_2$, 30 mol % of $\text{P}(\text{2-furyl})_3$, **2** (4.0 equiv), NaO^tBu (5.5 equiv).


was relatively inert, furnishing the desired product in low yield (10%). However, a moderate yield of 37% of **3n** could be obtained by the use of $\text{Pd}(\text{OAc})_2/\text{P}(\text{2-furyl})_3$ catalytic system with improving amounts of **2n** and the base. It is worth noting that *N*-tosylhydrazones (**2o** and **2p**) bearing two aryls were also compatible in the cascade reaction (**3o** and **3p**). In addition, *N*-tosylhydrazone **2q** bearing a phenyl and an ester as

the safe precursor of methyl phenyldiazoacetate afforded the desired product **3q** in 54% yield.

Then we turned our attention to the scope of the substrate **1**. As illustrated in Scheme 2, an array of *N,N*-disubstituted thioamides bearing α,α -diesters **1** could bifunctionalize a single carbon of *N*-tosylhydrazones **2** smoothly. Thioamides **1** bearing other esters, including ethyl esters (**1r**), isopropyl esters (**1s**), and benzyl esters (**1t**), were compatible (**3r–3t**). Thioamides **1** with *para*- or *meta*-electron-donating and electron-withdrawing substituted aryls on nitrogen both afforded the desired product **3u–3aa** in moderate to good yields. Methoxy (**1u**, **1z**), fluoro (**1v**), chloro (**1w**, **1aa**), trifluoromethyl (**1x**), and keto (**1y**) in aryls on nitrogen of **1** were all tolerated. In addition, Thioamides **1** containing other alkyl substitutions on nitrogen, including ethyl (**3ab**), isopropyl (**3ac**), *n*-hexyl (**3ad**), and 3-methoxypropyl (**3ae**), were all well tolerated, smoothly delivering the corresponding products **3ab–3ae** in moderate to good yields.

A one-pot reaction from 2.5 equiv of acetophenone, 2.5 equiv of 4-methylbenzenesulfonylhydrazide (which form **2a** followed by the removal of solvent), and 1.0 equiv of **1a** worked to afford **3a** in 61% yield (Scheme 3, a). Derivatization


Scheme 3. One-Pot Protocol and Product Derivatization



of the products was investigated. Product **3a** was reduced by NaBH_4 to furnish **9a** and **9a'** in 53% total yield with 4:1 diastereoselectivity, and the stereochemistry was assigned using 1D NOESY spectra of **9a** and **9a'** (Scheme 3 (b, 1) and SI). Product **3a** reacted with 1.2 equiv of EtMgBr and 2.0 equiv of MeLi under -78°C , and only the ester group in **3a** was transformed into ketone to furnish **10a** in moderate yield (Scheme 3 (b, 2)). As mentioned above, the reaction is compatible with halogen atoms which can be used as handles for functionalization. Product **3d** reacted with phenylboronic acid under the reaction conditions reported by Buchwald¹³ to afford **11a** in 87% yield (Scheme 3 (b, 3)). In addition, some chiral ligands instead of *t*BuXPhos were investigated, but none

of the enantioselective product **3a** was obtained (see the details in the SI). Thus, the anions VII (Scheme 4) are more likely to be the trapped intermediates in this cascade process.

Scheme 4. Proposed Mechanism



A plausible mechanism for this novel reaction is proposed in Scheme 4. NaO^tBu abstracts the α -H of thioacyl of **1** to form the thioenolate anion **I**, which might coordinate with the palladium catalyst to give the palladium complex **IV**. The complex **IV** and the diazo compound **II** derived from *N*-tosylhydrazone **2** under NaO^tBu might form the palladium carbene **V**, which undergoes carbene migratory insertion to give the intermediate **VI**. The organopalladium moiety in **VI** attacks the intramolecular ester activated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ to afford product **3**. Also, intermediate **VI** might produce the anion **VII** along with recovery of $\text{Pd(TFA)}_2/t\text{BuXPhos}$, and then the anion **VII** attacks the intramolecular ester activated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ to afford the product **3** via addition/elimination. Alternatively, $\text{Pd(TFA)}_2/t\text{BuXPhos}$ reacts with the diazo compound **II** derived from *N*-tosylhydrazone **2** to give the metal carbene intermediate **III**. Then the thioenolate anion **I** derived from **1** might undergo ligand exchange with **III** to give the intermediate **V**, which forms **VI** via carbene migratory insertion followed by the similar intramolecular cyclization as above to afford **3**. In addition, the thioenolate anion **I** directly attacks the carbene carbon of **III** also might form the intermediate **VI**, and then the intermediate **VI** directly or the anion **VII** derived from **VI** undergoes cyclization to furnish **3**.

In conclusion, *N,N*-disubstituted thioamides bearing α,α -diesters coupled with *N*-tosylhydrazones bearing alkyls with Pd(TFA)_2 as the catalyst, *t*BuXPhos as the ligand, and NaO^tBu as the base at 100°C and the formed intermediates were unprecedentedly trapped by intramolecular ester functionalities activated by co-catalyst $\text{BF}_3 \cdot \text{Et}_2\text{O}$, affording polyfunctional thiophen-3(2*H*)-ones with sulfur-containing tetrasubstituted carbon centers in moderate to good yields. This represents the first time that the formed organopalladium intermediates containing β -hydrogen from palladium carbene migratory insertion selectively form the second bond at the same carbon via 1,2-addition/elimination with esters instead of undergoing β -hydrogen eliminations. The novel cascade reaction utilizes odorless, nontoxic, safe, and easily available starting materials. Wide substrate scope and a broad range of functional groups are tolerated. Products possess rich and valuable structures in medicinal chemistry and materials. Developing other novel reactions to selectively construct

multiple bonds in a single process with odorless and nontoxic sulfur reagents as well as evaluation of the biological activities of novel thiophen-3(2*H*)-ones are currently being investigated in our laboratory and in our cooperative laboratory.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c03796>.

Experimental procedures and spectroscopic characterization data, ^1H and ^{13}C NMR spectra of the new compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Liqin Jiang – School of Chemistry and Molecular Engineering, East China Normal University, Shanghai 200241, China;
orcid.org/0000-0002-9751-7510; Email: lqjiang@sat.ecnu.edu.cn, jiangliqin_777@163.com

Authors

Zhongliang Cai – School of Chemistry and Molecular Engineering, East China Normal University, Shanghai 200241, China

Zhi Yao – School of Chemistry and Molecular Engineering, East China Normal University, Shanghai 200241, China

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.0c03796>

Author Contributions

† Z.C. and Z.Y. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for financial support from the Science and Technology Commission of Shanghai Municipality (No. 15ZR1410500).

■ REFERENCES

- (1) (a) Nicolaou, K. C.; Hale, C. R. H.; Nilewski, C.; Ioannidou, H. A. Constructing Molecular Complexity and Diversity: Total Synthesis of Natural Products of Biological and Medicinal Importance. *Chem. Soc. Rev.* **2012**, *41*, 5185–5238. (b) *Sulphur-Containing Drugs and Related Organic Compounds*; Damani, L. A., Ed.; Wiley: New York, 1989. (c) Ilardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-Mining for Sulfur and Fluorine: An Evaluation of Pharmaceuticals to Reveal Opportunities for Drug Design and Discovery. *J. Med. Chem.* **2014**, *57*, 2832–2842. (d) McFadden, J. M.; Medghalchi, S. M.; Thupari, J. N.; Pinn, M. L.; Vadlamudi, A.; Miller, K. I.; Kuhajda, F. P.; Townsend, C. A. Application of a Flexible Synthesis of (5*R*)-Thiolactomycin To Develop New Inhibitors of Type I Fatty Acid Synthase. *J. Med. Chem.* **2005**, *48*, 946–961. (e) Suhas, R.; Chandrashekar, S.; Gowda, D. C. Synthesis of Uriedo and Thiouriedo Derivatives of Peptide Conjugated Heterocycles - A New Class of Promising Antimicrobials. *Eur. J. Med. Chem.* **2012**, *48*, 179–191. (f) Rowley, J. A.; Reid, R. C.; Poon, E. K. Y.; Wu, K.-C.; Lim, J.; Lohman, R.-J.; Hamidon, J. K.; Yau, M.-K.; Halili, M. A.; Durek, T.; Iyer, A.; Fairlie, D. P. Potent Thiophene Antagonists of Human Complement C3a Receptor with Anti-Inflammatory Activity. *J. Med. Chem.* **2020**, *63*, 529–541. (g) Yang, Z.; Mao, Z.; Xie, Z.; Zhang, Y.; Liu, S.; Zhao, J.; Xu, J.; Chi, Z.; Aldred, M. P. Recent Advances in

Organic Thermally Activated Delayed Fluorescence Materials. *Chem. Soc. Rev.* **2017**, *46*, 915–1016.

(2) Yao, Z.; Cai, Z.; Zhen, L.; Jiang, L. Cobalt-Catalyzed Switchable Intramolecular Thioenolization/C–H Thiolation and C(sp²)-H/C(sp³)-H Dehydrogenative Coupling. *Org. Lett.* **2020**, *22*, 4505–4510.

(3) Zhen, L.; Fan, H.; Wang, X.; Jiang, L. Synthesis of Thiocarbonyl Fluorides and Isothiocyanates Using CF₃SiMe₃ and Elemental Sulfur or AgSCF₃ and KBr with Amines. *Org. Lett.* **2019**, *21*, 2106–2110.

(4) (a) Shen, C.; Zhang, P.; Sun, Q.; Bai, S.; Hor, T. S. A.; Liu, X. Recent Advances in C–S Bond Formation via C–H Bond Functionalization and Decarboxylation. *Chem. Soc. Rev.* **2015**, *44*, 291–314. (b) Zhang, G.; Liu, C.; Yi, H.; Meng, Q.; Bian, C.; Chen, H.; Jian, J.-X.; Wu, L.-Z.; Lei, A. External Oxidant-Free Oxidative Cross-Coupling: A Photoredox Cobalt-Catalyzed Aromatic C–H Thiolation for Constructing C–S Bonds. *J. Am. Chem. Soc.* **2015**, *137*, 9273–9280. (c) Huang, C.; Qian, X.-Y.; Xu, H.-C. Continuous-Flow Electrosynthesis of Benzofused S-Heterocycles by Dehydrogenative C–S Cross-Coupling. *Angew. Chem., Int. Ed.* **2019**, *58*, 6650–6653.

(5) (a) Fang, F. G.; Danishefsky, S. J. The Total Synthesis of Chilenine: Novel Constructions of Cyclic Enamides. *Tetrahedron Lett.* **1989**, *30*, 2747–2750. (b) Koduri, N. D.; Wang, Z.; Cannell, G.; Cooley, K.; Lemma, T. M.; Miao, K.; Nguyen, M.; Frohock, B.; Castaneda, M.; Scott, H.; Albinescu, D.; Hussaini, S. R. Enaminones via Ruthenium-Catalyzed Coupling of Thioamides and α -Diazocarbonyl Compounds. *J. Org. Chem.* **2014**, *79*, 7405–7414. (c) Mohammadi, L.; Zolfigol, M. A.; Ebrahimi, M.; Roberts, K. P.; Ansari, S.; Azadbakht, T.; Hussaini, S. R. A Copper(I)-Complexed Magnetic Nanoparticle Catalyst for Enaminone Synthesis. *Cat. Commun.* **2017**, *102*, 44–47. (d) Pal, A.; Koduri, N. D.; Wang, Z.; Quiroz, E. L.; Chong, A.; Vuong, M.; Rajagopal, N.; Nguyen, M.; Roberts, K. P.; Hussaini, S. R. Copper-Catalyzed Chemoselective Cross-Coupling Reaction of Thioamides and α -Diazocarbonyl Compounds: Synthesis of Enaminones. *Tetrahedron Lett.* **2017**, *58*, 586–589.

(6) (a) Bommineni, G. R.; Kapilashrami, K.; Cummings, J. E.; Lu, Y.; Knudson, S. E.; Gu, C.; Walker, S. G.; Slayden, R. A.; Tonge, P. J. Thiolactomycin-Based Inhibitors of Bacterial β -Ketoacyl-ACP Synthases with in Vivo Activity. *J. Med. Chem.* **2016**, *59*, 5377–5390. (b) Rothe, M. L.; Li, J.; Garibay, E.; Moore, B. S.; McKinnie, S. M. K. Synthesis, Bioactivity, and Enzymatic Modification of Antibacterial Thiotetromycin Derivatives. *Org. Biomol. Chem.* **2019**, *17*, 3416–3423. (c) Li, J.; Tang, X.; Awakawa, T.; Moore, B. S. Enzymatic C–H Oxidation-Amidation Cascade in the Production of Natural and Unnatural Thiotetronate Antibiotics with Potentiated Bioactivity. *Angew. Chem., Int. Ed.* **2017**, *56*, 12234–12239. (d) Jones, S. M.; Urch, J. E.; Kaiser, M.; Brun, R.; Harwood, J. L.; Berry, C.; Gilbert, I. H. Analogues of Thiolactomycin as Potential Antimalarial Agents. *J. Med. Chem.* **2005**, *48*, 5932–5941. (e) Jones, S. M.; Urch, J. E.; Brun, R.; Harwood, J. L.; Berry, C.; Gilbert, I. H. Analogues of Thiolactomycin as Potential Anti-malarial and Anti-trypansomal Agents. *Bioorg. Med. Chem.* **2004**, *12*, 683–692. (f) Senior, S. J.; Illarionov, P. A.; Gurucha, S. S.; Campbell, I. B.; Schaeffer, M. L.; Minnikin, D. E.; Besra, G. S. Acetylene-Based Analogues of Thiolactomycin, Active Against Mycobacterium Tuberculosis MtFabH Fatty Acid Condensing Enzyme. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 373–376. (g) Bhowruth, V.; Brown, A. K.; Senior, S. J.; Snaith, J. S.; Besra, G. S. Synthesis and Biological Evaluation of a C5-Biphenyl Thiolactomycin Library. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5643–5646. (h) Schaetzer, J. H.; Pitterna, T.; Lu, L.; Wu, Y.; Renold, P.; Perruccio, F.; Cassayre, J. Y.; Mueglebach, M. Spiroheterocyclic Derivatives as Pesticides and Their Preparation and Use. *PCT Int. Appl. WO 2010135914 A1* 2010-12-02.

(7) (a) Guo, X.; Hu, W. Novel Multicomponent Reactions via Trapping of Protic Onium Ylides with Electrophiles. *Acc. Chem. Res.* **2013**, *46*, 2427–2440. (b) Hu, W.; Xu, X.; Zhou, J.; Liu, W.-J.; Huang, H.; Hu, J.; Yang, L.; Gong, L.-Z. Cooperative Catalysis with Chiral Brønsted Acid-Rh₂(OAc)₄: Highly Enantioselective Three-

Component Reactions of Diazo Compounds with Alcohols and Imines. *J. Am. Chem. Soc.* **2008**, *130*, 7782–7783. (c) Guan, X.-Y.; Yang, L.-P.; Hu, W. Cooperative Catalysis in Multicomponent Reactions: Highly Enantioselective Synthesis of γ -Hydroxyketones with a Quaternary Carbon Stereocenter. *Angew. Chem., Int. Ed.* **2010**, *49*, 2190–2192. (d) Jiang, J.; Xu, H.-D.; Xi, J.-B.; Ren, B.-Y.; Lv, F.-P.; Guo, X.; Jiang, L.-Q.; Zhang, Z.-Y.; Hu, W.-H. Diastereoselectively Switchable Enantioselective Trapping of Carbamate Ammonium Ylides with Imines. *J. Am. Chem. Soc.* **2011**, *133*, 8428–8431. (e) Zhang, D.; Zhou, J.; Xia, F.; Kang, Z.; Hu, W. Bond Cleavage, Fragment Modification and Reassembly in Enantioselective Three-Component Reactions. *Nat. Commun.* **2015**, *6*, 5801. (f) Xiao, G.; Ma, C.; Xing, D.; Hu, W. Enantioselective Synthesis of α -Mercapto- β -amino Esters via Rh(II)/Chiral Phosphoric Acid-Cocatalyzed Three-Component Reaction of Diazo Compounds, Thiols, and Imines. *Org. Lett.* **2016**, *18*, 6086–6089. (g) Xiao, G.; Chen, T.; Ma, C.; Xing, D.; Hu, W. Rh(II)/Chiral Phosphoric Acid-Cocatalyzed Enantioselective Synthesis of Spirooxindole-Fused Thiaindanes. *Org. Lett.* **2018**, *20*, 4531–4535. (h) Qiu, H.; Li, M.; Jiang, L.-Q.; Lv, F.-P.; Zan, L.; Zhai, C.-W.; Doyle, M. P.; Hu, W.-H. Highly Enantioselective Trapping of Zwitterionic Intermediates by Imines. *Nat. Chem.* **2012**, *4*, 733–738. (i) Zhang, D.; Qiu, H.; Jiang, L.; Lv, F.; Ma, C.; Hu, W. Enantioselective Palladium(II) Phosphate Catalyzed Three-Component Reactions of Pyrrole, Diazoesters, and Imines. *Angew. Chem., Int. Ed.* **2013**, *52*, 13356–13360. (j) Jia, S.; Xing, D.; Zhang, D.; Hu, W. Asymmetric Functionalization of Aromatic C-H Bonds by Electrophilic Trapping of Metal-Carbene-Induced Zwitterionic Intermediates. *Angew. Chem., Int. Ed.* **2014**, *53*, 13098–13101.

(8) (a) Hunter, A. C.; Chinthapally, K.; Bain, A. I.; Stevens, J. C.; Sharma, I. Rhodium/Gold Dual Catalysis in Carbene sp^2 C-H Functionalization/Conia-ene Cascade for the Stereoselective Synthesis of Diverse Spirocarbocycles. *Adv. Synth. Catal.* **2019**, *361*, 2951–2958. (b) Bhat, A. H.; Alavi, S.; Grover, H. K. Tandem Carbenoid C-H Functionalization/Conia-ene Cyclization of *N*-Propargyl Indoles Generates Pyrroloindoles under Cooperative Rh(II)/Zn(II) Catalysis. *Org. Lett.* **2020**, *22*, 224–229.

(9) (a) Xia, Y.; Wang, J. Transition-Metal-Catalyzed Cross-Coupling with Ketones or Aldehydes via *N*-Tosylhydrazones. *J. Am. Chem. Soc.* **2020**, *142*, 10592–10605. (b) Xia, Y.; Qiu, D.; Wang, J. Transition-Metal-Catalyzed Cross-Couplings through Carbene Migratory Insertion. *Chem. Rev.* **2017**, *117*, 13810–13889. (c) Fulton, J. R.; Aggarwal, V. K.; de Vicente, J. The Use of Tosylhydrazone Salts as a Safe Alternative for Handling Diazo Compounds and Their Applications in Organic Synthesis. *Eur. J. Org. Chem.* **2005**, *2005*, 1479–1492. (d) Arunprasath, D.; Devi Bala, B.; Sekar, G. Stereoselective Construction of α -Tetralone-Fused Spirooxindoles via Pd-Catalyzed Domino Carbene Migratory Insertion/Conjugate Addition Sequence. *Org. Lett.* **2017**, *19*, 5280–5283. (e) Gutman, E. S.; Arredondo, V.; Van Vranken, D. L. Cyclization of η^3 -Benzylpalladium Intermediates Derived from Carbene Insertion. *Org. Lett.* **2014**, *16*, 5498–5501. (f) Khanna, A.; Maung, C.; Johnson, K. R.; Luong, T. T.; Van Vranken, D. L. Carbenylative Amination with *N*-Tosylhydrazones. *Org. Lett.* **2012**, *14*, 3233–3235. (g) Huang, Y.; Li, X.; Wang, X.; Yu, Y.; Zheng, J.; Wu, W.; Jiang, H. Copper-Catalyzed Cyanothiolation to Incorporate a Sulfur-Substituted Quaternary Carbon Center. *Chem. Sci.* **2017**, *8*, 7047–7051.

(10) Ishitobi, K.; Muto, K.; Yamaguchi, J. Pd-Catalyzed Alkenyl Thioether Synthesis from Thioesters and *N*-Tosylhydrazones. *ACS Catal.* **2019**, *9*, 11685–11690.

(11) (a) Teng, Q.; Liu, Y.; Hu, J.; Chai, H.; Li, K.; Zhang, H. Straightforward Synthesis of Ferrocenyl-Based Thioether and Application as Chemosensor for Hg^{2+} ion. *Tetrahedron* **2016**, *72*, 3890–3897. (b) García-Carrillo, M. A.; Guzmán, A.; Díaz, E. Metal Free Coupling of Heteroaryl *N*-tosylhydrazones and Thiols: Efficient Synthesis of Sulfides. *Tetrahedron Lett.* **2017**, *58*, 1952–1956. (c) Ding, Q.; Cao, B.; Yuan, J.; Liu, X.; Peng, Y. Synthesis of Thioethers via Metal-Free Reductive Coupling of Tosylhydrazones with Thiols. *Org. Biomol. Chem.* **2011**, *9*, 748–751. (d) Xu, Y.; Lv, G.; Yan, K.; He, H.; Li, J.; Luo, Y.; Lai, R.; Hai, L.; Wu, Y. Blue Light-

promoted Carbene Transfer Reactions of Tosylhydrazones. *Chem. - Asian J.* **2020**, *15*, 1945–1947.

(12) Wang, S.; Cheng, B.-Y.; Srsen, M.; König, B. Umpolung Difunctionalization of Carbonyls via Visible-light Photoredox Catalytic Radical-Carbanion Relay. *J. Am. Chem. Soc.* **2020**, *142*, 7524–7531.

(13) Old, D. W.; Wolfe, J. P.; Buchwald, S. L. A Highly Active Catalyst for Palladium-Catalyzed Cross-Coupling Reactions: Room-Temperature Suzuki Couplings and Amination of Unactivated Aryl Chlorides. *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723.