


# RASPA 1.0: Molecular Software Package for Adsorption and Diffusion in (Flexible) Nanoporous Materials



version 0.99-481

David Dubbeldam<sup>1</sup>

Van 't Hoff Institute, University of Amsterdam,  
Science Park 904, 1098XH, Amsterdam, The Netherlands

Sofia Calero<sup>2</sup>

Department of Physical, Chemical and Natural Systems, University Pablo de Olavide,  
Sevilla 41013, Spain

Donald E. Ellis<sup>3</sup>

Department of Physics and Astronomy, Northwestern University,  
2145 Sheridan Road, Evanston IL 60208 USA

Randall Q. Snurr<sup>4</sup>,

Chemical and Biological Engineering Department, Northwestern University,  
2145 Sheridan Road, Evanston IL 60208, USA

December 26, 2011

<sup>1</sup>email: D.Dubbeldam@uva.nl

<sup>2</sup>email: scalero@upo.es

<sup>3</sup>email: don-ellis@northwestern.edu

<sup>4</sup>email: snurr@northwestern.edu



# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Design philosophy . . . . .	5
1.2	Units and conventions . . . . .	6
1.3	Compiling and installing RASPA . . . . .	6
1.3.1	Requirements . . . . .	6
1.3.2	unpacking/getting RASPA . . . . .	7
1.3.3	installing RASPA . . . . .	7
1.3.4	compiling RASPA . . . . .	7
1.3.5	Running RASPA . . . . .	8
1.4	Output from RASPA . . . . .	10
1.5	Submitting batch-jobs on a cluster-system . . . . .	11
1.6	Future plans . . . . .	11
<b>2</b>	<b>Format of the Input Files</b>	<b>13</b>
2.1	Introduction . . . . .	13
2.2	Simulation input . . . . .	14
2.3	Force field . . . . .	36
2.3.1	Force fields . . . . .	36
2.3.2	'pseudo_atoms.def' . . . . .	36
2.3.3	'force_field_mixing_rules.def' . . . . .	36
2.3.4	'force_field.def' . . . . .	38
2.4	Molecules . . . . .	39
2.4.1	Rigid molecule . . . . .	39
2.4.2	Flexible molecule . . . . .	39
2.4.3	Rigid/Flexible molecule . . . . .	40
2.4.4	Chiral molecules . . . . .	41
2.5	Framework . . . . .	42
2.5.1	Asymmetric unit cell . . . . .	42
2.5.2	Fractional occupancies in zeolites . . . . .	44
2.5.3	Format of the framework atoms . . . . .	46
2.5.4	Typing the atoms of the framework . . . . .	47
2.6	Using CIF-files . . . . .	51
2.6.1	Definition of CIF-files . . . . .	51
	Bibliography . . . . .	64
<b>3</b>	<b>Potentials</b>	<b>65</b>
3.1	Functional forms of force fields . . . . .	65
3.2	Bonded potentials diagonal terms . . . . .	66
3.2.1	Bond-stretching potentials . . . . .	66
3.2.2	Urey-Bradley potentials . . . . .	67

3.2.3	Bending potential . . . . .	69
3.2.4	Wilson inversion-bend potential . . . . .	70
3.2.5	Torsion potential . . . . .	72
3.2.6	Improper torsion potential . . . . .	74
3.3	Non-bonded potentials . . . . .	77
3.3.1	Van der Waals potentials . . . . .	77
3.3.2	Tail corrections . . . . .	80
3.3.3	Electrostatics . . . . .	81
3.4	Bonded potentials cross terms . . . . .	82
3.4.1	Bond-bond potential . . . . .	82
3.4.2	Bond-bend potential . . . . .	82
3.4.3	Bend-bend potential . . . . .	83
3.4.4	Bond-torsion potential . . . . .	83
3.4.5	Bend-torsion potential . . . . .	83
3.5	Polarization . . . . .	84
3.5.1	Introduction . . . . .	84
3.5.2	Computational scheme . . . . .	85
	Bibliography . . . . .	86
<b>4</b>	<b>Examples</b>	<b>87</b>
4.1	Introduction . . . . .	87
4.2	Basic examples . . . . .	88
4.3	Non-basic examples . . . . .	95
4.4	Advanced examples . . . . .	104
4.5	Auxiliary examples . . . . .	113
4.6	Validation examples . . . . .	118
<b>5</b>	<b>The source code</b>	<b>121</b>
5.1	Introduction . . . . .	121
5.2	Data types . . . . .	121
5.3	Datastructures . . . . .	122
5.4	Modifying . . . . .	129
5.4.1	Monte Carlo . . . . .	129
5.4.2	Molecular Dynamics . . . . .	131
5.5	Debugging . . . . .	134
5.5.1	Linux . . . . .	134
5.5.2	Mac OSX . . . . .	134
5.6	RASPA as a project in Xcode 4 . . . . .	135
5.6.1	Getting started . . . . .	135
5.6.2	Fine tuning . . . . .	138
5.6.3	Using 'git' (version control) . . . . .	140
5.7	Using 'git' . . . . .	140
5.7.1	Installing git . . . . .	140
5.7.2	. . . . .	141
5.8	Coding conventions . . . . .	141
5.9	Optimization tips . . . . .	142
<b>6</b>	<b>Troubleshooting</b>	<b>143</b>

# Chapter 1

## Introduction

### 1.1 Design philosophy

This software is a general purpose classical simulation package. It has been developed at Northwestern University (Evanston, USA) during 2006, 2007 and 2008 in active collaboration with University Pablo de Olavide (Seville, Spain). It can be used for the simulation of molecules in gases, fluids, zeolites, aluminosilicates, metal-organic frameworks, carbon nanotubes and external fields. It has incorporated most of the capabilities of DLPOLY, BIGMAC, BIGGERMAC, TINKER, MUSIC, GULP, and COTA.

Programs can be written in various ways, but often it is true that the fastest codes are probably the hardest to read, while programs strictly based on readability lacks efficiency. RASPA is based on the following ideas:

- Correctness and accuracy

For all the techniques and algorithms available in RASPA we have implemented the 'best' ones available in literature. For example, RASPA uses Configurational-Bias Monte-Carlo, it uses the Ewald summation for electrostatics, molecular dynamics is based on 'symplectic' integrators, all Monte-Carlo moves obey detailed balance etc.

- Functional design

Looking at the source, you will notice that there are not a lot of files. The program is splitted up according to its function: 'grid.c' contains the code to make and use a grid of a framework, 'ewald.c' handles all the electrostatic, 'mc\_moves.c' contains all the moves to be used in Monte-Carlo, 'potentials.c' contains all the VDW potentials etc.

- Input made easy

The requirements for the input files is kept as minimal as possible. Only for more advanced options extra commands in the input file are needed. Also the format of the input is straightforward. Default settings are usually the best ones. Fugacity coefficients and excess adsorption are automatically computed.

- Integrated simulation environment

The code is built up of many functions and routines which can be easily combined to do what you want. Molecular dynamics can be used in Monte Carlo and visa versa. Extension and modification of the code is relatively straightforward.

RASPA used three 'types' or 'groups' for the particles: 1) Framework atoms, 2) Adsorbates, and 3) Cations. The advantage is that all the energies are splitted and the interactions can be examined (also the energies are splitted in the Ewald Fourier part). Another example is when using thermostats in e.g. LTA5A where a different thermostat operates in the framework atoms, the adsorbates, and the cations. These all move at different length- and time scales. Note that it is not possible to exchange types during Identity-change moves (if defined they are ignored).

## 1.2 Units and conventions

- The standard units in RASPA from which all other units are derived are:

quantity	symbol	unit	value
length	$l$	Ångstrom	$10^{-10}$ m
temperature	$T$	Kelvin	K
mass	$m$	atomic mass	$1.6605402 \times 10^{-27}$ kg
time	$t$	pico seconds	$10^{-12}$ s
charge	$q$	atomic charge	$1.60217733 \times 10^{-19}$ C/particle

Some examples of derived units:

quantity	symbol	units	conversion value
energy	$U$	$J = \text{mass} \times \text{length}^2/\text{time}^2$	$1.66054 \times 10^{-23}$ (=10 J/mol)
pressure	$p$	$\text{Pa} = \text{mass}/(\text{length} \times \text{time}^2)$	$1.66054 \times 10^7$
diffusion constant	$D$	$D = \text{length}^2/\text{time}$	$1 \times 10^{-8}$
force	$f$	$f = \text{length}/\text{time}^2$	$1.66054 \times 10^{-13}$
...	...	...	...

A pressure input of 10 Pascal in the input file, is converted to 'internal units' by dividing by  $1.66054 \times 10^7$ . In the output any internal pressure is printed, multiplied by  $1.66054 \times 10^7$ . It is not necessary to convert units besides input and output, with a few exceptions. One of them is the Coulombic conversion factor

$$\frac{q_i q_j}{4\pi\epsilon_0} = \frac{\text{charge}^2}{4\pi \times \text{electric constant} \times \text{length} \times \text{energy}} = 138935.4834964017 \quad (1.1)$$

with the electric constant as  $8.8541878176 \times 10^{-12}$  in units of  $\text{C}^2/(\text{N.m}^2)$ . This factor is needed to convert the electrostatic energy to the internal units at every evaluation.

The Boltzmann's constant  $k_B$  is

$$k_B = \text{Boltzmann constant}/\text{energy} = 0.8314464919 \quad (1.2)$$

with the Boltzmann constant as  $1.380650324 \times 10^{-23}$  in units of  $\text{J/K}$ , and  $k_B = 0.8314464919$  in internal units.

- Numbering is based on the C-convention, i.e. starting from zero.
- Files in the current directory always have preference.  
Sometimes one would like to try various parameters for force field fitting for example. In order to avoid making a lot of directories for each force field it is more convenient to have the 'pseudo\_atoms.def', 'force\_field\_mixing\_rule.def' and 'force\_field.def' files in the *current* directory.

## 1.3 Compiling and installing RASPA

### 1.3.1 Requirements

RASPA needs a C compiler, like 'gcc' or intel's 'icc' compilers, and the libraries 'blas' and 'lapack'.

### 1.3.2 unpacking/getting RASPA

The 'tar' command can be used to extract the files from the archive

```
tar -zxvf Raspa-x.xx.tar.gz
```

To create a compressed archive use

```
tar -zcvf Raspa-x.xx.tar.gz raspa-x.xx/
```

executing from the directory where the source code of RASPA is in.

Alternatively, the source code of RASPA is available via 'git'. Working with 'git' and a remote repository means that you will have to distinguish between two locations of the code:

1. The repository (visible to everyone)
2. your local copy (only visible to you)

To check-out the code for the first time do:

```
git clone ssh://raspa@itsaio.science.uva.nl/~raspa/RASPA.git
```

and to check-out the documentation for the first time do:

```
git clone ssh://raspa@itsaio.science.uva.nl/~raspa/RASPA-Docs.git
```

### 1.3.3 installing RASPA

The *RASPA\_DIR* environment variable should be set to where you would like to install RASPA. A common way of defining it is using the bash-shell

```
export RASPA_DIR=${HOME}/Research/simulations/
```

or

```
setenv RASPA_DIR "${HOME}/Research/simulations/"
```

for 'csh' and 'tcsh' shells. It is possible to add this line to ".bashrc", "/etc/bashrc", "/etc/profile" etc, depending on the unix-version and shell version to automatically have the environment variable set at login.

Note that the source-code of RASPA is kept separate from the installation data. RASPA needs the environment variable to locate various files it needs, e.g. molecule definitions, framework definitions, force and field definitions. It looks for these files relative to the *RASPA\_DIR* directory.

Before installing RASPA with

```
make install
```

from the top-directory, the code needs to be compiled.

### 1.3.4 compiling RASPA

RASPA uses the standard 'configure' utilities (autoconf, automake, libtool, and make). The steps to install from scratch, i.e. after a 'make distclean' are

```
rm -rf autom4te.cache  
aclocal  
automake --add-missing  
autoconf  
.configure --prefix=${RASPA_DIR}  
make
```

where ‘\${RASPA\_DIR}’ is the directory you would like to install RASPA, and the commands are executed in the top directory.

You can use the ‘CFLAGS’ environment variable to set compiler options and ‘CC’ to set the compiler. For example, for a gcc compiler one could use

```
export CFLAGS="-Wall -O4 -ffast-math"
export CC="gcc"
```

### 1.3.5 Running RASPA

Running RASPA is based on two files:

- A ‘run’ file to execute the program  
an example file is:

```
#!/bin/sh -f
export RASPA_DIR=${HOME}/Research/simulations/
$RASPA_DIR/bin/simulate
```

This type of file is known as a ‘shell script’. RASPA needs the variable ‘RASPA\_DIR’ to be set in order to look up the molecules, frameworks, etc. The script sets the variable and runs RASPA. RASPA can then be run from any directory you would like.

- An ‘input’-file describing the type of simulation and the parameters  
In the same directory as the ‘run’-file, there needs to be a file called ‘simulation.input’. An example file is:

```
SimulationType          MonteCarlo
NumberOfCycles          100000
NumberOfInitializationCycles 10000
PrintEvery               1000

Box 0
BoxLengths 30 30 30
ExternalTemperature 300.0

component 0 methane
    TranslationProbability      1.0
    CreateNumberOfMolecules     100
```

This tells RASPA to run a Monte-Carlo simulation of 100 methane molecules in a  $30 \times 30 \times 30$  Å cubic box (with 90° angles) at 300 Kelvin. It will start with 10000 cycles to equilibrate the system and will use 100000 cycles to obtain thermodynamic properties of interest. Every 1000 cycles a status-report is printed to the output. The Monte-Carlo program will use only the ‘translation move’ where a particle is given a random translation and the move is accepted or rejected based on the energy difference.

In order to run it on a cluster using a queueing system one needs an additional file ‘bsub.job’ (arbitrary name), for ‘gridengine’

```
#!/bin/csh
# Serial sample script for Grid Engine
# Replace items enclosed by {}
```

```

## -S /bin/csh
## -N MinimizationIRMOF-1
## -V
## -cwd
echo $PBS_JOBID > jobid
setenv RASPA_DIR ${HOME}/RASPA/simulations/
$RASPA_DIR/bin/simulate

```

or for a 'torque' system

```

#!/bin/bash
#PBS -N Test
#PBS -o pbs.out
#PBS -e pbs.err
#PBS -r n
#PBS -V
#PBS -mba
cd $PBS_O_WORKDIR
echo $PBS_JOBID > jobid
export RASPA_DIR=${HOME}/RASPA/simulations
${RASPA_DIR}/bin/simulate

```

The job can be submitted using 'qsub < bsub.job'.

**University of Amsterdam** has a rocks-linux cluster (based on centos 5.6). You can log in to the headnode

```
ssh username@carbon.science.uva.nl
```

Off-campus access to 'carbon' is provided using 'vpn'. The queuing system is based on 'gridengine'. The carbon-cluster has a filesystem based on ZFS with automatic compression and snapshots made for the last 3 days. These 'backups' are accessible in your home-account, in the directory `~/.zfs`.

**SARA** has a rocks-linux cluster (based on redhat). You can log in to the headnode

```
ssh username@lisa.sara.nl
```

'lisa' is accessible from everywhere. The queuing system is based on 'torque'.

**Georgia Tech** has a cluster of linux redhat nodes called 'pace'. You can log in to the headnode

```
ssh username@neojoey.pace.gatech.edu
```

Off-campus access to 'pace' is provided using 'vpn' or by first hopping to

```
ssh username@acme.gatech.edu
```

A serial job can be submitted using 'qsub < bsub.job' and

```

#!/bin/bash
#PBS -N job-name
#PBS -l nodes=1:ppn=1
#PBS -l walltime=120:00:00
#PBS -m n
#PBS -q joe-penguin
cd $PBS_O_WORKDIR
export RASPA_DIR=${HOME}/RASPA/simulations/
$RASPA_DIR/bin/simulate $1

```

The 'pace'-cluster does not really have a limit on the amount of time that you can run. In the example script we ask for 120 hours of run-time. That means the jobs will be killed after 120 hours so make sure the walltime is always higher than what you actually need. However, the downside of specifying a long walltime is that you will have to wait longer in the queue before your job starts. Note that with the 'BinaryRestartFile' options you can restart RASPA and continue where it left off.

The 'pace'-userguide can be found here:

<http://share-it.gatech.edu/oit/pace/user-guide>

and the status of the cluster can be viewed graphically

<http://neospace.pace.gatech.edu/ganglia/>

The compilers on 'pace' are a bit old, RASPA has been compiled using "gcc version 4.1.2 20080704 (Red Hat 4.1.2-48)" with the following options

```

#!/bin/sh
rm config.cache
export CFLAGS="-Wall -march=opteron -ffast-math -O3 -funroll-loops -ftree-vectorize
             -msse3 -m64"
export CC="gcc"
./configure --prefix=${HOME}/RASPA/simulations

```

Note the code using this version of gcc does not compile with '-ftree-loop-linear' which is known to cause incorrect code in rare circumstances. The 'pace' nodes are AMD opteron, and the intel compiler does not optimize well for an AMD-cpu.

## 1.4 Output from RASPA

RASPA generates output from the simulation. Some data is just information on the status, while other data are written because you specifically asked the program to compute it for you. The output is written to be used with other programs like:

- gnuplot
- VTK
- VMD

The main output is written to the directory 'Output/System[0]/', 'Output/System[1]/', ... for each of the simulated systems. Usually one simulates only a single system. However, the Gibbs ensemble requires 2 system, one for vapor phase and one for the liquid phase, while  $n$  systems are used by the (hyper-)parallel-tempering technique(s).

## 1.5 Submitting batch-jobs on a cluster-system

In the directory 'scripts' several perl job-scripts can be found:

- **script\_adsorption.pl**  
A script to compute single component isotherms (many pressures) at several temperatures for many frameworks and adsorbates.
- **script\_adsorption\_mixture.pl**  
A script to compute mixture isotherms (many pressures) at several temperatures for many frameworks and adsorbates.
- **script\_adsorption\_get\_results.pl**  
A script to get the results back from single component isotherms.
- **script\_adsorption\_mixture\_get\_results.pl**  
A script to get the results back from mixture isotherms.
- **script\_md.pl**  
A script to compute diffusion using molecular dynamics for many adsorbates in many systems at many tempereratures.
- **script\_md\_mixture.pl**  
A script to compute mixture diffusion using molecular dynamics for many adsorbates in many systems at many tempereratures.
- **script\_idealgas\_rosenbluth.pl\_md\_mixture.pl**  
A script to compute Rosenbluth weights for many adsorbates in an empty box at many temperatures.
- **script\_idealgas\_single\_molecule.pl\_md\_mixture.pl**  
A script to compute average energies for many ideal-gas (single molecule in an empty box) adsorbates at many temperatures.

## 1.6 Future plans

In future versions we plan to implement the following extensions

- Speed optimization (parallel implementations usign MPI)
- Symplectic NosePointCare integrators for NVT, NPT, and NPT-PR
- More MC moves, like the 'reptation' move
- Verlet lists/cell lists
- Rigid/quaternion minimization
- Symmetry/spacegroup for force/hessian
- Finish all inverse space groups
- Find space group(s)
- Saddle point search
- VDW using Ewald
- Histogram-reweighting (done run-time)

- Ewald 2D
- Analytical Hessian, Born term, and cross terms for all potentials
- Add clays, carbon-nanotubes, and metal-oxides structures
- Lattice Boltzmann
- Lattice gases
- Kinetic Monte Carlo
- Graphical User Interface (Cocoa)

# Chapter 2

## Format of the Input Files

### 2.1 Introduction

In order to run a simulation you need several input-files:

- **'simulation.input'**

This file contains the information on the type of simulation, the amount of steps, the framework unit cell, number of unit cells in each directions, the used molecules, the type of used Monte-Carlo moves etc.

- **'structure-name.cif'**

If a framework (e.g. a zeolite or MOF) is used, then the definition of the structure needs to be provided. CIF-files are supported and the default input. The name of the file should be equal to the one provided in **'simulation.input'**, e.g. IRMOF-1.cif if `Frameworkname IRMOF-1` is listed in **'simulation.input'**.

- **'pseudo\_atoms.def'**

The **'pseudo\_atoms.def'** file list all the information on used pseudo-atoms, e.g. charge, mass. Usually a pseudo-atom is an atom, but there are exceptions like united atoms (where CH3 is lumped into one unit) and off-atom sites in Tip5p water that represent oxygen lone pairs. Because in CIF-files for frameworks you can provide also information on atoms, there is no need to list framework atoms here if a CIF-file is used. On reading the CIF-file these defined atoms are added to the pseudo-atoms. If also provided in the **'pseudo\_atoms.def'** then the definition in the **'pseudo\_atoms.def'** file has priority.

- **'force\_field\_mxing\_rules.def','force\_field.def'**

The force field defined on the pseudo-atoms in **'pseudo\_atoms.def'**. These files list the Van der Waals potential types, the parameters, whether to use tail-corrections, whether to shift to zero at the cutoff, and the type of mixing rule. Force fields in literature are usually published in two forms: 1) a list of potentials parameters per atom and a mixing rule, or 2) pairs of atoms and parameters. The first option corresponds to the file **'force\_field\_mxing\_rules.def'** and the latter option to the file **'force\_field.def'**. You can use both at the same time, where **'force\_field.def'** has precedence over **'force\_field\_mxing\_rules.def'**.

- **'molecule-name.def'**

The definition of the used molecules. The name of the file should be equal to the one provided in **'simulation.input'**, e.g. propane.def if `MoleculeName propane` is listed in **'simulation.input'**.

- **'framework.def'**

Used for a flexible framework to define all the bonds, bends, torsions, core-shells, etc.

The format of these files will be described in the remaining sections. Chapter 4 provides lots of examples to see everything in action. In addition to the input-files you will need either a 'run' file that is executable, or a queuing-script to submit the job to the queue (see 1.3.5).

## 2.2 Simulation input

Leading spaces and comments at the end of each line are omitted. Empty lines are skipped, and case is not important except in file names (i.e. framework and molecule names).

### Simulation types

- **SimulationType MonteCarlo**  
Starts the Monte Carlo part of RASPA. The particular ensemble is not specified but implicitly deduced from the specified Monte Carlo moves. Note that a MD-move can be used for hybrid MC/MD.
- **SimulationType MolecularDynamics**  
Starts the Molecular Dynamics part of RASPA. The ensemble is explicitly specified.
- **SimulationType Spectra**  
Starts the computation of the vibrational analysis. Possible options include infra red spectrum at zero Kelvin, powder diffraction, and mode analysis.
- **SimulationType Minimization**  
Starts the minimization routine. It produces configurations and crystal structures at zero Kelvin.
- **SimulationType Visualization**  
Output VTK-files for snapshots and crystal structures, including energy surface pictures.
- **SimulationType BarrierCrossing**  
Routine for the dynamical correction of dynamically corrected Transition State Theory.
- **SimulationType Numerical**  
Computes all the forces numerically from the energy and compares them to the analytical expressions. Also the strain-derivative tensor (related to the stress tensor), and the second derivative of the energy with respect to strain, as well as the Hessian matrix can be checked.
- **SimulationType MakeGrid**  
Creates pre-tabulated energy-grids for use in rigid frameworks.

### Simulation duration

- **NumberOfCycles [int]**  
The number of cycles for the production run. For Monte Carlo a cycle consists of  $N$  steps, where  $N$  is the amount of molecules with a minimum of 20 steps. This means that on average during each cycle on each molecule a Monte Carlo move has been attempted (either successful or unsuccessful). For MD the number of cycles is simply the amount of integration steps.
- **NumberOfInitializationCycles [int]**  
The number of cycles used to initialize the system using Monte Carlo. This can be used for both Monte Carlo as well as Molecular Dynamics to quickly equilibrate the positions of the atoms in the system.
- **NumberOfEquilibrationCycles [int]**  
The number of Molecular Dynamics steps to equilibrate the velocities in the systems. After this equilibration the production run is started.

## Restart and crash-recovery

- **RestartFile** [yes|no]  
Reads the positions, velocities, and force from the directory 'RestartInitial'. Any creation of molecules in the 'simulation.input' file will be in addition and after this first read from file. This is useful to load initial positions of cations for example, and after that create adsorbates. The restart file is written at 'PrintEvery' intervals.
- **ContinueAfterCrash** [yes|no]  
Write a binary file containing the complete status of the program. The file name is 'binary\_restart.dat' and is located in the directory 'CrashRestart'. With this option to 'yes' the presence of this file will result in continuation from the point where the program was at the moment of outputting this file. The file can be quite big (several hundreds of megabytes) and will be outputted every 'WriteBinaryRestartFileEvery' cycles.
- **WriteBinaryRestartFileEvery** [int]  
The output frequency (i.e. every [int] cycles) of the crash-recovery file.

## Printing options

- **PrintEvery** [int]  
Prints the loadings (when a framework is present) and energies every [int] cycles. For MD information like energy conservation and stress are printed.
- **PrintPropertiesEvery** [int]  
Output running averages of many properties (i.e. Henry coefficients and elastic constants).
- **PrintForcefieldToOutput** [yes—no]  
Prints the force field information to the output-file. Default: yes.
- **PrintPseudoAtomsToOutput** [yes—no]  
Prints the pseudo-atom information to the output-file. Default: yes.
- **PrintMoleculeDefinitionToOutput** [yes—no]  
Prints the molecule definition information to the output-file. Default: yes.

## Force field definitions

- **ChargeFromChargeEquilibration** [yes|no]  
Compute the charges of the framework using the 'charge-equilibration'-method.
- **SymmetrizeFrameworkCharges** [yes|no]  
All charges of the framework are made equivalent for equivalent framework atoms. Using regular charge-equilibration the charges are different for symmetrically equivalent framework atoms, and this option restores the symmetry.
- **ForceField** [string]  
Reads in the force field [string], first the file 'pseudo\_atoms.def' is read, then 'force\_field\_mixing\_rules.def' and finally 'force\_field.def'. The latter overwrites general settings for interactions based on mixing rules with specific ones for individual interactions.

Note that if any of these files are in the working directory then these will be read and used instead of the ones in '\${RASPA\_DIR}/simulations/share/raspa/forcefield/[string]'.

- **CutOffVDW** [real]

The cutoff of the Van der Waals potentials. Interactions longer than this distance are omitted from the energy and force computations. The potential can either be shifted to zero at the cutoff, or interactions can just be neglected after the cut off, or the remainder of the potential energy can be approximated using tail corrections. This is specified in the force field files and can be specified globally or for each interaction individually.

- **CutOffVDWSwitch** [real]

The distance at which VDW switching will start. The smoothing will make sure the value and derivatives are zero at the cutoff. The default: 0.9 times the CutOff.

- **CutOffChargeCharge** [real]

The cutoff of the charge-charge potential. The potential is truncated at the cutoff and only shifted when 'ChargeMethod CoulombShifted' or 'ChargeMethod CoulombSmoothed' is used. No tail-corrections are (or can be) applied. The only way to include the long-range part is to use 'ChargeMethod Ewald'. The parameter is also used in combination with the Ewald precision to compute the number of wave vectors and Ewald parameter  $\alpha$ . For the Ewald summation using rather large unit cells, a charge-charge cutoff of about half the smallest box-length would be advisable in order to avoid the use of an excessive amount of wave-vectors in Fourier space. For non-Ewald methods the cutoff should be as large as possible (greater than about 30 Å).

- **CutOffChargeChargeSwitch** [real]

The distance at which charge-charge switching will start. The smoothing will make sure the value and derivatives are zero at the cutoff. The default: 0.65 times the CutOff.

- **CutOffChargeBondDipole** [real]

The cutoff of the charge-bonddipole potential.

- **CutOffChargeBondDipoleSwitch** [real]

The distance at which charge-bonddipole switching will start. The smoothing will make sure the value and derivatives are zero at the cutoff. The default: 0.70 times the CutOff.

- **CutOffBondDipoleBondDipole** [real]

The cutoff of the bonddipole-bonddipole potential.

- **CutOffBondDipoleBondDipoleSwitch** [real]

The distance at which bonddipole-bonddipole switching will start. The smoothing will make sure the value and derivatives are zero at the cutoff. The default: 0.75 times the CutOff.

- **OmitAdsorbateAdsorbateVDWInteractions** [yes|no]

Omits the Van der Waals interactions between adsorbates.

- **OmitAdsorbateAdsorbateCoulombInteractions** [yes|no]

Omits the Coulombic (i.e. Ewald) interactions between adsorbates.

- **OmitInterMolecularInteractions** [yes|no]

Omits the interactions between all molecules (only interactions with the framework). This also works with the Ewald summation on. The options implies the setting of both

  - **OmitAdsorbateAdsorbateVDWInteractions** [yes|no]

  - **OmitAdsorbateAdsorbateCoulombInteractions** [yes|no]

- **InternalFrameworkLennardJonesInteractions** [yes|no]

Compute the Van der Waals interaction of the flexible framework. The Demontis flexible model for silicalite is defined with only bond, bend, and torsion for example. One can use this option and also use 'Charge None'.

- RemoveBondNeighboursFromLongRangeInteraction [yes|no]
- RemoveBendNeighboursFromLongRangeInteraction [yes|no]  
RemoveTorsionNeighboursFromLongRangeInteraction [yes|no]  
After construction of the connectivity table all interactions are removed from Van der Waals and charge interactions that are defined as 1-2 (i.e. bonds), 1-3 (i.e. bends, Urey-Bradley) and 1-4 (i.e. torsion, inversion-bend) respectively.
- Remove12NeighboursFromChargeChargeInteraction [yes|no]  
Remove13NeighboursFromChargeChargeInteraction [yes|no]  
Remove14NeighboursFromChargeChargeInteraction [yes|no]  
Remove all 1-2, 1-3, and/or 1-4 interactions within the framework from the long-range charge-charge interaction within the flexible framework respectively.
- Remove12NeighboursFromChargeBondDipoleInteraction [yes|no]  
Remove13NeighboursFromChargeBondDipoleInteraction [yes|no]  
Remove14NeighboursFromChargeBondDipoleInteraction [yes|no]  
Remove all 1-2, 1-3, and/or 1-4 interactions within the framework from the long-range charge-bond dipole interaction within the flexible framework respectively.
- Remove12NeighboursFromBondDipoleBondDipoleInteraction [yes|no]  
Remove13NeighboursFromBondDipoleBondDipoleInteraction [yes|no]  
Remove14NeighboursFromBondDipoleBondDipoleInteraction [yes|no]  
Remove all 1-2, 1-3, and/or 1-4 interactions within the framework from the long-range bond dipole-bond dipole interaction within the flexible framework respectively.
- ComputePolarization [yes|no]  
Turns on the computation of polarization.
- BackPolarization [yes|no] (not working yet)  
Turns on the self-consistent iteration. If "BackPolarization" is off the polarization algorithm just uses the first term in the expansion, e.g. the static electric field. If "BackPolarization" is on the algorithm iterates for "NumberOfBackPolarizationSteps" to take into account that the induced dipoles alter the electric field, which in turn alters the electric field etc.
- NumberOfBackPolarizationSteps [int]  
The number of self-consistent iterations to reach convergence.
- OmitIntraFrameworkPolarization [yes|no]  
Omits the polarization of framework atoms due to other framework atoms. If the Ewald summation is used, the intra framework contribution is exactly removed (i.e. the polarization due to the framework as well as *all* the periodic images of the framework atoms are removed). Note that  $\mathbf{E} \neq \mathbf{F}/q$  because the periodic images are included in the force but not in the electric field.
- OmitAdsorbateAdsorbatePolarization [yes|no]  
Omits the adsorbate polarization due to other adsorbate atoms.
- OmitCationCationPolarization [yes|no]  
Omits the cation polarization due to other cation atoms.

## Thermostat and barostat parameters

- ExternalTemperature [list-of-reals]  
The external temperature in Kelvin for each system. Because the system is in contact with this imaginary reservoir the average temperature of the system can be controlled. Default: 298K.

- ExternalPressure [list-of-reals]  
The external pressure in Pascal for each system. Because the system is in contact with this imaginary reservoir the average pressure of the system can be controlled.
- ThermostatChainLength [int]  
The length of the chain to thermostat the system. Default: 5.
- BarostatChainLength [int]  
The length of the chain to thermostat the volume and/or cell parameters. Default 5.
- NumberOfYoshidaSuzukiSteps [int]  
The number of Yoshida/Suzuki multiple timesteps.
- TimeScaleParameterThermostat [real]  
The time scale on which the system thermostat evolves. Default: 0.15 ps.
- TimeScaleParameterBarostat [real]  
The time scale on which the thermostat for the volume and/or cell parameters evolve. Default: 0.15 ps.

## Molecular dynamics parameters

- TimeStep [real]  
The time step in picoseconds for MD integration. Default value: 0.0005 ps (0.5 fs).
- Ensemble [list-of-NVE|NVT|NPT|NPH|NPTPR|NPHPR]  
Sets the ensemble as a list of NVE,NVT, NPT, NPH, NPTPR, or NPHPR for each system. If only a single ensemble is given, it is used for all systems. The given ensemble will be used for both initialization as well as the production run.
  - NVE  
The micro canonical ensemble, the number of particle  $N$ , the volume  $V$ , and the energy  $E$  are constant.
  - NVT  
The canonical ensemble, the number of particle  $N$ , the volume  $V$ , and the average temperature  $\langle P \rangle$  are constant. Instantaneous values for the temperature are fluctuating.
  - NPT  
The isobaric-isothermal ensemble, the number of particle  $N$ , the average pressure  $\langle P \rangle$ , and the average temperature  $\langle P \rangle$  are constant. Instantaneous values for the pressure and temperature are fluctuating.
  - NPH  
The isoenthalpic-isobaric ensemble, the number of particle  $N$ , the average pressure  $\langle P \rangle$ , and the enthalpy  $H$  are constant. Instantaneous values for the pressure and temperature are fluctuating.
  - NPTPR  
The isobaric-isothermal ensemble with a fully flexible cell (Parrinello-Rahman).
- InitEnsemble [list-of-NVE|NVT|NPT|NPH|NPTPR|NPHPR]  
Sets the ensemble as a list of NVE,NVT, NPH, NPTPR, or NPHPR for each system. If only a single ensemble is given, it is used for all systems. The given ensemble will be only used for the initialization run.
- RunEnsemble [list-of-NVE|NVT|NPT|NPH|NPTPR|NPHPR]  
Set the ensemble as a list of NVE,NVT, NPH, NPTPR, or NPHPR for each system. If only a single ensemble is given, it is used for all systems. The given ensemble will be only used for the production run.

- NPTPRCellType [list-of-Regular|Monoclinic|RegularUpperTriangle|MonoclinicUpperTriangle|Isotropic|Anisotropic]  
 The type of constraints on the cell-matrix **h**. Default: RegularUpperTriangle.
  - Regular  
 If the pressure tensor is asymmetric ( $P_{\alpha\beta} \neq P_{\beta\alpha}$ ) at a given instant of time, then there will be a net torque acting on the cell that will cause it to rotate. Cell rotations can be eliminated by using the symmetrized tensor  $P_{\alpha\beta} = (P_{\alpha\beta} + P_{\beta\alpha})/2$  in the equations of motion and setting the initial total angular momentum of the cell to zero. This approach is formally implemented by constraining the force on the cell  $\mathbf{g} = \mathbf{g}^T$ . All three angles  $\alpha, \beta, \gamma$  are allowed to change, as well as the box lengths **a**, **b**, **c**.
  - Monoclinic  
 All three box lengths **a**, **b**, **c** are allowed to vary, as well as cell angle  $\beta$ , but  $\alpha = \gamma = 90^\circ$ .
  - RegularUpperTriangle  
 Only the upper triangular part of the cell matrix is used to eliminate rotation of the box. All three angles  $\alpha, \beta, \gamma$  are allowed to change, as well as the box lengths **a**, **b**, **c**.
  - MonoclinicUpperTriangle  
 Only the upper triangular part of the cell matrix is used to eliminate rotation of the box. All three box lengths **a**, **b**, **c** are allowed to vary, as well as cell angle  $\beta$ , but  $\alpha = \gamma = 90^\circ$ .
  - Isotropic  
 All three box lengths **a** = **b** = **c** are allowed to vary isotropically, and the angles remain fixed  $\alpha = \beta = \gamma = 90^\circ$ .
  - Anisotropic  
 All three box lengths **a**, **b**, **c** are allowed to vary *independently*, but the angles remain fixed  $\alpha = \beta = \gamma = 90^\circ$ .

## Box parameters

- Box [int]  
 Set the current system to type 'Box [int]' (other option is 'Framework' when a framework is present).
- BoxLengths [real] [real] [real]  
 The cell dimensions of the box in Angstroms. Default: 25 25 25 Å.
- BoxAngles [real] [real] [real]  
 The cell angles of the box in degrees. Default: 90° 90° 90°.
- BoxMatrix [int]  
 [real] [real] [real]  
 [real] [real] [real]  
 [real] [real] [real]  
 Set the system [int] to type 'Box' (other option is 'Framework' when a framework is present). The  $3 \times 3$  cell matrix of system [int], given as three vectors (as columns). This is the most general form and any box can be specified in this way. Units of the vectors are Angstrom.

## Framework parameters

- Framework [int]  
 Set the current system to type 'Framework [int]' (other option is 'Box' when no framework is present). All other options listed in the section framework parameters refer to this system, so make sure this is before any other framework options.

- **FrameworkName** [string]  
Loads the framework with name [string]. Several frameworks can be read per system, which is useful for to study interpenetration of frameworks. Here the frameworks are allowed to move independently from each other.
- **HeliumVoidFraction** [real]  
The void fraction as measure by probing the structure with helium a room temperature. This quantity has to be obtained from a separate simulation and is essential to compute the *excess*-adsorption during the simulation.
- **InputFileType** [cif|xyz|mol|cssr]  
Sets the framework input type to CIF, XYZ, MOL, or CSSR-format.
- **UnitCells** [int] [int] [int]  
The number of unit cells in x,y, and z direction for the system. The full cell will contain the unit cells, and periodic boundary conditions will be applied on the box level (*not* on a unit cell level).
- **ShiftUnitCells** [real] [real] [real]  
Shift the fractional positions so that the center of a framework can be altered.
- **FlexibleFramework** [yes|no]  
Allow the current framework of the current system to be fully flexible. The name of the flexible model is provided using the 'FrameworkDefinitions [string]' input option.
- **FrameworkDefinitions** [string]  
The force field name [string] of the flexible framework. The file is read even when 'FlexibleFramework no' is specified (the reason is that framework bond-dipoles are defined using the 'framework.def' file).
- **ModifyFrameworkAtomConnectedTo** [atom-type-1] [atom-type-2] [atom-type-3] [atom-type-4]  
Modifies the atom-type-1 to atom-type-2, always if atom-type-3 and atom-type-4 are omitted, or only it is connected to atom-type-3 when atom-type-3 is specified, or only when it is connected to both atom-type-3 and atom-type-4 if both are specified.
- **ModifyFrameworkDimer** [atom-type-1] [atom-type-2] [atom-type-3] [atom-type-4]  
Modifies the connected atom-type-1 and atom-type-2 dimer to atom-type-3 and atom-type-4.
- **ModifyFrameworkTriple** [atom-type-1] [atom-type-2] [atom-type-3] [atom-type-4] [atom-type-5] [atom-type-6]  
Modifies the connected triple atom-type-1,atom-type-2,atom-type-3 to atom-type-4,atom-type-5,atom-type-6.
- **RemoveAtomNumberCodeFromLabel** [yes|no]  
Reading structure-files: the number is removed from the framework atom-types, e.g. '01', '02', '03', etc. are mapped to '0'.
- **AddAtomNumberCodeToLabel** [yes|no]  
Writing structure-files: the number is added to the framework atom-types, e.g. '0' are mapped to '01', '02', '03', etc.
- **RestrictFrameworkAtomsToBox** [yes|no]  
Restricts (places back) atoms to the unit cell dimensions, i.e. fractional positions between 0 and 1.
- **ReadCIFAsCartesian** [yes|no]  
Reads the position listed in the CIF-file as Cartesian. Only applicable to P1 systems (no symmetry).

## System moves

- FrameworkChangeMoveProbability [real]

The probability per cycle to randomly translate a framework atom. During this move the number of inner cycles is the amount of framework atoms, with a maximum of 500. This move is applicable to relatively rigid structures like zeolites. For other structure where movement is caused by collective behavior (for example, the rotation of a phenyl-ring in a metal-organic framework) the MC/MD move is more convenient. Such movement is hardly sampled at all by individual MC translation moves.

- VolumeChangeProbability [real]

The probability per cycle to attempt a volume-change. Rigid molecules are scaled by center-of-mass, while flexible molecules and the framework is atomically scaled.

- BoxShapeChangeProbability [real]

The probability per cycle to attempt a shape-change of the box. One of the 6 upper triangular elements of the box matrix is randomly chosen. Rigid molecules are scaled by center-of-mass, while flexible molecules and the framework is atomically scaled.

- GibbVolumeChangeProbability [real]

The probability per cycle to attempt a Gibbs volume-change MC move during a Gibbs ensemble simulation. The total volume of the two boxes (usually one for the gas phase, one for the liquid phase) remains constant, but the individual volume of the boxes are changed. The volumes are changed by a random change in  $\ln(V_I/V_{II})$ .

- HybridNVEMoveProbability [real]

The probability per cycle to attempt a hybrid Monte Carlo move using Molecular Dynamics in the NVE-ensemble. The whole system is integrated using Newton's equations of motion. The new configuration is then accepted or rejected using the standard MC rule. Note that the difference in energy  $\Delta U$  is the integration error. The integration time step is set using 'TimeStep'.

- NumberOfHybridNVESteps [int]

The number of integration steps for the hybrid MC/MD NVE move. Default: 5.

- HybridNPHMoveProbability [real]

The probability per cycle to attempt a hybrid Monte Carlo move using Molecular Dynamics in the NPH-ensemble. The whole system is integrated using Newton's equations of motion. The new configuration is then accepted or rejected using the standard MC rule. Note that the difference in energy  $\Delta U$  is the integration error. The integration time step is set using 'TimeStep'.

- NumberOfHybridNPHSteps [int]

The number of integration steps for the hybrid MC/MD NPH move. Default: 5.

- HybridNPHPRMoveProbability [real]

The probability per cycle to attempt a hybrid Monte Carlo move using Molecular Dynamics in the NPH-ensemble with full cell fluctuations (Parrinello-Rahman). The whole system is integrated using Newton's equations of motion. The new configuration is then accepted or rejected using the standard MC rule. Note that the difference in energy  $\Delta U$  is the integration error. The integration time step is set using 'TimeStep'.

- NumberOfHybridNPHPRSteps [int]

The number of integration steps for the hybrid MC/MD NPH move (Parrinello-Rahman) with full cell fluctuations. Default: 5.

- ParallelTemperingProbability [real]

A move where two neighboring systems are swapped that differ in their temperature.

- HyperParallelTemperingProbability [real]  
A move where two neighboring systems are swapped that differ in their temperature and chemical potentials.
- ParallelMolFractionProbability [real]  
A move where two neighboring systems (similar to parallel tempering) are swapped that differ in their mol-fraction of components *A* and *B*.
- ParallelMolFractionComponentA [int]  
The identifier of the first component.
- ParallelMolFractionComponentB [int]  
The identifier of the second component.
- ChiralInversionProbability [real]  
A move specifically designed for systems with chiral molecules to change all *S*-molecules into *R*-molecules and vice versa. Note that the spacegroup needs to be set. If you have a framework that is P1 but has higher symmetry then use '`CalculateSpaceGroup yes`' to determine the true space group of the framework. An error will be given if this move is impossible for your system (e.g. when the framework is chiral).

## Component information

- Component [int] MoleculeName [string]  
Reads in the definition of component [int] using the file '*molecule-name-string.def*' from the directory '``${RASPA_DIR}/share/raspa/molecules/molecule-definitions-string``'.
- MoleculeDefinitions [string]  
The type of the molecule. For example, there could be an OPLS version of the molecule, or a TraPPE version, etc. This *molecule-definitions-string* is actually the directory name under which the molecule file is found in '``${RASPA_DIR}/share/raspa/molecules/``'.
- StartingBead [int]  
The starting bead for the configurational bias Monte Carlo (CBMC). In CBMC the molecule is grown bead by bead biasing the growth towards energetically favorable configurations. Certain operations, like the rotation MC move and Widom particle insertion, use this bead as the center of rotation and position of the probe molecule, respectively.
- BlockPockets [yes|no]  
Blocks certain pockets in the simulation volume. The growth of a molecule is not allowed in a blocked pocket. A typical example is the sodalite cages in FAU and LTA-type zeolites, these are not accessible to molecules like methane and bigger.
- BlockPocketsFileName [string]  
The file name for the definitions of all the blocking spheres.
- MolFraction [real]  
The mol fraction of this component in the mixture. The values can be specified relative to other components, as the fractions are normalized afterwards. The partial pressures for each component are computed from the total pressure and the mol fraction per component.
- FugacityCoefficient [real]  
The fugacity coefficient for the current component. For values 0 (or by not specifying this line), the fugacity coefficients are automatically computed using the Peng-Robinson equation of state. Note the critical pressure, critical temperature, and acentric factor need to be specified in the molecule file.

- Intra14VDWScalingValue [real]

The scaling factor for intra-molecular 1-4 van der Waals interactions. For example: OPLS uses a factor of  $\frac{1}{2}$ .

- Intra14ChargeChargeScalingValue [real]

The scaling factor for intra-molecular 1-4 charge/charge interactions. For example: OPLS uses a factor of  $\frac{1}{2}$ .

- IdealGasRosenbluthWeight [real]

The ideal Rosenbluth weight is the growth factor of the CBMC algorithm for a single chain in an empty box. The value only depends on temperature and therefore needs to be computed only once. For adsorption, specifying the value in advance is convenient because the applied pressure does not need to be corrected afterwards (the Rosenbluth weight corresponds to a shift in the chemical potential reference value, and the chemical potential is directly obtained from the fugacity). For equimolar mixtures this is essential.

- GibbSwapProbability [real]

The relative probability to attempt a Gibbs swap MC move for the current component. The "GibbsSwapMove" transfers a randomly selected particle from one box to the other (50% probability to transfer a particle from box I to II, an 50% visa versa).

- TranslationProbability [real]

The relative probability to attempt a translation move for the current component. A random displacement is chosen in the allowed directions (see 'TranslationDirection'). Note that the internal configuration of the molecule is unchanged by this move. The maximum displacement is scaled during the simulation to achieve an acceptance ratio of 50%.

- TranslationDirection [X|Y|Z|XY|XZ|YZ|XYZ|A|B|C|AB|AC|BC|ABC|

ORTHOGONAL\_TO\_AB\_DIR|ORTHOGONAL\_TO\_AC\_DIR|ORTHOGONAL\_TO\_BC\_DIR|

ORTHOGONAL\_TO\_O\_AB\_DIR|ORTHOGONAL\_TO\_O\_AC\_DIR|ORTHOGONAL\_TO\_O\_BC\_DIR|

ORTHOGONAL\_TO\_A\_BC\_DIR|ORTHOGONAL\_TO\_B\_AC\_DIR|ORTHOGONAL\_TO\_C\_AB\_DIR|

ORTHOGONAL\_TO\_O\_ABC\_DIR]

Specifies the allowed translation direction for the current component. Useful to sampling configuration with the starting bead restricted to a plane, i.e. see dcTST. Default: XYZ.

- RandomTranslationProbability [real]

The relative probability to attempt a random translation move for the current component. The displacement is chosen such that any position in the box can be reached. It is therefore similar as reinsertion, but 'reinsertion' changes the internal conformation of a molecule and uses biasing.

- RotationProbability [real]

The relative probability to attempt a random rotation move for the current component. The rotation is around the starting bead. A random vector on a sphere is generated, and the rotation is random around this vector.

- CBMCPromability [real]

The relative probability to attempt a partial reinsertion move for the current component. Part of the molecule is regrown, while part of the molecule can remain fixed. The list of partial reinsertion moves is specified in the 'molecule.def' file.

- ReinsertionProbability [real]

The relative probability to attempt a full reinsertion move for the current component. Multiple first beads are chosen, and one of these is selected according to its Boltzmann weight. The remaining part of the molecule is grown using biasing. This move is very useful, and often necessary, to change the internal configuration of flexible molecules.

- SwapProbability [real]

The relative probability to attempt a insertion or deletion move. Whether to insert or delete is decided randomly with a probability of 50% for each. The swap move imposes a chemical equilibrium between the system and an imaginary particle reservoir for the current component. The move starts with multiple first bead, and grows the remainder of the molecule using biasing.

- WidomProbability [real]

The relative probability to attempt a Widom particle insertion move for the current component. The Widom particle insertion moves measure the chemical potential and can be directly related to Henry coefficients and heats of adsorption.

- SurfaceAreaProbability [real]

The relative probability to attempt a surface-area move for the current component.

- ReinsertionInPlaceProbability [real]

The relative probability to attempt a reinsertion-in-place move for the current component. The reinsertion position is the current position of the starting bead of the randomly selected molecule. Alternatively, one can use the partial reinsertion move leaving one bead fixed. The move is very useful to sample configuration on a plane for dcTST to change the internal configuration, e.g. bonds, bends, torsions, etc.

- IdentityChangeProbability [real]

The relative probability to attempt an identity-change move for the current component. A molecule of type *A* is reinsertion, in the same place as the starting bead of *A*, as type *B* using the starting bead of component *B*. The *A – B* list is defined using 'IdentityChangesList' defining *B* for each component *A*, i.e. the current component can be reinserted into any component defined in the 'IdentityChangesList' list, and from that list the component is chosen randomly.

- NumberOfIdentityChanges [int]

The number of 'IdentityChangesList' elements for the current component.

- IdentityChangesList [list-of-int]

The list of components that the current component can be changed into. The identity-change move will randomly choose the new component from this list.

- GibbIdentityChangeProbability [real]

The relative probability to attempt an identity change for the current component in the Gibbs ensemble. It is a very useful move to for mixture of *n* components. Out of the *n* components, two components *i* ≠ *j* are selected at random. At random, it is selected to switch the identity of component *i* in box *I* or in box *II*, and the identity of the component *j* in the other box. In each box, a particle is selected at random which matches the desired identity.

- NumberOfGibbsIdentityChanges [int]

The number of 'GibbsIdentityChangesList' elements for the current component.

- GibbsIdentityChangesList [list-of-int]

The list of components that the current component can be changed into. The Gibbs-identity-change move will randomly choose the new component from this list.

- ExtraFrameworkMolecule [yes|no]

There are two major types of molecules, 'Adsorbates' and 'Cations'. The 'ExtraFrameworkMolecule' keyword sets whether the current component is a 'Cation' (yes) or a 'Adsorbate' (no). Energies in the output as splitted in Host-Host, Host-Adsorbate, Host-Cation, Adsorbate-Adsorbate, Cation-Cation, and Adsorbate-Cation. The distinction in two types of molecule is sometimes necessary. For example, consider a mixture of components, where polarization needs to be neglected between certain components (because they are parameterized without). The water model 'rpol' is defined including polarization,

but CO<sub>2</sub> using TraPPE is not. One can define water as 'Adsorbate', CO<sub>2</sub> as 'Cation' and neglect polarization between cations.

- RestrictEnantionface [yes|no]

Restricts all MC-moves to the enantioface defined by 'Enantioface'. Moves that result in an opposite enantioface are rejected.

- Enantioface [Re|Si]

The enantioface of the component, either 'Re' or 'Si'.

- EnantiofaceAtoms [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int]

The definition of the enantioface based on 5 atoms. The first 4 form a torsion, as well as the first 3 and the last atom. These two torsions form the definition of the enantioface.

- CreateNumberOfMolecules [int]

The number of molecule to create for the current component. Note these molecules are *in addition* to anything read in by using a restart-file. Usually, when the restart-file is used the amount here should be put back to zero. A warning, putting this value unreasonably high results in an infinite loop. The routine accepts molecules that are grown causing no overlap (energy smaller than 'EnergyOverlapCriteria'). Also the initial starting configurations are far from optimal and substantial equilibration is needed to reduce the energy. However, the CBMC growth is able to reach very high densities.

## Options to measure properties

- ComputeDistanceHistograms [yes|no]

Sets whether or not to compute the histograms of specified distance pairs for the current system. A directory 'DistanceHistograms' is created containing the histograms for each system.

- WriteDistanceHistogramEvery [int]

Output the distance histograms every [int] cycles.

- MaxRangeDistanceHistogram [real]

The range of the histograms.

- NumberOfElementsDistanceHistogram [int]

The number of elements of the histograms.

- DistanceHistogramDefinition [F|A|C] [int] [int] [F|A|C] [int] [int]

Define a distance histogram between two atoms.

- ComputeBendAngleHistograms [yes|no]

Sets whether or not to compute the bend-angle histograms of specified trimers of atoms for the current system. A directory 'BendAngleHistograms' is created containing the histograms for each system.

- WriteBendAngleHistogramEvery [int]

Output the distance histograms every [int] cycles.

- MaxRangeBendAngleHistogram [real]

- NumberOfElementsBendAngleHistogram [int]

- BendAngleHistogramDefinition [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int]

- ComputeDihedralAngleHistograms [yes|no]

Sets whether or not to compute the dihedral-angle histograms of specified quads of atoms for the current system. A directory 'DihedralAngleHistograms' is created containing the histograms for each system.

- WriteDihedralAngleHistogramEvery [int]  
Output the distance histograms every [int] cycles.
  - MaxRangeDihedralAngleHistogram [real]
  - NumberOfElementsDihedralAngleHistogram [int]
  - DihedralAngleHistogramDefinition [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int]
- ComputeAngleBetweenPlanesHistograms [yes|no]  
Sets whether or not to compute the histograms of angles between specified planes for the current system. A directory 'AngleBetweenPlanesHistograms' is created containing the histograms for each system.
  - WriteAngleBetweenPlanesHistogramEvery [int]  
Output the distance histograms every [int] cycles.
  - MaxRangeAngleBetweenPlanesHistogram [real]
  - NumberOfElementsAngleBetweenPlanesHistogram [int]
  - AngleBetweenPlanesHistogramDefinition [F|A|C] [int] [int] [F|A|C] [int] [int]
- ComputePSD [yes|no]  
Sets whether or not to compute the pore-size distribution (PSD) for the current system. A directory 'PoreSizeDistributionHistogram' is created containing the output 'HistogramPoreSizeDistribution.dat' per system.
  - WritePSDEvery [int]  
Output the PSD every [int] cycles.
  - PSDProbeDistance [Minimum|Sigma]  
Sets whether to use the minimum of the potential  $\sigma^{1/6}$  as the probe distance or whether to use  $\sigma$ .
  - HistogramSizePoreSizeDistribution [int]  
default: 100.
  - MaxRangePoreSizeDistribution [real]  
default: 10.
- ComputeRDF [yes|no]  
Sets whether or not to compute the radial distribution function (RDF) for the current system. A directory 'RadialDistributionFunctions' is created containing the output per system. The RDF is computed for each atom type pair unless the option 'print' flag in 'pseudo\_atoms.def' is 'no'.
  - WriteRDFEvery [int]  
Output the RDF every [int] cycles.
- ComputeMSD [yes|no]  
Sets whether or not to compute the mean-squared displacement (MSD) for the current system using a modified order-N algorithm. A directory 'MSDOrderN' is created containing the output per system. The output consists of files containing self-msd data per component, the total self-msd, the Onsager msd for each component pair, and the the total Onsager msd. The units in the files are Å<sup>2</sup> for the msd, and ps for time.

- WriteMSDEvery [int]  
Output the MSD every [int] cycles.
  - SampleMSDEvery [int]  
Samples every [int] integration steps. Default: 1.
  - ComputeIndividualMSD [yes|no]  
Computes the msd, not only per component, but also per molecule.
  - NumberOfBlocksMSD [int]  
The number of 'blocks' for the order- $n$  correlation measurement. Each block represent a different time-scale of sampling. Default: 25.
  - NumberOfBlockElementsMSD [int]  
The number of elements in each block. For example, if the number is 10, then the first block samples: 1, 2, 3, ..., 10, the second block 10, 20, 30, ..., 100, the third block 100, 200, 300, ..., 1000, etc. Default: 25.
- ComputeVACF [yes|no]  
Sets whether or not to compute the velocity autocorrelation function (VACF) for the current system using a modified order-N algorithm. A directory 'VACFOrderN' is created containing the output per system. The output consists of files containing self-vacf data per component, the total self-vacf, the Onsager vacf for each component pair, and the the total Onsager vacf. The files start with the integration diffusivity-values, computed using a generalization of the Simpson's rule (in the sense that it is exact for cubic polynomials and is valid for an odd as well as even number of intervals). The units in the files are Å<sup>2</sup>/ps for velocity, and ps for time.
  - WriteVACFEvery [int]  
Output the VACF every [int] cycles.
  - SampleVACFEvery [int]  
Samples every [int] integration steps. Default: 5.
  - ComputeIndividualVACF [yes|no]  
Computes the vacf, not only per component, but also per molecule.
  - NumberOfBlocksVACF [int]  
The number of 'blocks' for the order- $n$  correlation measurement. Each block represent a different time-scale of sampling. Default: 10.
  - NumberOfBlockElementsVACF [int]  
The number of elements in each block. For example, if the number is 10, then the first block samples: 1, 2, 3, ..., 10, the second block 10, 20, 30, ..., 100, the third block 100, 200, 300, ..., 1000, etc. Default: 5000.
- ComputeRVACF [yes|no]  
Sets whether or not to compute the rotational velocity autocorrelation function (RVACF) for the current system using a modified order-N algorithm. A directory 'RVACFOrderN' is created containing the output per system. The output consists of files containing self-rvacf data per component, the total self-rvacf, the Onsager rvacf for each component pair, and the the total Onsager rvacf. The files start with the integration diffusivity-values, computed using a generalization of the Simpson's rule (in the sense that it is exact for cubic polynomials and is valid for an odd as well as even number of intervals). The units in the files are Å<sup>2</sup>/ps for velocity, and ps for time.
  - WriteRVACFEvery [int]  
Output the RVACF every [int] cycles.
  - SampleRVACFEvery [int]  
Samples every [int] integration steps. Default: 5.

- ComputeIndividualRVACF [yes|no]  
Computes the vacf, not only per component, but also per molecule.
  - NumberOfBlocksRVACF [int]  
The number of 'blocks' for the order-*n* correlation measurement. Each block represent a different time-scale of sampling. Default: 10.
  - NumberOfBlockElementsRVACF [int]  
The number of elements in each block. For example, if the number is 10, then the first block samples: 1, 2, 3, ..., 10, the second block 10, 20, 30, ..., 100, the third block 100, 200, 300, ..., 1000, etc. Default: 5000.
- ComputeMOACF [yes|no]  
Sets whether or not to compute the molecular orientation velocity autocorrelation function (MOACF) for the current system using a modified order-N algorithm. A directory 'MOACFOrderN' is created containing the output per system. The output consists of files containing self-moacf data per component and the total self-rvacf. The units in the files are rad<sup>2</sup>/ps for velocity, and ps for time.
  - WriteMOACFEvery [int]  
Output the MOACF every [int] cycles.
  - SampleMOACFEvery [int]  
Samples every [int] integration steps. Default: 5.
  - ComputeIndividualMOACF [yes|no]  
Computes the moacf, not only per component, but also per molecule.
  - NumberOfBlocksMOACF [int]  
The number of 'blocks' for the order-*n* correlation measurement. Each block represent a different time-scale of sampling. Default: 10.
  - NumberOfBlockElementsMOACF [int]  
The number of elements in each block. For example, if the number is 10, then the first block samples: 1, 2, 3, ..., 10, the second block 10, 20, 30, ..., 100, the third block 100, 200, 300, ..., 1000, etc. Default: 5000.
- ComputeMSDConventional [yes|no]  
Sets whether or not to compute the mean-squared displacement (MSD) for the current system using the conventional algorithm. A directory 'MSD' is created containing the output per system. The routine is available for legacy reasons, the same results can be obtained using the order-N method and 1 block of size 'BufferLengthMSD'. The units in the files are Å<sup>2</sup> for the msd, and ps for time.
  - WriteMSDConventionalEvery [int]  
Output the MSD every [int] cycles. Default: 5000.
  - SampleMSDConventionalEvery [int]  
Samples every [int] integration steps. Default: 1.
  - NumberOfBuffersMSDConventional [int]  
The number of (overlapping) buffers with a different offset in time. Default: 20.
  - BufferLengthMSDConventional [int]  
The length of the buffers. Default: 5000.
- ComputeVACFConventional [yes|no]  
Sets whether or not to compute the velocity autocorrelation function (VACF) for the current system using the conventional algorithm. A directory 'VACF' is created containing the output per system. The routine is available for legacy reasons, the same results can be obtained using the order-N method and 1 block of size 'BufferLengthVACF'. The units in the files are Å<sup>2</sup>/ps for velocity, and ps for time.

- WriteVACFConventionalEvery [int]  
Output the VACF every [int] cycles. Default: 5000.
  - SampleVACFConventionalEvery [int]  
Samples every [int] integration steps. Default: 1.
  - NumberOfBuffersVACFConventional [int]  
The number of (overlapping) buffers with a different offset in time. Default: 20.
  - BufferLengthVACFConventional [int]  
The length of the buffers. Default: 5000.
- ComputeDensityHistograms [yes|no]  
Sets whether or not to compute a density histogram for the current system. For example, during adsorption it keeps track of the amount of molecules.
- ComputeEnergyHistogram [yes|no]  
Sets whether or not to compute a histogram of the energy for the current system. For example, during adsorption it keeps track of the total energy, the VDW energy, the Coulombic energy, and the polarization energy. Output is written to the directory 'EnergyHistograms'.
  - WriteEnergyHistogramEvery [int]  
Sets to print the energy histogram of the system every [int] cycles.
  - EnergyHistogramSize [int]  
Sets the number of elements of the histogram. Default: 1000.
  - EnergyHistogramLowerLimit [real]  
Sets the lower limit of the histogram. Default: -10000.
  - EnergyHistogramUpperLimit [real]  
Sets the upper limit of the histogram. Default: 0.
- ComputeThermoDynamicFactor [yes|no]  
Sets whether or not to compute the thermodynamic factors of the energy for the current system. The output is written to the directory 'ThermoDynamicFactor'.
  - WriteThermoDynamicFactorEvery [int]  
Sets to print the thermodynamic factors every [int] cycles.
- ComputeEndToEndDistanceHistograms [yes|no]  
Sets whether or not to compute a histogram for end-to-end distances of molecules for the current system.
- ComputePrincipleMomentsOfInertia [yes|no]  
Sets whether or not to compute the average principle moments of inertia of molecules for the current system.
- ComputeSpectra [yes|no]  
Sets whether or not to compute the Infra-Red (IR) spectra of molecules for the current system.
  - WriteSpectraEvery [int]  
Sets to print the spectra of molecules every [int] cycles.
- ComputeMoleculeProperties [yes|no]  
Sets whether or not to compute properties of molecules like average bond-lengths, average bend-angles etc. for the current system.
- PrintMoleculePropertiesEvery [int]  
Sets to print the properties of molecules every [int] cycles.

- ComputeSurfaceArea [yes|no]
 

Sets whether or not to compute the surface.

  - SurfaceAreaProbeAtom [string]
  - SurfaceAreaSamplingPointsPerSphere [int]
 

Sets the number of points to sampling a sphere per iteration.
  - SurfaceAreaProbeDistance [Minimum|Sigma]
 

Sets whether to use the minimum of the potential  $\sigma^{1/6}$  as the probe distance or whether to use  $\sigma$ .
- DensityProfile [yes|no]
- DensityProfileGridPoints [int] [int] [int]
- ComputeElasticConstants [yes|no]
 

Sets whether to compute elastic constants.
- ComputePowderDiffractionPattern [yes|no]
 

Sets whether to compute the powder diffraction pattern for the framework.

  - DiffractionType [Xray|Neutron|Electron]
 

Sets the diffraction type as xray-scattering, neutron-scattering, or electron-scattering, respectively.
  - DiffractionRadiationType [chromium|iron|copper|molybdenum|silver|synchrotron]
 

Sets the type of the diffraction radiation as chromium, iron, copper, molybdenum, silver, or synchrotron, respectively.
  - WaveLengthType [Single|Double]
 

Set the type of the beam as single or as a doublet.
  - PeakShape [Gaussian|Lorentzian|PseudoVoigt]
 

Sets the shape of the peaks as Gaussian, Lorentzian, or Pseudo-Voigt, respectively.
  - WaveLength [real]
 

Sets the wavelength of the diffraction beam.
  - TwoThetaMin [real]
 

Sets the minimum value of  $2\theta$ .
  - TwoThetaMax [real]
 

Sets the maximum value of  $2\theta$ .
  - TwoThetaStep [real]
 

Sets the step size of  $2\theta$ .
  - PeakWidthModifierU [real]
  - PeakWidthModifierV [real]
  - PeakWidthModifierW [real]
- ComputerNormalModes [yes|no]
 

Sets whether to compute normal modes.

  - MinimumMode [int]
 

Sets the minimum normal to compute.

- MaximumMode [int]  
Sets the maximum normal to compute.
- ModeResolution [int]

## Energy/force grid options

- UseTabularGrid [yes|no]  
Use a pre-tabulated grid for the energy and forces. Default: no.
- SpacingVDWGrid [real]  
The grid spacing of the Van der Waals potentials. Default: 0.15 Angstrom.
- SpacingCoulombGrid [real]  
The grid spacing of the Coulomb potential. Default: 0.15 Angstrom.
- GridTypes [list-of-strings]  
A list of atom-types for each of the used grids.

## Minimization/Saddle point search

- MinimizationMethod [Baker|NewtonRaphson|SaddlePointSearch]  
The Baker minimization method uses the eigenvalues/vectors to find a true minimum where all eigenvalues are positive. Newton-Raphson uses the first and second derivatives, but not the eigenvalues/vectors. The saddle point search can best be started from a minimum energy configuration. The algorithm walks up hill along the softest eigen mode to find a first order saddle point.
- MinimizationVariables [Cartesian|Fractional]  
Whether the minimization is performed in Cartesian or fractional positions. For some crystal minimizations it might be more convenient to choose fractional positions. An example is when one wants to keep a particular fractional position fixed during the minimization.
- MaximumNumberOfMinimizationSteps [int]  
The maximum number of minimization steps after which the minimization is stopped. Default: 10000.
- RMSGradientTolerance [real]  
Stopping criteria: the maximum allowed RMS gradient. Default:  $10^{-6}$ .
- MaxGradientTolerance [real]  
Stopping criteria: the maximum allowed gradient for each and every atom (and the strain elements for cell minimizations). Default:  $10^{-6}$ .
- MaximumStepLength [real]  
The maximum length of a minimization step. The length is dependent on the problem at hand. A too low value converges slowly (i.e. the minimization takes more steps), while a too high value might not converge at all. Default value: 0.3.
- FrameworkFixedInitialization [free|fixed]  
Sets all framework atoms as 'free' or 'fixed'. This command must precede individual overwrites and applies to the current system.
- AdsorbateFixedInitialization [free|fixed]  
Sets all adsorbate groups and atoms as 'free' or 'fixed'. This command must precede individual overwrites and applies to the current system.

- **CationFixedInitialization** [free|fixed]  
Sets all cation groups and atoms as 'free' or 'fixed'. This command must precede individual overwrites and applies to the current system.
- **ActiveFrameworkAtom** [int]  
Sets the atom of the current framework and system as 'active'.
- **ActiveFrameworkAtoms** [int] [list-of-ints]  
Sets the [int] atoms listed in [list-of-ints] of the current framework and system as 'active'.
- **FixedFrameworkAtom** [int]  
Sets the atom of the current framework and system as 'fixed'.
- **FixedFrameworkAtoms** [int] [list-of-ints]  
Sets the [int] atoms listed in [list-of-ints] of the current framework and system as 'fixed'.
- **ActiveAdsorbateMolecule** [int]  
Sets all atom and groups of the adsorbate molecule [int] as 'active'. Applies to the current system.
- **FixedAdsorbateMolecule** [int]  
Sets all atom and groups of the adsorbate molecule [int] as 'fixed'. Applies to the current system.
- **ActiveAdsorbateAtom** [int] [int]  
Sets an atom (second argument) of an adsorbate molecule (first argument) as 'active'. Applies to the current system.
- **FixedAdsorbateAtom** [int] [int]  
Sets an atom (second argument) of an adsorbate molecule (first argument) as 'fixed'. Applies to the current system.
- **ActiveAdsorbateGroup** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'active'. Applies to the current system and both center of mass and the orientation are set as 'active'.
- **FixedAdsorbateGroup** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'fixed'. Applies to the current system and both center of mass and the orientation are set as 'fixed'.
- **ActiveAdsorbateGroupCenterOfMass** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'active'. Applies to the current system and only the center of mass is set as 'active'.
- **FixedAdsorbateGroupCenterOfMass** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'fixed'. Applies to the current system and only the center of mass is set as 'fixed'.
- **ActiveAdsorbateGroupOrientation** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'active'. Applies to the current system and only the orientation is set as 'active'.
- **FixedAdsorbateGroupOrientation** [int] [int]  
Sets a group (second argument) of an adsorbate molecule (first argument) as 'fixed'. Applies to the current system and only the orientation is set as 'fixed'.
- **ActiveCationMolecule** [int]  
Sets all atom and groups of the cation molecule [int] as 'active'. Applies to the current system.

- FixedCationMolecule [int]  
Sets all atom and groups of the cation molecule [int] as 'fixed'. Applies to the current system.
- ActiveCationAtom [int] [int]  
Sets an atom (second argument) of an cation molecule (first argument) as 'active'. Applies to the current system.
- FixedCationAtom [int] [int]  
Sets an atom (second argument) of an cation molecule (first argument) as 'fixed'. Applies to the current system.
- ActiveCationGroup [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'active'. Applies to the current system and both center of mass and the orientation are set as 'active'.
- FixedCationGroup [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'fixed'. Applies to the current system and both center of mass and the orientation are set as 'fixed'.
- ActiveCationGroupCenterOfMass [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'active'. Applies to the current system and only the center of mass is set as 'active'.
- FixedCationGroupCenterOfMass [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'fixed'. Applies to the current system and only the center of mass is set as 'fixed'.
- ActiveCationGroupOrientation [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'active'. Applies to the current system and only the orientation is set as 'active'.
- FixedCationGroupOrientation [int] [int]  
Sets a group (second argument) of an cation molecule (first argument) as 'fixed'. Applies to the current system and only the orientation is set as 'fixed'.
- FixAtomType [string]
- FixAtomTypes [int] [list-of-strings]  
The atom-types that are considered fixed during the minimization. All other atoms/groups will be optimized. If the atom-type is contained in a rigid unit, the entire unit will be frozen.
- DistanceConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [real]  
Defines a 'hard' distance constraint between two atoms and/or groups, and the distance.
- AngleConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [real]  
Defines a 'hard' angular constraint between three atoms and/or groups, and the constraint angle.
- DihedralConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [real]  
Defines a 'hard' dihedral constraint between four atoms and/or groups, and constraint dihedral.
- HarmonicDistanceConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [real] [real]  
Defines a 'hard' distance constraint between two atoms and/or groups, and the distance.
- HarmonicAngleConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [real] [real]  
Defines a 'hard' angular constraint between three atoms and/or groups, and the constraint angle.
- HarmonicDihedralConstraint [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [F|A|C] [int] [int] [real] [real]  
Defines a 'hard' dihedral constraint between four atoms and/or groups, and constraint dihedral.

## Monte Carlo settings

- MinimumInnerCycles [int]  
The minimum number of inner cycles for each cycle. Default: 20.
- NumberOfTrialPositions [int]  
The number of trial positions during the growth of a molecule. Default: 10.
- NumberOfTrialPositionsForTheFirstBead [int]  
The number of trial positions for the first bead. Default: 10.
- NumberOfTrialPositionsTorsion [int]  
The number of trial positions for torsions over a single bond. Default: 100.
- NumberOfTrialMovesPerOpenBead [int]  
The number of trial moves per open bead during CBMC. Default: 200.
- TargetAccRatioSmallMCScheme [real]
- TargetAccRatioTranslation [real]
- EnergyOverlapCriteria [real]  
The energy criteria to consider an energy as 'overlap'. Default:  $10^5$  K.
- MinimumRosenbluthFactor [real]  
The minimum Rosenbluth weight, values lower are consider to be 'overlapping'. Default:  $10^{-150}$ .

## Biasing options

- BiasingDirection [A|B|C|AB\_DIAGONAL|AC\_DIAGONAL|BC\_DIAGONAL|  
A\_BC\_DIAGONAL|B\_AC\_DIAGONAL|C\_AB\_DIAGONAL|  
O\_ABC\_DIAGONAL]
- BiasingMethod [UMBRELLA|RUIZMONTERO]
- BiasingProfile [string]  
The name of the file containing the biasing profile.
- RuizMonteroFactor [real]
- UmbrellaFactor [real]  
The biasing free energy is multiplied by the UmbrellaFactor. This is useful when the biasing free energy goes to infinity in certain regions. If the exact free energy would be used to biased, then the histogram would be flat, even very close to atoms. To keep the repulsion one can lower the used free energy biasing by e.g. multiplying by 0.9.
- RestrictMovesToUnitCell [yes|no]  
Restrict the Monte-Carlo moves to the first unitcell for this component.
- BoxAxisABC\_Min [real]  
When a particle is restricted in all Monte-Carlo moves (RestrictMovesToUnitCell or RestrictMovesToBox) then do not allow trial moves with a fractional position smaller than BoxAxisABC\_Min.
- BoxAxisABC\_Max [real]  
When a particle is restricted in all Monte-Carlo moves (RestrictMovesToUnitCell or RestrictMovesToBox) then do not allow trial moves with a fractional position greater than BoxAxisABC\_Max.

## Transition State Theory settings

FreeEnergyMappingType [A\_MAPPING|B\_MAPPING|C\_MAPPING|ABC\_MAPPING|

MAP\_AB\_DIAGONAL|MAP\_AC\_DIAGONAL|MAP\_BC\_DIAGONAL|

MAP\_A\_BC\_DIAGONAL|MAP\_B\_AC\_DIAGONAL|MAP\_C\_AB\_DIAGONAL|

MAP\_O\_ABC\_DIAGONAL]

- Determines how the free energy profile is constructed from the contributions of points in the unit cell. The free energy is computed using Widom insertion by inserting probe molecules at many random position inside the unit cell. The 'FreeEnergyMappingType' maps a Cartesian position on a reaction coordinate 'q'. The mappings 'A\_MAPPING', 'B\_MAPPING', 'C\_MAPPING' map the Cartesian position onto the 'a', 'b', 'c' lattice vectors. The diagonal mapping maps onto diagonal, either in 2D or in 3D. For example, 'MAP\_A\_BC\_DIAGONAL' maps onto the line from 'A' to 'B+C' where 'A', 'B', and 'C' are the end points of the lattive vectors; and 'MAP\_O\_ABC\_DIAGONAL' maps onto the line from the origin to the opposite point 'A+B+C' on the diagonal.

PositionHistogramMappingType [A\_MAPPING|B\_MAPPING|C\_MAPPING|ABC\_MAPPING|

MAP\_AB\_DIAGONAL|MAP\_AC\_DIAGONAL|MAP\_BC\_DIAGONAL|

MAP\_A\_BC\_DIAGONAL|MAP\_B\_AC\_DIAGONAL|MAP\_C\_AB\_DIAGONAL|

MAP\_O\_ABC\_DIAGONAL]

- Determines how the position histogram is constructed from the contributions of points in the unit cell. The free energy is computed from the histogram by using  $F(q) = -\log[P(q)]$ . The 'PositionHistogramMappingType' maps a Cartesian position on a reaction coordinate 'q'. The mapings 'A\_MAPPING', 'B\_MAPPING', 'C\_MAPPING' map the Cartesian position onto the 'a', 'b', 'c' lattice vectors. The diagonal mapping maps onto diagonal, either in 2D or in 3D. For example, 'MAP\_A\_BC\_DIAGONAL' maps onto the line from 'A' to 'B+C' where 'A', 'B', and 'C' are the end points of the lattive vectors; and 'MAP\_O\_ABC\_DIAGONAL' maps onto the line from the origin to the opposite point 'A+B+C' on the diagonal.

- PutMoleculeOnBarrier [yes|no]

Places the first molecule of component 0 at the position given by 'BarrierPosition'. This is used e.g. to start sampling configuration on top of a free energy barrier.

- BarrierPosition [real] [real] [real]

The location of the free energy barrier in fractional units of the first unit cell.

- MaxBarrierDistance [real]

The maximum distance in Ångstrom of the dcTST trajectory.

- MaxBarrierTime [real]

The maximum time of the dcTST trajetory in picoseconds.

- NumberOfVelocities [int]

The number of times the same initial position of the sampled dcTST starting configurations is used with different initial velocities.

- WritedcTSTSnapShotsToFile [yes|no]

Whether to write out sampled configuration to a file. The file is stored in the directory 'dcTST\_starting\_configurations' and used as the tarting point to compute the transmission coefficient in dcTST.

- WritedcTSTSnapShotsEvery [int]

The frequency in MC cycles of writing out the sampled configurations. Default: 1000.

## 2.3 Force field

### 2.3.1 Force fields

#### 2.3.2 'pseudo\_atoms.def'

The 'pseudo\_atoms.def' files describes the (pseudo-)atoms to be used in the simulation. An example is the definitions for the tip5p water model:

```
#number of pseudo atoms
3
#type    print   as     chem   oxidation   mass      charge   polarization   B-factor   radii   connectivity   anisotropic   anisotropic-type   tinker-type
Ow      yes     O     -       -          15.9996   0.0      0.0        1.0      0.5      2       0           absolute      0
Hw      yes     H     -       -          1.0008    0.241    0.0        1.0      1.00     1       0           absolute      0
Lw      no      L     -       -          0.0       -0.241   0.0        1.0      1.00     1       0           absolute      0
```

The first line is skipped, the second line is the number of (pseudo-)atoms, the third line is skipped again, and next all the (pseudo-)atoms are specified. The format and meaning is:

name	An unique string of character to be used to identify the atom. The same name has be used in other files to refer to this atom.
print	Whether or not this atom should be printed to movies. The dummy 'L' atoms of the tip5p water model are an example where you would like them to be skipped, only the 'O' and 'H' atoms should be printed.
as	The string to be printed to movies.
chem	The chemical symbol, e.g. O, O <sup>-</sup> , O <sup>2-</sup> . They are defined in 'scattering_factors.c' and are used only in powder diffraction and spectra.
oxidation state	
mass	The mass of the atom in atomic units.
charge	The charge of the atom in atomic units.
polarization	polarizability in units of Å <sup>3</sup> .
B-factor	The temperature factor of the atom, used only in powder diffraction.
radius	The radius of the atom to be used to decide what atoms are considered as 'neighbors'. The current rule is that two atoms <i>i</i> and <i>j</i> are considered 'bonded' if the distance between the atoms is smaller then 0.56+Radius <sub><i>i</i></sub> +Radius <sub><i>j</i></sub> .
connectivity	The connectivity of the atoms (not yet used).
anisotropic factor	The magntitude of the anisotropy.
anisotropic-type	The type of anisotropy, either 'relative' or 'absolute'. For example, a relative anisotropic factor of hydrogen of -0.077, used in the MM3 force field, means the site is pulled inward by 7.7% (and located at 92.3% of the C-H bondlength). An absolute anisotropic factor of e.g. 0.3 means the site is displaced outward by 0.3Å.
Tinker-type	The type of the atom in other codes, e.g. Tinker. This is used for output-files in formats used in other codes.

#### 2.3.3 'force\_field\_mixing\_rules.def'

```
# general rule for shifted vs truncated
shifted
# general rule tailcorrections
no
# number of defined interactions
9
# type interaction
Zn1           lennard-jones    0.42      2.7
```

```

01      lennard-jones    700.0    2.98
02      lennard-jones    70.5     3.11
C1      lennard-jones    48.5     3.76
C2      lennard-jones    47.86    3.47
C3      lennard-jones    47.86    3.47
H1      lennard-jones    7.65     2.85
O_co2    lennard-jones   80.507   3.033
C_co2    lennard-jones   28.129   2.757
# general mixing rule for Lennard-Jones
Jorgensen

```

The first line is skipped, the second line is the general cutoff rule for shifted vs truncated, the third line is skipped, the fourth line is the general rule for tail corrections, the fifth line is skipped, the sixth line is the number of defined self-interactions for the (pseudo-) atoms. The next line is skipped again followed by the defined potentials for the (pseudo-)atoms. The file is ended with a skipped line and the general rule for the mixing rule. Note all these interactions can be subsequently overwritten using the 'force\_field.def' file for specific interactions.

For convenience, you can use pattern matching of the interactions. Any string  $s_1$  ending with an underscore will match any string  $s_2$  that starts with the substring  $s_1$ . Note that usually the '\*' symbol is used, but this symbol has already a different meaning for CIF-files. Of course patterns can match more than one atom, e.g. 'C\_' matches 'C1' (a carbon atom) but also 'Cl' (chloride), and the rules are applied top to bottom. Therefore, list the generic ones first and the more specific after the generic patterns.

Example of a generic UFF/TraPPE force field for united atom alkanes in MOFs:

```

# general rule for shifted vs truncated
shifted
# general rule tailcorrections
no
# number of defined interactions
32
# type interaction, parameters.    IMPORTANT: define generic matches first
O_      lennard-jones    48.1581   3.03315
N_      lennard-jones    38.9492   3.26256
C_      lennard-jones    47.8562   3.47299
F_      lennard-jones    36.4834   3.0932
B_      lennard-jones    47.8058   3.58141
Cl_     lennard-jones    142.562   3.51932
Br_     lennard-jones    186.191   3.51905
H_      lennard-jones    7.64893   2.84642
Zn_     lennard-jones    62.3992   2.46155
Be_     lennard-jones    42.7736   2.44552
Cr_     lennard-jones    7.54829   2.69319
Fe_     lennard-jones    6.54185   2.5943
Mn_     lennard-jones    6.54185   2.63795
Cu_     lennard-jones    2.5161    3.11369
Co_     lennard-jones    7.04507   2.55866
Ga_     lennard-jones    208.836   3.90481
Ti_     lennard-jones    8.55473   2.8286
Sc_     lennard-jones    9.56117   2.93551
V_      lennard-jones    8.05151   2.80099
Ni_     lennard-jones    7.54829   2.52481
Zr_     lennard-jones    34.7221   2.78317
Mg_     lennard-jones    55.8574   2.69141

```

```

Ne_          lennard-jones    21.1352   2.88918
Ag_          lennard-jones    18.1159   2.80455
In_          lennard-jones    301.428   3.97608
Cd_          lennard-jones    114.734   2.53728
Sb_          lennard-jones    225.946   3.93777
Te_          lennard-jones    200.281   3.98232
He           lennard-jones    10.9      2.64
CH4_sp3     lennard-jones    158.5     3.72
CH3_sp3     lennard-jones    108.0     3.76
CH2_sp3     lennard-jones    56.0      3.96
# general mixing rule for Lennard-Jones
Lorentz-Berthelot

```

Here, 'CH4\_sp3', 'CH3\_sp3', and 'CH2\_sp3' are first matched by 'C\_-' but later overwritten with the correct values. However, carbon atoms listed in the MOF CIF-file, like 'C1', 'C2', 'C3', etc. will be set to the 'C\_-' value as intended.

general cutoff rule	'shifted' or 'truncated'	'shifted' shifts the potentials to zero at the cutoff radius, 'truncated' leaves them unchanged.
general tail corrections rule	'yes' or 'no'	'yes' applies the tail corrections to all interactions, 'no' omits the tail corrections for all interactions
general mixing-rule (only used for Lennard-Jones)	'Jorgensen' or 'Lorentz-Berthelot'	'Jorgensen' $\{\epsilon_{ij} = \sqrt{\epsilon_i \epsilon_j}, \sigma_{ij} = \sqrt{\sigma_i \sigma_j}\}$ 'Lorentz-Berthelot' $\{\epsilon_{ij} = \sqrt{\epsilon_i \epsilon_j}, \sigma_{ij} = \frac{1}{2}(\sigma_i + \sigma_j)\}$
self interaction type	'zero-potential', '12-6', 'Lennard-Jones', 'Buckingham', 'MCY', 'generic', 'HIW', 'MIE', 'BHM', or 'Hydrogen'	the type of the potential determines the subsequent parameters, i.e. Lennard-Jones expects a strength parameter $\epsilon_{ii}$ and a size parameter $\sigma_{ii}$ .

### 2.3.4 'force\_field.def'

The 'force\_field\_mixing\_rules.def' file given above can be used for the flexible model of the metal-organic framework IRMOF-1. It is defined using the Jorgensen mixing rule and uses shifted potentials cutoff at 12 Å. The EMP2-CO<sub>2</sub> model however uses the Lorentz-Berthelot for CO<sub>2</sub>-CO<sub>2</sub> interactions and uses a truncated potential with tail corrections. Moreover, if we also want to use the DREIDING model for the CO<sub>2</sub>-framework interactions the correction-file 'force\_field.def' would look like:

```

# rules to overwrite
3
# pair      truncated/shifted tailcorrections
O_co2 O_co2 truncated      yes
O_co2 C_co2 truncated      yes
C_co2 C_co2 truncated      yes
# number of defined interactions
14
# type      type2      interaction
Zn1       C_co2      lennard-jones 27.34776042 3.420
Zn1       O_co2      lennard-jones 46.77926891 3.545
O1        C_co2      lennard-jones 36.07117963 2.915
O1        O_co2      lennard-jones 61.70097244 3.04

```

```

02      C_co2      lennard-jones 36.07117963 2.915
02      O_co2      lennard-jones 61.70097244 3.04
C1      C_co2      lennard-jones 35.94746166 3.135
C1      O_co2      lennard-jones 61.48934867 3.26
C2      C_co2      lennard-jones 35.94746166 3.135
C2      O_co2      lennard-jones 61.48934867 3.26
C3      C_co2      lennard-jones 35.94746166 3.135
C3      O_co2      lennard-jones 61.48934867 3.26
H1      C_co2      lennard-jones] 14.37184748 2.825
H1      O_co2      lennard-jones] 24.58353107 2.95
# mixing rules to overwrite
1
#
0_co2 C_co2 Lorentz-Berthelot

```

Tip: always double check the complete interaction set printed in the output !

## 2.4 Molecules

The format of the molecules is designed to allow for a combination of flexible and rigid subunits. A molecule is made up of "groups", where a group is a collection of either rigid or flexible atoms.

### 2.4.1 Rigid molecule

An example of CO<sub>2</sub> as a rigid molecule.

```

# critical constants: Temperature [T], Pressure [Pa], and Acentric factor [-]
304.1282
7377300.0
0.22394
#Number Of Atoms
3
# Number of groups
1
# CO2-group
3
rigid
0 O_co2    0.0      0.0      1.16
1 C_co2    0.0      0.0      0.0
2 O_co2    0.0      0.0     -1.16
# Chiral centers Bond BondDipoles Bend UrayBradley InvBend Torsion Imp. Torsion Bond/Bond Stretch/Bend/Bend Stretch/Torsion Bend/Torsion IntraVDW IntraCoulomb
0      2      0      0      0      0      0      0      0      0      0      0      0      0      0
* Bond stretch: atom n1-n2, type, parameters
0 1 RIGID_BOND
1 2 RIGID_BOND
# Number of config moves
0

```

The first three numbers are the critical constants: the critical temperature, the critical pressure, and the acentric factor. They are used to automatically compute the fugacity from the pressure using an equation of state, e.g. Peng Robinson. Then the number of atoms and the number of groups. The groups are listed one by one with first the number of atoms in the group, whether it is rigid or flexible, and the atoms as number, type, and for a rigid molecule the relative positions. After the groups follows the bond, bend, torsion etc. parameters. The file is ended with the config moves.

### 2.4.2 Flexible molecule

An example of a flexible molecule is the united-atom 2-methylbutane molecule. Note that for flexible units there is not need to list relative positions. Bond-potentials are listed as the two atoms on which the potential operates, the potential type and the corresponding parameters. At the end we have 2 config moves defined, one where atoms 0,1,2 are kept fixed and the rest is regrown, and another config move where only atoms 2,3 are kept fixed.

```

# critical constants: Temperature [T], Pressure [Pa], and Acentric factor [-]
460.35
3395700.0
0.2296
# Number Of Atoms
5
# Number Of Groups
1
# Alkane-group
5
flexible
0 CH3_sp3
1 CH_sp3
2 CH2_sp3
3 CH3_sp3
4 CH3_sp3
# Chiral centers Bond BondDipoles Bend UrayBradley InvBend Torsion Imp. Torsion Bond/Bond Stretch/Bend/Bend Stretch/Torsion Bend/Torsion IntraVDW IntraCoulomb
0 4 0 4 0 0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
# Bond stretch: atom n1-n2, type, parameters
0 1 HARMONIC_BOND 96500 1.54
1 2 HARMONIC_BOND 96500 1.54
1 4 HARMONIC_BOND 96500 1.54
2 3 HARMONIC_BOND 96500 1.54
# Bond bending: atom n1-n2-n3, type, parameters
0 1 2 HARMONIC_BEND 62500 112
0 1 4 HARMONIC_BEND 62500 112
4 1 2 HARMONIC_BEND 62500 112
1 2 3 HARMONIC_BEND 62500 114
# Torsion n1-n2-n3-n4 type
0 1 2 3 OPLS_DIHEDRAL -251.06 428.73 -111.85 441.27
4 1 2 3 OPLS_DIHEDRAL -251.06 428.73 -111.85 441.27
# Number of config moves
2
# nr_fixed followed by a list
3 0 1 2
2 2 3

```

### 2.4.3 Rigid/Flexible molecule

Flexible and rigid units can easily be combined, as shown for the 1,4-benzenedicarboxylate (BDC) molecule. Note that the relative positions in the rigid units are recomputed in the molecular reference frame.

```

# critical constants: Temperature [T], Pressure [Pa], and Acentric factor [-]
0.0
0.0
0.0
#Number Of Atoms
16
# Number of groups
3
# carboxyl-group
3
flexible
0 Mof_Db
1 Mof_Ca
2 Mof_Db
# phenyl-ring
10
rigid
3 Mof_Cb 6.458 6.458 11.526
4 Mof_Cc 7.308 7.308 12.221
5 Mof_Cc 5.608 5.608 12.221
6 Mof_H 7.876 7.876 11.759
7 Mof_H 5.04 5.04 11.759
8 Mof_Cc 7.308 7.308 13.611
9 Mof_Cc 5.608 5.608 13.611
10 Mof_H 7.876 7.876 14.073
11 Mof_H 5.04 5.04 14.073
12 Mof_Cb 6.458 6.458 14.306
# carboxyl-group
3
flexible
13 Mof_Ca
14 Mof_Db
15 Mof_Db
# Chiral centers Bond BondDipoles Bend UrayBradley InvBend Torsion Imp. Torsion Bond/Bond Stretch/Bend/Bend Stretch/Torsion Bend/Torsion IntraVDW IntraCoulomb
0 16 0 10 0 0 16 0 0 0 0 0 0 0 0 0 0 0 0 80 80
# Bond stretch: atom n1-n2, type, parameters
0 1 HARMONIC_BOND 543840.64928424 1.27
1 2 HARMONIC_BOND 543840.64928424 1.27
1 3 HARMONIC_BOND 353750.919316375 1.44
3 4 RIGID_BOND
3 5 RIGID_BOND
4 6 RIGID_BOND
5 7 RIGID_BOND
4 8 RIGID_BOND
5 9 RIGID_BOND
8 10 RIGID_BOND
9 11 RIGID_BOND
8 12 RIGID_BOND
9 12 RIGID_BOND
12 13 HARMONIC_BOND 353750.919316375 1.44
13 14 HARMONIC_BOND 543840.64928424 1.27
13 15 HARMONIC_BOND 543840.64928424 1.27
...
...
```

## 2.4.4 Chiral molecules

44methylethyoctane, the left-handed form:

```

# critical constants: Temperature [T], Pressure [Pa], and Acentric factor [-]
535.6
2847232.5
0.325
# Number Of Atoms
11
# Number of groups
1
# octane+group
11
flexible
0 CH3_sp3
1 CH2_sp3
2 CH2_sp3
3 C_sp3
4 CH2_sp3
5 CH2_sp3
6 CH2_sp3
7 CH3_sp3
8 CH3_sp3
9 CH2_sp3
10 CH3_sp3
# Chiral centers Bond BondDipoles Bend UrayBradley InvBend Torsion Imp. Torsion Bond/Bond Stretch/Bend Bend/Bend Stretch/Torsion Bend/Torsion IntraVDW IntraCoulomb
    1   10      0   12      0     0       6      0     0     0     0     0     0     0     0     0     0     21     0
# chiral center
2 3 4 8 L
# Bond stretch: atom n1-n2, type, parameters
0 1 HARMONIC_BOND 96500 1.54
1 2 HARMONIC_BOND 96500 1.54
2 3 HARMONIC_BOND 96500 1.54
3 4 HARMONIC_BOND 96500 1.54
4 5 HARMONIC_BOND 96500 1.54
5 6 HARMONIC_BOND 96500 1.54
6 7 HARMONIC_BOND 96500 1.54
3 8 HARMONIC_BOND 96500 1.54
3 9 HARMONIC_BOND 96500 1.54
9 10 HARMONIC_BOND 96500 1.54
# Bond bending: atom n1-n2-n3, type, parameters
0 1 2 HARMONIC_BEND 62500 114
1 2 3 HARMONIC_BEND 62500 114
2 3 4 HARMONIC_BEND 62500 109.47
2 3 8 HARMONIC_BEND 62500 109.47
2 3 9 HARMONIC_BEND 62500 109.47
8 3 4 HARMONIC_BEND 62500 109.47
9 3 4 HARMONIC_BEND 62500 109.47
3 4 5 HARMONIC_BEND 62500 114
9 3 8 HARMONIC_BEND 62500 109.47
3 9 10 HARMONIC_BEND 62500 114
4 5 6 HARMONIC_BEND 62500 114
5 6 7 HARMONIC_BEND 62500 114
# Torsion: atom n1-n2-n3-n4, type, parameters
0 1 2 3 TRAPPE_DIHEDRAL 0.0 355.03 -68.19 791.32
1 2 3 4 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
1 2 3 8 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
1 2 3 9 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
2 3 4 5 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
2 3 9 10 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
8 3 4 5 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
10 9 3 4 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
9 3 4 5 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
3 4 5 6 TRAPPE_DIHEDRAL 0.0 355.03 -68.19 791.32
10 9 3 8 TRAPPE_DIHEDRAL 0.0 0.0 0.0 461.29
4 5 6 7 TRAPPE_DIHEDRAL 0.0 355.03 -68.19 791.32
# Intra VDW: atom n1-n2
0 4
0 5
0 6
0 7
0 8
0 9
0 10
1 5
1 6
1 7
1 10
2 6
2 7
3 7
5 10
6 8
6 9
6 10
7 8
7 9
7 10
# Number of config moves
0

```

while the right-handed form has

```

# chiral center
2 3 4 8 R

```

## 2.5 Framework

### 2.5.1 Asymmetric unit cell

Frameworks are often presented in literature using as much symmetry as possible to reduced the amount of atoms needed to describe the structure. Usually only the fractional positions of the atoms in the *asymmetric unit cell* are given. Given a space group and the unit cell parameters (length and angles) all other positions in the full unit cell can be generated. For example, the isoreticular metal-organic framework IRMOF-1 is published as 7 fractional positions, space group 225, a cubic unit cell with cell lengths of 25.832 Å, and  $\alpha = \beta = \gamma = 90^\circ$ . RASPA can read cif-files, and the structure can be put into a file (see 'IRMOF-1.cif' in 'structures/mofs/cif'):

```
data_IRMOF-1

_cell_length_a      25.832
_cell_length_b      25.832
_cell_length_c      25.832
_cell_angle_alpha   90
_cell_angle_beta    90
_cell_angle_gamma   90
_cell_volume        17237.5

_symmetry_cell_setting      cubic
_symmetry_space_group_name_Hall '-F 4 2 3'
_symmetry_space_group_name_H-M 'F m -3 m'
_symmetry_Int_Tables_number 225

loop_
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
Zn1      Zn      0.2934    0.2066    0.2066
O1       O       0.25      0.25      0.25
O2       O       0.2819    0.2181    0.134
C1       C       0.25      0.25      0.1113
C2       C       0.25      0.25      0.0538
C3       C       0.2829    0.2171    0.0269
H1       H       0.3049    0.1951    0.0448
```


RASPA can then be run using:

```
SimulationType      MonteCarlo
NumberOfCycles      0
InitializationCycles 0

Forcefield          GenericMOFs

Framework           0
FrameworkName       IRMOF-1
UnitCells           1 1 1
```

and in the directory `Movies/System_0/` several files appear:



(a) The seven asymmetric atoms of IRMOF-1. (b) The full unit cell of IRMOF-1 has 424 atoms.

Figure 2.1: Asymmetric unit cells: the left figure shows the seven crystallographically different atoms in the IRMOF-1 structure in ball-and-stick format. The 'copies' (crystallographically identical atoms) are shown as lines. The right figure shows the full unit cell of IRMOF-1 in ball-and-stick.

- **Framework\_0\_initial\_1\_1\_1.cif**

The framework in CIF-format at the start of the simulation.

- **Framework\_0\_initial\_1\_1\_1\_P1.cif**

The framework in CIF-format at the start of the simulation converted to P1 (no symmetry).

- **Framework\_0\_initial.pdb**

The framework in PDB-format at the start of the simulation converted to P1 (no symmetry).

The files named 'final' are the structures at the end of the simulation. There are several programs that can read and view CIF-files: e.g. Jmol (free), Mercury (free), Crystal Maker (commercial, free demo), Materials Studio (commercial), and Gaussview (commercial). The PDB-files can be viewed in the freely available VMD-program.

**Tip:** always double check the `Framework_0_initial_1_1_1_P1.cif`, if you see something strange then check `_symmetry_space_group_name_Hall` and the fractional positions.

Space group 225 has 192 elements and the first 10 elements look like (see the file 'src/spacegroup.c' for the complete set):

$x' = x$	$y' = y$	$z' = z$
$x' = -x$	$y' = -y$	$z' = z$
$x' = -x$	$y' = y$	$z' = -z$
$x' = x$	$y' = -y$	$z' = -z$
$x' = z$	$y' = x$	$z' = y$
$x' = z$	$y' = -x$	$z' = -y$
$x' = -z$	$y' = -x$	$z' = y$
$x' = -z$	$y' = x$	$z' = -y$
$x' = y;$	$y' = z$	$z' = x$
$x' = -y$	$y' = z$	$z' = -x$
$x' = y$	$y' = -z$	$z' = -x$
...		

The procedure to generate a unit cell is to loop over the elements of the spacegroup and the atoms in the asymmetric unit cell, and to apply simply all the rule. For each new  $x', y', z'$  position a check is needed whether the same position has already been added (doubles have to be removed). After this procedure the 7 positions have been expanded to 424 positions. The fractional positions are transformed in the final step to Cartesian positions.

### 2.5.2 Fractional occupancies in zeolites

The procedure from asymmetric to full unit cell is rather simple when the fractional occupancies are unity. However, quite often there is some disorder the type of atoms. For example, in zeolites like FAU the Si/Al ratio is specified, but it is unknown where the aluminum actually is. Zeolite X is faujasite with a high amount of aluminum. The FAU structure with a Si/Al ratio of unity is given by

```

data_NaX

_audit_creation_method RASPA-1.0
_audit_creation_date 2011-2-20
_audit_author_name 'David Dubbeldam'

_cell_length_a    25.099
_cell_length_b    25.099
_cell_length_c    25.099
_cell_angle_alpha 90
_cell_angle_beta  90
_cell_angle_gamma 90
_cell_volume       14273.9

_symmetry_cell_setting      cubic
_symmetry_space_group_name_Hall '-F 2uv 2vw 3'
_symmetry_space_group_name_H-M 'F d -3'
_symmetry_Int_Tables_number 203

loop_
_atom_site_label

```

```

_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
Si1      Si4+  -0.05381   0.12565   0.03508
Al1      Al3+  -0.05524   0.03639   0.12418
O1       O2-   -0.1099    0.0003    0.1056
O2       O2-   -0.0011    -0.0028   0.1416
O3       O2-   -0.0346    0.0758    0.0711
O4       O2-   -0.0693    0.0726    0.18

```

Now the aluminum and silicon are alternating and Löwestein rule is obeyed. For higher Si/Al ratios the 'Al' position is fractionally occupied and a certain percentage might actually be silicon. The procedure here is to first generate the full unit cell of FAU with 96 aluminum (the maximum amount) and judiciously replace aluminum by silicon in the full unit cell.

```

SimulationType          MC
NumberOfCycles          0
NumberOfInitializationCycles 0
PrintEvery              10

Forcefield               Local

Substitute 0 Al1 Si1
Substitute 5 Al1 Si1
Substitute 10 Al1 Si1
Substitute 15 Al1 Si1
Substitute 20 Al1 Si1
RandomlySubstitute 75 Al1 Si1

Framework 0
FrameworkName NaX
UnitCells 1 1 1
ExternalTemperature 300.0

```

It reads the CIF-file which is Nax with 96 aluminum. You can use two types of commands to replace an atom:

- Substitute  
For example. `Substitute 10 Al1 Si1` means replace the 10th Al1 by Si1.
- RandomlySubstitute  
For example, `RandomlySubstitute 75 Al1 Si1` means randomly substitute 75 Al1 by Si1.

When you do them both, first the fixed rules are substituted and next the random ones with the "left-overs". The first one "Substitute" is useful to always have the same structure. You could make a random structure once, look in the output which Al was substituted and use the next time the "Substitute" command. In this way, you always work with the spacegroup NaX structure (not in P1) which is afterwards change by specifying rules.

More problematic are when several atoms have fractional occupancies lower than unity. Consider IRMOF-8 shown in Fig. 2.2. The linker molecules are disordered over two possible positions. One of these needs to be selected per linker. First the unit cell is generated from the asymmetric unit cell and subsequently the unit cell needs to be edited. Program which can do just that are Materials Studio, Gaussview, etc. After

the cell has been created and edited, the file needs to be placed in 'structures/mofs/cif'. Structures with disorder needs to be created at unit cell level (P1).

Even more difficult is MOF-1. Here the cif-file also contains several possibilities, but is not a priori known which ones to choose, i.e. what is the structure of the Dabco unit (1,4-diazabicyclo[2.2.2]octane) within the framework? One possibility is to choose a structure and use a quantum code and minimize the periodic unit cell. The result is shown in Fig. 2.3.

Note that all these procedures are necessary, but it is still an open question, especially for MOFs, whether you can keep the framework rigid or not. However, it is very hard to calibrate a flexible framework model and for this a substantial amount of reliable experimental data is required.

### 2.5.3 Format of the framework atoms

The atom-types in CIF-files are constructed from the name of the element and an identifier, e.g. 'C10' carbon type 10. Usually these carbon atoms are different because they have either different charges or different Van der Waals parameters.

Sometimes a force field is defined to have interactions on an atom-type which depends on its neighbors. For example, the oxygen atom is different whether it is connected to a silicon or to an aluminum atom. Therefore the atom are labelled using

```
ModifyFrameworkAtomConnectedTo 01 Oa1 Al1
ModifyFrameworkAtomConnectedTo 02 Oa2 Al1
ModifyFrameworkAtomConnectedTo 03 Oa3 Al1
ModifyFrameworkAtomConnectedTo 04 Oa4 Al1
```

which modifies '01' to 'Oa1' when connected to 'Al1', etc. In the CIF-file you can list the new framework atom with unknown position '?'.

```
loop_
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
Si1      Si4+ -0.05381  0.12565  0.03508
Al1      Al3+ -0.05524  0.03639  0.12418
O1       O2-  -0.1099   0.0003   0.1056
O2       O2-  -0.0011  -0.0028   0.1416
O3       O2-  -0.0346   0.0758   0.0711
O4       O2-  -0.0693   0.0726   0.18
Oa1      O2-    ?       ?       ?
Oa2      O2-    ?       ?       ?
Oa3      O2-    ?       ?       ?
Oa4      O2-    ?       ?       ?
```

Alternatively, you can list the atom types 'Oa1'-'Oa4' in your 'pseudo\_atoms.def' file.

Other times a force field is defined as

```
# rules to overwrite
0
# number of defined interactions
4
# type      type2      interaction
0          0          lennard-jones      29.4338257      3.062219744
0          Si         lennard-jones      49.05711264     3.483346249
```

```

Si           Si           lennard-jones      81.76308187      3.962387454
CH4_sp3     O            lennard-jones      115.00          3.47
# mixing rules to overwrite
0

```

Here, we have that all oxygens in the framework are of the same type, and all silicon is of the same type. In this case, we would like to map '01', '02', '03', etc. to '0', and 'Si1', 'Si2', etc. to 'Si'. You can achieve this using

```
RemoveAtomNumberCodeFromLabel yes
```

Suppose you want to use MFI with only '0' and 'Si'. MFI is defined using

```

Si1      Si4+    0.42238   0.0565   -0.33598
Si2      Si4+    0.30716   0.02772  -0.1893
\.
01      02-     0.3726    0.0534   -0.2442
02      02-     0.3084    0.0587   -0.0789
\.

```

The force field in 'force\_field.def'

```

# rules to overwrite
0
# number of defined interactions
1
# type      type2      interaction
CH4_sp3     O            lennard-jones      115.00          3.47
# mixing rules to overwrite
0

```

The 'pseudo\_atom.def'

```

#number of pseudo atoms
3
#type      print    as   scatt mass      charge  polarization B-factor radii  connectivity anisotropic anisotropic-type tinker-type
0         yes      O     O   15.9994  -1.025   0.0       1.0     0.5      2        0        absolute   0
Si        yes      Si    Si   28.0855   2.05    0.0       1.0     1.18     4        0        absolute   0
CH4_sp3   yes      C     C   16.04246  0.0     0.0       1.0     1.00     0        0        absolute   0

```

and the output file will show:


```

Pseudo atoms: 2
=====
Pseudo Atom[ 0] Name Si      Oxydation:           Element: Si4+ pdb-name: Si   Scat. Types: 111 14 Mass=28.085498706 B-factor:0.000
               Charge=2.050  Polarization=0.017 [A^-3] (considered a charged atom and no polarization) Interactions: no
               Anisotropic factor: 0.000 [-] (Absolute), Radius: 1.110 [A]
Pseudo Atom[ 1] Name O      Oxydation:           Element: O2- pdb-name: O    Scat. Types: 105 8 Mass=15.999404927 B-factor:0.000
               Charge=-1.025  Polarization=3.880 [A^-3] (considered a charged atom and no polarization) Interactions: no
               Anisotropic factor: 0.000 [-] (Absolute), Radius: 0.660 [A]


```

## 2.5.4 Typing the atoms of the framework

Atoms from a pdb- or cif-file are usually labeled e.g. 'C' for a carbon atom. In many force fields different carbon types have different charges. It is necessary to 'type' the structure and RASPA contains tools to do this. Let's assume the original structure always contains elements like 'H', 'C', 'N', 'O', etc. and we want to type them 'Mof\_Ha', 'Mof\_Hb', etc. A force field type called 'Typing' preexists. It only defines the 'pseudo\_atoms.def' file:




(a) The IRMOF-8 structure as directly computed from the asymmetric positions and the space group.




(b) The IRMOF-8 after making a selection, shown is only one of the possibilities.

Figure 2.2: IRMOF-8 has linkers which are disordered, the linker atoms have a fractional occupancy of 0.5, The atoms however are not individually disorder and there are two disordered linker, one out of two possibilities needs to be selected per linker position.



(a) The MOF-1 structure from the cif-file.



(b) The MOF-1 structure edited and optimized with the quantum program dmol (plane wave code).

Figure 2.3: The MOF-1 structure is synthesized as  $[Zn_2(1,4\text{-bdc})_2(\text{Dabco})]$ . The Dabco (1,4-diazabicyclo[2.2.2]octane) is very disordered with occupancies of 0.38 for the carbon and 0.5 for the hydrogen. The cif-file shown on the left shows all possibilities on top of each other. Here, just choosing one of the possibilities is difficult and it is not obvious which atoms to select. The brute force method is to select one possible choice and use a quantum plane wave for periodic structures and optimize the full unit cell. In this case it is feasible because of the low amount of atoms in the unit cell (only 54 atoms).

```

#number of pseudo atoms
43
#type    print   as   scat   mass   charge   polarization   B-factor   radii   connectivity
UNIT      no      H     H      1.0     1.0      0.0          1.0       1.0      0
He        yes     He    He    4.002602  0.0      0.0          1.0       1.0      0
Zn        yes     Zn1   Zn    65.37    0.0      0.0          1.0      1.448     0
Zn1       yes     Zn1   Zn    65.37    0.0      0.0          1.0      1.448     0
Cu        yes     Cu1   Cu    63.546   0.0      0.0          1.0       1.4      0
Cu1       yes     Cu1   Cu    63.546   0.0      0.0          1.0       1.4      0
O         yes     O     O    15.9994   0.0      0.0          1.0       0.68     2
O1        yes     O1    O    15.9994   0.0      0.0          1.0       0.68     2
O2        yes     O2    O    15.9994   0.0      0.0          1.0       0.68     2
O3        yes     O3    O    15.9994   0.0      0.0          1.0       0.68     2
O4        yes     O4    O    15.9994   0.0      0.0          1.0       0.68     2
C          yes     C     C    12.0107   0.0      0.0          1.0      0.720     0
C1        yes     C1    C    12.0107   0.0      0.0          1.0      0.720     0
C2        yes     C2    C    12.0107   0.0      0.0          1.0      0.720     0
C3        yes     C3    C    12.0107   0.0      0.0          1.0      0.720     0
C4        yes     C4    C    12.0107   0.0      0.0          1.0      0.720     0
C5        yes     C5    C    12.0107   0.0      0.0          1.0      0.720     0
C6        yes     C6    C    12.0107   0.0      0.0          1.0      0.720     0
C7        yes     C7    C    12.0107   0.0      0.0          1.0      0.720     0
C8        yes     C8    C    12.0107   0.0      0.0          1.0      0.720     0
C9        yes     C9    C    12.0107   0.0      0.0          1.0      0.720     0
C10       yes     C10   C    12.0107   0.0      0.0          1.0      0.720     0
C11       yes     C11   C    12.0107   0.0      0.0          1.0      0.720     0
C12       yes     C12   C    12.0107   0.0      0.0          1.0      0.720     0
C13       yes     C13   C    12.0107   0.0      0.0          1.0      0.720     0
C14       yes     C14   C    12.0107   0.0      0.0          1.0      0.720     0
C15       yes     C15   C    12.0107   0.0      0.0          1.0      0.720     0
C16       yes     C16   C    12.0107   0.0      0.0          1.0      0.720     0
N          yes     N     N    14.00674  0.0      0.0          1.0       0.68     0
N1         yes     N1    N    14.00674  0.0      0.0          1.0       0.68     0
N2         yes     N2    N    14.00674  0.0      0.0          1.0       0.68     0
N3         yes     N3    N    14.00674  0.0      0.0          1.0       0.68     0
N4         yes     N4    N    14.00674  0.0      0.0          1.0       0.68     0
H          yes     H     H    1.00794   0.0      0.0          1.0      0.320     0
H1         yes     H1    H    1.00794   0.0      0.0          1.0      0.320     0
H2         yes     H2    H    1.00794   0.0      0.0          1.0      0.320     0
H3         yes     H3    H    1.00794   0.0      0.0          1.0      0.320     0
H4         yes     H4    H    1.00794   0.0      0.0          1.0      0.320     0
H5         yes     H5    H    1.00794   0.0      0.0          1.0      0.320     0
H6         yes     H6    H    1.00794   0.0      0.0          1.0      0.320     0
H7         yes     H7    H    1.00794   0.0      0.0          1.0      0.320     0
H8         yes     H8    H    1.00794   0.0      0.0          1.0      0.320     0
H9         yes     H9    H    1.00794   0.0      0.0          1.0      0.320     0

```

As an example, let's type the structure 'NU-100' [1]. Figure 2.4 shows the NU-100 cluster with linkers and metal-corners. The pictures shows the different types of atoms and has been used to compute CHelpG charges. In the RASPA input-file you can use the typing command:

```
ModifyFrameworkAtomConnectedTo C Mof_Ca O
```

Look for a 'C' atom, check if it is connect to an 'O' atom and if so, type it 'Mof\_Ca'. It is also possible to define two neighbors:

```
ModifyFrameworkAtomConnectedTo C Mof_Cc Mof_Cb Mof_Cb
```




Figure 2.4: Cluster used for deriving partial charges on atoms in NU-100SP [1].

Look for an 'C' atom, if it is connect to a 'Mof\_Cb' and to another 'Mof\_Cb' atom, then type is 'Mof\_Cc'.  
The input-file to type 'NU-100' is

```

SimulationType          MC
NumberOfCycles          0

Forcefield               Local

Framework 0
FrameworkName NU-100SP
UnitCells 1 1 1
InputFileType cssr
ExternalTemperature 298.0

ModifyFrameworkAtomConnectedTo C C1 0
ModifyFrameworkAtomConnectedTo C C2 C1
ModifyFrameworkAtomConnectedTo C C3 C2 C2
ModifyFrameworkAtomConnectedTo C C4 C2
ModifyFrameworkAtomConnectedTo C C5 C4
ModifyFrameworkAtomConnectedTo C C6 C5
ModifyFrameworkAtomConnectedTo C C7 C6
ModifyFrameworkAtomConnectedTo C C8 C7

```

```

ModifyFrameworkAtomConnectedTo C C9 C8
ModifyFrameworkAtomConnectedTo C C10 C9
ModifyFrameworkAtomConnectedTo C C11 C10
ModifyFrameworkAtomConnectedTo C C12 C11
ModifyFrameworkAtomConnectedTo C C13 C12
ModifyFrameworkAtomConnectedTo C C14 C13
ModifyFrameworkAtomConnectedTo C C15 C14
ModifyFrameworkAtomConnectedTo H H1 C3
ModifyFrameworkAtomConnectedTo H H2 C4
ModifyFrameworkAtomConnectedTo H H3 C9
ModifyFrameworkAtomConnectedTo H H4 C10
ModifyFrameworkAtomConnectedTo H H5 C15
ModifyFrameworkAtomConnectedTo O O2 C1
ModifyFrameworkAtomConnectedTo Cu Cu O2

```

For MOFs, the easiest start-point to type is the carboxylate group. The carbon connected to the oxygen is typed 'Mof\_Ca', the carbon connected to 'Mof\_Cb' is typed 'Mof\_Cc'. The third line is important: the carbon should only be typed 'Mof\_Cc' when it is connected to an 'Mof\_Cb' and another 'Mof\_Cb'. This must be done like this, otherwise the atom which is above called 'Mof\_Cd' would also be wrongly labeled 'Mof\_Cc'. After running RASPA, the 'Movie'-directory contains the file 'Framework\_intitial.cssr' which is the cssr-file with complete typing. This file can be copied to 'structures/mofs/cssr' and given an appropriate name. Each pseudatom type can now be assigned a different charge in the 'pseudo\_atoms.def' file of the 'NU-100' forcefield.

Note that the lines containing the typing-rules are performed top to bottom and in later rules one can use the new names of the previous rules.

## 2.6 Using CIF-files

### 2.6.1 Definition of CIF-files

CIF files present crystallographic data in an human readable free format. Let's look at an example:

```

data_FAU_SI

_audit_creation_method RASPA-1.0
_audit_creation_date 2011-2-19
_audit_author_name 'David Dubbeldam'

_citation_author_name      'J.J. Hriljac, M.M. Eddy, A.K. Cheetham, J.A. Donohue, and G.J. Ray'
_citation_title            'Powder Neutron Diffraction and Si-29 MAS NMR Studies of Siliceous Zeolite-Y'
_citation_journal_abbrev   'J. Solid State Chem.'
_citation_journal_volume  106
_citation_page_first      66
_citation_page_last       72
_citation_year             1993

_cell_length_a    24.2576
_cell_length_b    24.2576
_cell_length_c    24.2576
_cell_angle_alpha 90
_cell_angle_beta  90
_cell_angle_gamma 90
_cell_volume      14273.9

_symmetry_cell_setting cubic
_symmetry_space_group_name_Hall '-F 4vw 2vw 3'
_symmetry_space_group_name_H-M  'F d -3 m'
_symmetry_Int_Tables_number 227

```

```

loop_
_symmetry_equiv_pos_as_xyz
'x,y,z'
'-x+3/4,-y+1/4,z+1/2'
.....
.....
.....
'z,-y+3/4,-x+3/4'
'z+1/2,y+1/2,x'

loop_
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_charge
_atom_site_polarization
Si1      Si4+   -0.05392   0.1253    0.03589   2.05   0
O1        O2-     0         -0.10623   0.10623  -1.025   0
O2        O2-   -0.00323  -0.00323   0.14066  -1.025   0
O3        O2-     0.0757   0.0757   -0.03577  -1.025   0
O4        O2-   0.07063   0.07063   0.32115  -1.025   0

```

The 'data\_' string signal the start of a data block. Each data block corresponds to a different structures, and typically only one structure is present (although it possible to combine more than one structure in a single file). The CIF instructions are divided into *data name categories*, such as '\_atom\_site\_' to describe atomic site parameters, '\_cell\_' to describe the cell parameters, '\_symmetry\_' to specify space group symmetry, etc. CIF data names begin with an underscore. For some data name the data can be provided using a list of data items. Such data items are preceded by a 'loop\_' string. The '\_atom\_site\_' section is typical example. The order of the data items correspond to the order in which the actual data is provided.

A nice feature of CIFs is that one can easily extend the syntax to include non-standard data item. For example, in the '\_atom\_site\_' section, the data items '\_atom\_site\_label', '\_atom\_site\_type\_symbol', '\_atom\_site\_fract\_x', '\_atom\_site\_fract\_y', '\_atom\_site\_charge' belong to the official CIF specification, but '\_atom\_site\_charge', '\_atom\_site\_polarization', '\_atom\_site\_anisotropic\_displacement', '\_atom\_site\_anisotropic\_type', and '\_atom\_site\_print\_to\_pdb' have been added in RASPA CIFs. Used in this fashion, they provide a replacement for the 'pseudo\_atoms.def' file. Note that if a 'pseudo\_atoms.def' file is used, the value in that file will have preference over the CIF-file values (if they both defined the same atom-type).

Note: It is possible to give crystallographically identical framework atoms different charges. If you specify the charges via the CIF-file while not defining them using the 'pseudo\_atoms.def' file, then the charges will be set per atom. The space group will 'expand' these charges to the copies of the asymmetric framework. If, however, the type of the framework atom is also present in the 'pseudo\_atoms.def' file then the values from the 'pseudo\_atoms.def' have precedence. Charge can only be set via CIF-files and MOL-files. For MOL-files, the mol-file has precedence over the 'pseudo\_atoms.def' file (opposite behavior as CIF-file). For CSSR-files the charge will always be taken from the 'pseudo\_atoms.def' file. Charges read from a restart-file have precedence over the 'pseudo\_atoms.def' file.

## Appendix: space group information

triclinic									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
1	1	P 1	P 1	cell choice 1	primitive	1	yes	no	no
2	2	P -1	-P 1	cell choice 1	primitive	2	yes	no	no

Table 2.1: Triclinic spacegroup information.

monoclinic									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
3	3	P 1 2 1	P 2y	unique axis b	primitive	2	no	yes	no
4	3	P 1 1 2	P 2	unique axis c	primitive	2	no	yes	no
5	3	P 2 1 1	P 2x	unique axis a	primitive	2	no	yes	no
6	4	P 1 21 1	P 2yb	unique axis b	primitive	2	no	yes	no
7	4	P 1 1 21	P 2c	unique axis c	primitive	2	no	yes	no
8	4	P 21 1 1	P 2xa	unique axis a	primitive	2	no	yes	no
9	5	C 1 2 1	C 2y	b, cell choice 1	c	4	no	yes	no
10	5	A 1 2 1	A 2y	b, cell choice 2	a	4	no	yes	no
11	5	I 1 2 1	I 2y	b, cell choice 3	body	4	no	yes	no
12	5	A 1 1 2	A 2	c, cell choice 1	a	4	no	yes	no
13	5	B 1 1 2	B 2	c, cell choice 2	b	4	no	yes	no
14	5	I 1 1 2	I 2	c, cell choice 3	body	4	no	yes	no
15	5	B 2 1 1	B 2x	a, cell choice 1	b	4	no	yes	no
16	5	C 2 1 1	C 2x	a, cell choice 2	c	4	no	yes	no
17	5	I 2 1 1	I 2x	a, cell choice 3	body	4	no	yes	no
18	6	P 1 m 1	P -2y	unique axis b	primitive	2	no	no	no
19	6	P 1 1 m	P -2	unique axis c	primitive	2	no	no	no
20	6	P m 1 1	P -2x	unique axis a	primitive	2	no	no	no
21	7	P 1 c 1	P -2yc	b, cell choice 1	primitive	2	no	no	no
22	7	P 1 n 1	P -2yac	b, cell choice 2	primitive	2	no	no	no
23	7	P 1 a 1	P -2ya	b, cell choice 3	primitive	2	no	no	no
24	7	P 1 1 a	P -2a	c, cell choice 1	primitive	2	no	no	no
25	7	P 1 1 n	P -2ab	c, cell choice 2	primitive	2	no	no	no
26	7	P 1 1 b	P -2b	c, cell choice 3	primitive	2	no	no	no
27	7	P b 1 1	P -2xb	a, cell choice 1	primitive	2	no	no	no
28	7	P n 1 1	P -2xbc	a, cell choice 2	primitive	2	no	no	no
29	7	P c 1 1	P -2xc	a, cell choice 3	primitive	2	no	no	no
30	8	C 1 m 1	C -2y	b, cell choice 1	c	4	no	no	no
31	8	A 1 m 1	A -2y	b, cell choice 2	a	4	no	no	no
32	8	I 1 m 1	I -2y	b, cell choice 3	body	4	no	no	no
33	8	A 1 1 m	A -2	c, cell choice 1	a	4	no	no	no
34	8	B 1 1 m	B -2	c, cell choice 2	b	4	no	no	no
35	8	I 1 1 m	I -2	c, cell choice 3	body	4	no	no	no
36	8	B m 1 1	B -2x	a, cell choice 1	b	4	no	no	no
37	8	B m 1 1	C -2x	a, cell choice 2	c	4	no	no	no
38	8	I m 1 1	I -2x	a, cell choice 3	body	4	no	no	no
39	9	C 1 c 1	C -2yc	b, cell choice 1	c	4	no	no	no
40	9	A 1 n 1	A -2yab	b, cell choice 2	a	4	no	no	no
41	9	I 1 a 1	I -2ya	b, cell choice 3	body	4	no	no	no
42	9	A 1 a 1	A -2ya	-b, cell choice 1	a	4	no	no	no
43	9	C 1 n 1	C -2yac	-b, cell choice 2	c	4	no	no	no
44	9	I 1 c 1	I -2yc	-b, cell choice 3	body	4	no	no	no

45	9	A 1 1 a	A -2a	c, cell choice 1	a	4	no	no	no
46	9	B 1 1 n	B -2ab	c, cell choice 2	b	4	no	no	no
47	9	I 1 1 b	I -2b	c, cell choice 3	body	4	no	no	no
48	9	B 1 1 b	B -2b	-c, cell choice 1	b	4	no	no	no
49	9	A 1 1 n	A -2ab	-c, cell choice 2	a	4	no	no	no
50	9	I 1 1 a	I -2a	-c, cell choice 3	body	4	no	no	no
51	9	B b 1 1	B -2xb	a, cell choice 1	b	4	no	no	no
52	9	C n 1 1	C -2xac	a, cell choice 2	c	4	no	no	no
53	9	I c 1 1	I -2xc	a, cell choice 3	body	4	no	no	no
54	9	C c 1 1	C -2xc	-a, cell choice 1	c	4	no	no	no
55	9	B n 1 1	B -2xab	-a, cell choice 2	b	4	no	no	no
56	9	I b 1 1	I -2xb	-a, cell choice 3	body	4	no	no	no
57	10	P 1 2/m 1	-P 2y	unique axis b	primitive	4	yes	no	no
58	10	P 1 1 2/m	-P 2	unique axis c	primitive	4	yes	no	no
59	10	P 2/m 1 1	-P 2x	unique axis a	primitive	4	yes	no	no
60	11	P 1 21/m 1	-P 2yb	unique axis b	primitive	4	yes	no	no
61	11	P 1 1 21/m	-P 2c	unique axis c	primitive	4	yes	no	no
62	11	P 21/m 1 1	-P 2xa	unique axis a	primitive	4	yes	no	no
63	12	C 1 2/m 1	-C 2y	b, cell choice 1	c	8	yes	no	no
64	12	A 1 2/m 1	-A 2y	b, cell choice 2	a	8	yes	no	no
65	12	I 1 2/m 1	-I 2y	b, cell choice 3	body	8	yes	no	no
66	12	A 1 1 2/m	-A 2	c, cell choice 1	a	8	yes	no	no
67	12	B 1 1 2/m	-B 2	c, cell choice 2	b	8	yes	no	no
68	12	I 1 1 2/m	-I 2	c, cell choice 3	body	8	yes	no	no
69	12	B 2/m 1 1	-B 2x	a, cell choice 1	b	8	yes	no	no
70	12	C 2/m 1 1	-C 2x	a, cell choice 2	c	8	yes	no	no
71	12	I 2/m 1 1	-I 2x	a, cell choice 3	body	8	yes	no	no
72	13	P 1 2/c 1	-P 2yc	b, cell choice 1	primitive	4	yes	no	no
73	13	P 1 2/n 1	-P 2yac	b, cell choice 2	primitive	4	yes	no	no
74	13	P 1 2/a 1	-P 2ya	b, cell choice 3	primitive	4	yes	no	no
75	13	P 1 1 2/a	-P 2a	c, cell choice 1	primitive	4	yes	no	no
76	13	P 1 1 2/n	-P 2ab	c, cell choice 2	primitive	4	yes	no	no
77	13	P 1 1 2/b	-P 2b	c, cell choice 3	primitive	4	yes	no	no
78	13	P 2/b 1 1	-P 2xb	a, cell choice 1	primitive	4	yes	no	no
79	13	P 2/n 1 1	-P 2xbc	a, cell choice 2	primitive	4	yes	no	no
80	13	P 2/c 1 1	-P 2xc	a, cell choice 3	primitive	4	yes	no	no
81	14	P 1 21/c 1	-P 2ybc	b, cell choice 1	primitive	4	yes	no	no
82	14	P 1 21/n 1	-P 2yn	b, cell choice 2	primitive	4	yes	no	no
83	14	P 1 21/a 1	-P 2yab	b, cell choice 3	primitive	4	yes	no	no
84	14	P 1 1 21/a	-P 2ac	c, cell choice 1	primitive	4	yes	no	no
85	14	P 1 1 21/n	-P 2n	c, cell choice 2	primitive	4	yes	no	no
86	14	P 1 1 21/b	-P 2bc	c, cell choice 3	primitive	4	yes	no	no
87	14	P 21/b 1 1	-P 2xab	a, cell choice 1	primitive	4	yes	no	no
88	14	P 21/n 1 1	-P 2xn	a, cell choice 2	primitive	4	yes	no	no
89	14	P 21/c 1 1	-P 2xac	a, cell choice 3	primitive	4	yes	no	no
90	15	C 1 2/c 1	-C 2yc	b, cell choice 1	c	8	yes	no	no
91	15	A 1 2/n 1	-A 2yab	b, cell choice 2	a	8	yes	no	no
92	15	I 1 2/a 1	-I 2ya	b, cell choice 3	body	8	yes	no	no
93	15	A 1 2/a 1	-A 2ya	-b, cell choice 1	a	8	yes	no	no
94	15	C 1 2/n 1	-C 2yac	-b, cell choice 2	c	8	yes	no	no
95	15	I 1 2/c 1	-I 2yc	-b, cell choice 3	body	8	yes	no	no
96	15	A 1 1 2/a	-A 2a	c, cell choice 1	a	8	yes	no	no
97	15	B 1 1 2/n	-B 2ab	c, cell choice 2	b	8	yes	no	no
98	15	I 1 1 2/b	-I 2b	c, cell choice 3	body	8	yes	no	no
99	15	B 1 1 2/b	-B 2b	-c, cell choice 1	b	8	yes	no	no
100	15	A 1 1 2/n	-A 2ab	-c, cell choice 2	a	8	yes	no	no

101	15	I 1 1 2/a	-I 2a	-c, cell choice 3	body	8	yes	no	no
102	15	B 2/b 1 1	-B 2xb	a, cell choice 1	b	8	yes	no	no
103	15	C 2/n 1 1	-C 2xac	a, cell choice 2	c	8	yes	no	no
104	15	I 2/c 1 1	-I 2xc	a, cell choice 3	body	8	yes	no	no
105	15	C 2/c 1 1	-C 2xc	-a, cell choice 1	c	8	yes	no	no
106	15	B 2/n 1 1	-B 2xab	-a, cell choice 2	b	8	yes	no	no
107	15	I 2/b 1 1	-I 2xb	-a, cell choice 3	body	8	yes	no	no

Table 2.2: Monoclinic spacegroup information.

orthorhombic									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
108	16	P 2 2 2	P 2 2	cell choice 1	primitive	4	yes	yes	no
109	17	P 2 2 21	P 2c 2	abc	primitive	4	yes	yes	no
110	17	P 2 1 2 2	P 2a 2a	cab	primitive	4	yes	yes	no
111	17	P 2 21 2	P 2 2b	bca	primitive	4	yes	yes	no
112	18	P 2 1 21 2	P 2 2ab	abc	primitive	4	yes	yes	no
113	18	P 2 21 21	P 2bc 2	cab	primitive	4	yes	yes	no
114	18	P 2 1 2 21	P 2ac 2ac	bca	primitive	4	yes	yes	no
115	19	P 2 1 21 21	P 2ac 2ab	cell choice 1	primitive	4	yes	yes	no
116	20	C 2 2 21	C 2c 2	abc	c	8	yes	yes	no
117	20	A 2 1 2 2	A 2a 2a	cab	a	8	yes	yes	no
118	20	B 2 21 2	B 2 2b	bca	b	8	yes	yes	no
119	21	C 2 2 2	C 2 2	abc	c	8	no	yes	no
120	21	A 2 2 2	A 2 2	cab	a	8	no	yes	no
121	21	B 2 2 2	B 2 2	bca	b	8	no	yes	no
122	22	F 2 2 2	F 2 2	cell choice 1	face	16	no	yes	no
123	23	I 2 2 2	I 2 2	cell choice 1	body	8	no	yes	no
124	24	I 21 21 21	I 2b 2c	cell choice 1	body	8	no	yes	no
125	25	P m m 2	P 2 -2	abc	primitive	4	no	no	no
126	25	P 2 m m	P -2 2	cab	primitive	4	no	no	no
127	25	P m 2 m	P -2 -2	bca	primitive	4	no	no	no
128	26	P m c 21	P 2c -2	abc	primitive	4	no	no	no
129	26	P c m 21	P 2c -2c	ba-c	primitive	4	no	no	no
130	26	P 21 m a	P -2a 2a	cab	primitive	4	no	no	no
131	26	P 21 a m	P -2 2a	-cba	primitive	4	no	no	no
132	26	P b 21 m	P -2 -2b	bca	primitive	4	no	no	no
133	26	P m 21 b	P -2b -2	a-cb	primitive	4	no	no	no
134	27	P c c 2	P 2 -2c	abc	primitive	4	no	no	no
135	27	P 2 a a	P -2a 2	cab	primitive	4	no	no	no
136	27	P b 2 b	P -2b -2b	bca	primitive	4	no	no	no
137	28	P m a 2	P 2 -2a	abc	primitive	4	no	no	no
138	28	P b m 2	P 2 -2b	ba-c	primitive	4	no	no	no
139	28	P 2 m b	P -2b 2	cab	primitive	4	no	no	no
140	28	P 2 c m	P -2c 2	-cba	primitive	4	no	no	no
141	28	P c 2 m	P -2c -2c	bca	primitive	4	no	no	no
142	28	P m 2 a	P -2a -2a	a-cb	primitive	4	no	no	no
143	29	P c a 21	P 2c -2ac	abc	primitive	4	no	no	no
144	29	P b c 21	P 2c -2b	ba-c	primitive	4	no	no	no
145	29	P 21 a b	P -2b 2a	cab	primitive	4	no	no	no
146	29	P 21 c a	P -2ac 2a	-cba	primitive	4	no	no	no
147	29	P c 21 b	P -2bc -2c	bca	primitive	4	no	no	no
148	29	P b 21 a	P -2a -2ab	a-cb	primitive	4	no	no	no
149	30	P n c 2	P 2 -2bc	abc	primitive	4	no	no	no
150	30	P c n 2	P 2 -2ac	ba-c	primitive	4	no	no	no

151	30	P 2 n a	P -2ac 2	cab	primitive	4	no	no	no
152	30	P 2 a n	P -2ab 2	-cba	primitive	4	no	no	no
153	30	P b 2 n	P -2ab -2ab	bca	primitive	4	no	no	no
154	30	P n 2 b	P -2bc -2bc	a-cb	primitive	4	no	no	no
155	31	P m n 21	P 2ac -2	abc	primitive	4	no	no	no
156	31	P n m 21	P 2bc -2bc	ba-c	primitive	4	no	no	no
157	31	P 21 m n	P -2ab 2ab	cab	primitive	4	no	no	no
158	31	P 21 n m	P -2 2ac	-cba	primitive	4	no	no	no
159	31	P n 21 m	P -2 -2bc	bca	primitive	4	no	no	no
160	31	P m 21 n	P -2ab -2	a-cb	primitive	4	no	no	no
161	32	P b a 2	P 2 -2ab	abc	primitive	4	no	no	no
162	32	P 2 c b	P -2bc 2	cab	primitive	4	no	no	no
163	32	P c 2 a	P -2ac -2ac	bca	primitive	4	no	no	no
164	33	P n a 21	P 2c -2n	abc	primitive	4	no	no	no
165	33	P b n 21	P 2c -2ab	ba-c	primitive	4	no	no	no
166	33	P 21 n b	P -2bc 2a	cab	primitive	4	no	no	no
167	33	P 21 c n	P -2n 2a	-cba	primitive	4	no	no	no
168	33	P c 21 n	P -2n -2ac	bca	primitive	4	no	no	no
169	33	P n 21 a	P -2ac -2n	a-cb	primitive	4	no	no	no
170	34	P n n 2	P 2 -2n	abc	primitive	4	no	no	no
171	34	P 2 n n	P -2n 2	cab	primitive	4	no	no	no
172	34	P n 2 n	P -2n -2n	bca	primitive	4	no	no	no
173	35	C m m 2	C 2 -2	abc	c	8	no	no	no
174	35	A 2 m m	A -2 2	cab	a	8	no	no	no
175	35	B m 2 m	B -2 -2	bca	b	8	no	no	no
176	36	C m c 21	C 2c -2	abc	c	8	no	no	no
177	36	C c m 21	C 2c -2c	ba-c	c	8	no	no	no
178	36	A 21 m a	A -2a 2a	cab	a	8	no	no	no
179	36	A 21 a m	A -2 2a	-cba	a	8	no	no	no
180	36	B b 21 m	B -2 -2b	bca	b	8	no	no	no
181	36	B m 21 b	B -2b -2	a-cb	b	8	no	no	no
182	37	C c c 2	C 2 -2c	abc	c	8	no	no	no
183	37	A 2 a a	A -2a 2	cab	a	8	no	no	no
184	37	B b 2 b	B -2b -2b	bca	b	8	no	no	no
185	38	A m m 2	A 2 -2	abc	a	8	no	no	no
186	38	B m m 2	B 2 -2	ba-c	b	8	no	no	no
187	38	B 2 m m	B -2 2	cab	b	8	no	no	no
188	38	C 2 m m	C -2 2	-cba	c	8	no	no	no
189	38	C m 2 m	C -2 -2	bca	c	8	no	no	no
190	38	A m 2 m	A -2 -2	a-cb	a	8	no	no	no
191	39	A b m 2	A 2 -2b	abc	a	8	no	no	no
192	39	B m a 2	B 2 -2a	ba-c	b	8	no	no	no
193	39	B 2 c m	B -2a 2	cab	b	8	no	no	no
194	39	C 2 m b	C -2a 2	-cba	c	8	no	no	no
195	39	C m 2 a	C -2a -2a	bca	c	8	no	no	no
196	39	A c 2 m	A -2b -2b	a-cb	a	8	no	no	no
197	40	A m a 2	A 2 -2a	abc	a	8	no	no	no
198	40	B b m 2	B 2 -2b	ba-c	b	8	no	no	no
199	40	B 2 m b	B -2b 2	cab	b	8	no	no	no
200	40	C 2 c m	C -2c 2	-cba	c	8	no	no	no
201	40	C c 2 m	C -2c -2c	bca	c	8	no	no	no
202	40	A m 2 a	A -2a -2a	a-cb	a	8	no	no	no
203	41	A b a 2	A 2 -2ab	abc	a	8	no	no	no
204	41	B b a 2	B 2 -2ab	ba-c	b	8	no	no	no
205	41	B 2 c b	B -2ab 2	cab	b	8	no	no	no
206	41	C 2 c b	C -2ac 2	-cba	c	8	no	no	no

207	41	C c 2 a	C -2ac -2ac	bca	c	8	no	no	no
208	41	A c 2 a	A -2ab -2ab	a-cb	a	8	no	no	no
209	42	F m m 2	F 2 -2	abc	face	16	no	no	no
210	42	F 2 m m	F -2 2	cab	face	16	no	no	no
211	42	F m 2 m	F -2 -2	bca	face	16	no	no	no
212	43	F d d 2	F 2 -2d	abc	face	16	no	no	no
213	43	F 2 d d	F -2d 2	cab	face	16	no	no	no
214	43	F d 2 d	F -2d -2d	bca	face	16	no	no	no
215	44	I m m 2	I 2 -2	abc	body	8	no	no	no
216	44	I 2 m m	I -2 2	cab	body	8	no	no	no
217	44	I m 2 m	I -2 -2	bca	body	8	no	no	no
218	45	I b a 2	I 2 -2c	abc	body	8	no	no	no
219	45	I 2 c b	I -2a 2	cab	body	8	no	no	no
220	45	I c 2 a	I -2b -2b	bca	body	8	no	no	no
221	46	I m a 2	I 2 -2a	abc	body	8	no	no	no
222	46	I b m 2	I 2 -2b	ba-c	body	8	no	no	no
223	46	I 2 m b	I -2b 2	cab	body	8	no	no	no
224	46	I 2 c m	I -2c 2	-cba	body	8	no	no	no
225	46	I c 2 m	I -2c -2c	bca	body	8	no	no	no
226	46	I m 2 a	I -2a -2a	a-cb	body	8	no	no	no
227	47	P 2/m 2/m 2/m	-P 2 2	cell choice 1	primitive	8	yes	no	no
228	48	P 2/n 2/n 2/n:1	P 2 2 -1n	cell choice 1	primitive	8	yes	no	no
229	48	P 2/n 2/n 2/n:2	-P 2ab 2bc	cell choice 2	primitive	8	yes	no	no
230	49	P 2/c 2/c 2/m	-P 2 2c	abc	primitive	8	yes	no	no
231	49	P 2/m 2/a 2/a	-P 2a 2	cab	primitive	8	yes	no	no
232	49	P 2/b 2/m 2/b	-P 2b 2b	bca	primitive	8	yes	no	no
233	50	P 2/b 2/a 2/n:1	P 2 2 -1ab	cell choice 1	primitive	8	yes	no	no
234	50	P 2/b 2/a 2/n:2	-P 2ab 2b	cell choice 2	primitive	8	yes	no	no
235	50	P 2/n 2/c 2/b:1	P 2 2 -1bc	cab	primitive	8	yes	no	no
236	50	P 2/n 2/c 2/b:2	-P 2b 2bc	cab, cell choice 2	primitive	8	yes	no	no
237	50	P 2/c 2/n 2/a:1	P 2 2 -1ac	bca	primitive	8	yes	no	no
238	50	P 2/c 2/n 2/a:2	-P 2a 2c	bca, cell choice 2	primitive	8	yes	no	no
239	51	P 21/m 2/m 2/a	-P 2a 2a	abc	primitive	8	yes	no	no
240	51	P 2/m 21/m 2/b	-P 2b 2	ba-c	primitive	8	yes	no	no
241	51	P 2/b 21/m 2/m	-P 2 2b	cab	primitive	8	yes	no	no
242	51	P 2/c 2/m 21/m	-P 2c 2c	-cba	primitive	8	yes	no	no
243	51	P 2/m 2/c 21/m	-P 2c 2	bca	primitive	8	yes	no	no
244	51	P 21/m 2/a 2/m	-P 2 2a	a-cb	primitive	8	yes	no	no
245	52	P 2/n 21/n 2/a	-P 2a 2bc	abc	primitive	8	yes	no	no
246	52	P 21/n 2/n 2/b	-P 2b 2n	ba-c	primitive	8	yes	no	no
247	52	P 2/b 2/n 21/n	-P 2n 2b	cab	primitive	8	yes	no	no
248	52	P 2/c 21/n 2/n	-P 2ab 2c	-cba	primitive	8	yes	no	no
249	52	P 21/n 2/c 2/n	-P 2ab 2n	bca	primitive	8	yes	no	no
250	52	P 2/n 2/a 21/n	-P 2n 2bc	a-cb	primitive	8	yes	no	no
251	53	P 2/m 2/n 21/a	-P 2ac 2	abc	primitive	8	yes	no	no
252	53	P 2/n 2/m 21/b	-P 2bc 2bc	ba-c	primitive	8	yes	no	no
253	53	P 21/b 2/m 2/n	-P 2ab 2ab	cab	primitive	8	yes	no	no
254	53	P 21/c 2/n 2/m	-P 2 2ac	-cba	primitive	8	yes	no	no
255	53	P 2/n 21/c 2/m	-P 2 2bc	bca	primitive	8	yes	no	no
256	53	P 2/m 21/a 2/n	-P 2ab 2	a-cb	primitive	8	yes	no	no
257	54	P 21/c 2/c 2/a	-P 2a 2ac	abc	primitive	8	yes	no	no
258	54	P 2/c 21/c 2/b	-P 2b 2c	ba-c	primitive	8	yes	no	no
259	54	P 2/b 21/a 2/a	-P 2a 2b	cab	primitive	8	yes	no	no
260	54	P 2/c 2/a 21/a	-P 2ac 2c	-cba	primitive	8	yes	no	no
261	54	P 2/b 2/c 21/b	-P 2bc 2b	bca	primitive	8	yes	no	no
262	54	P 21/b 2/a 2/b	-P 2b 2ab	a-cb	primitive	8	yes	no	no

263	55	P 21/b 21/a 2/m	-P 2 2ab	abc	primitive	8	yes	no	no
264	55	P 2/m 21/c 21/b	-P 2bc 2	cab	primitive	8	yes	no	no
265	55	P 21/c 2/m 21/a	-P 2ac 2ac	bca	primitive	8	yes	no	no
266	56	P 21/c 21/c 2/n	-P 2ab 2ac	abc	primitive	8	yes	no	no
267	56	P 2/n 21/a 21/a	-P 2ac 2bc	cab	primitive	8	yes	no	no
268	56	P 21/b 2/n 21/b	-P 2bc 2ab	bca	primitive	8	yes	no	no
269	57	P 2/b 21/c 21/m	-P 2c 2b	abc	primitive	8	yes	no	no
270	57	P 21/c 2/a 21/m	-P 2c 2ac	ba-c	primitive	8	yes	no	no
271	57	P 21/m 2/c 21/a	-P 2ac 2a	cab	primitive	8	yes	no	no
272	57	P 21/m 21/a 2/b	-P 2b 2a	-cba	primitive	8	yes	no	no
273	57	P 21/b 21/m 2/a	-P 2a 2ab	bca	primitive	8	yes	no	no
274	57	P 2/c 21/m 21/b	-P 2bc 2c	a-cb	primitive	8	yes	no	no
275	58	P 21/n 21n 2/m	-P 2 2n	abc	primitive	8	yes	no	no
276	58	P 2/m 21/n 21/n	-P 2n 2	cab	primitive	8	yes	no	no
277	58	P 21/n 2/m 21/n	-P 2n 2n	bca	primitive	8	yes	no	no
278	59	P 21/m 21/m 2/n:1	P 2 2ab -1ab	cell choice 1	primitive	8	yes	no	no
279	59	P 21/m 21/m 2/n:2	-P 2ab 2a	cell choice 2	primitive	8	yes	no	no
280	59	P 2/n 21/m 21/m:1	P 2bc 2 -1bc	cab	primitive	8	yes	no	no
281	59	P 2/n 21/m 21/m:2	-P 2c 2bc	cab, cell choice 2	primitive	8	yes	no	no
282	59	P 21/m 2/n 21/m:1	P 2ac 2ac -1ac	bca	primitive	8	yes	no	no
283	59	P 21/m 2/n 21/m:2	-P 2c 2a	bca, cell choice 2	primitive	8	yes	no	no
284	60	P 21/b 2/c 21/n	-P 2n 2ab	abc	primitive	8	yes	no	no
285	60	P 2/c 21/a 21/n	-P 2n 2c	ba-c	primitive	8	yes	no	no
286	60	P 21/n 21/a 2/b	-P 2a 2n	cab	primitive	8	yes	no	no
287	60	P 21/n 2/a 21/b	-P 2bc 2n	-cba	primitive	8	yes	no	no
288	60	P 2/b 21/n 21/a	-P 2ac 2b	bca	primitive	8	yes	no	no
289	60	P 21/c 21/n 2/b	-P 2b 2ac	a-cb	primitive	8	yes	no	no
290	61	P 21/b 21/c 21/a	-P 2ac 2ab	abc	primitive	8	yes	no	no
291	61	P 21/c 21/a 21/b	-P 2bc 2ac	ba-c	primitive	8	yes	no	no
292	62	P 21/n 21/m 21/a	-P 2ac 2n	abc	primitive	8	yes	no	no
293	62	P 21/m 21/n 21/b	-P 2bc 2a	ba-c	primitive	8	yes	no	no
294	62	P 21/b 21/n 21/m	-P 2c 2ab	cab	primitive	8	yes	no	no
295	62	P 21/c 21/m 21/n	-P 2n 2ac	-cba	primitive	8	yes	no	no
296	62	P 21/m 21/c 21/n	-P 2n 2a	bca	primitive	8	yes	no	no
297	62	P 21/n 21/a 21/m	-P 2c 2n	a-cb	primitive	8	yes	no	no
298	63	C 2/m 2/c 21/m	-C 2c 2	abc	c	16	yes	no	no
299	63	C 2/c 2/m 21/m	-C 2c 2c	ba-c	c	16	yes	no	no
300	63	A 21/m 2/m 2/a	-A 2a 2a	cab	a	16	yes	no	no
301	63	A 21/m 2/a 2/m	-A 2 2a	-cba	a	16	yes	no	no
302	63	B 2/b 21/m 2/m	-B 2 2b	bca	b	16	yes	no	no
303	63	B 2/m 21/m 2/b	-B 2b 2	a-cb	b	16	yes	no	no
304	64	C 2/m 2/c 21/a	-C 2ac 2	abc	c	16	yes	no	no
305	64	C 2/c 2/m 21/b	-C 2ac 2ac	ba-c	c	16	yes	no	no
306	64	A 21/b 2/m 2/a	-A 2ab 2ab	cab	a	16	yes	no	no
307	64	A 21/c 2/a 2/m	-A 2 2ab	-cba	a	16	yes	no	no
308	64	B 2/b 21/c 2/m	-B 2 2ab	bca	b	16	yes	no	no
309	64	B 2/m 21/a 2/b	-B 2ab 2	a-cb	b	16	yes	no	no
310	65	C 2/m 2/m 2/m	-C 2 2	abc	c	16	yes	no	no
311	65	A 2/m 2/m 2/m	-A 2 2	cab	a	16	yes	no	no
312	65	B 2/m 2/m 2/m	-B 2 2	bca	b	16	yes	no	no
313	66	C 2/c 2/c 2/m	-C 2 2c	abc	c	16	yes	no	no
314	66	A 2/m 2/a 2/a	-A 2a 2	cab	a	16	yes	no	no
315	66	B 2/b 2/m 2/b	-B 2b 2b	bca	b	16	yes	no	no
316	67	C 2/m 2/m 2/a	-C 2a 2	abc	c	16	yes	no	no
317	67	C 2/m 2/m 2/b	-C 2a 2a	ba-c	c	16	yes	no	no
318	67	A 2/b 2/m 2/m	-A 2b 2b	cab	a	16	yes	no	no

319	67	A 2/c 2/m 2/m	-A 2 2b	-cba	a	16	yes	no	no
320	67	B 2/m 2/c 2/m	-B 2 2a	bca	b	16	yes	no	no
321	67	B 2/m 2/a 2/m	-B 2a 2	a-cb	b	16	yes	no	no
322	68	C 2/c 2/c 2/a:1	C 2 2 -1ac	cell choice 1	c	16	yes	no	no
323	68	C 2/c 2/c 2/a:2	-C 2a 2ac	cell choice 2	c	16	yes	no	no
324	68	C 2/c 2/c 2/b:1	C 2 2 -1ac	ba-c	c	16	yes	no	no
325	68	C 2/c 2/c 2/b:2	-C 2a 2c	ba-c, cell choice 2	c	16	yes	no	no
326	68	A 2/b 2/a 2/a:1	A 2 2 -1ab	cab	a	16	yes	no	no
327	68	A 2/b 2/a 2/a:2	-A 2a 2b	cab, cell choice 2	a	16	yes	no	no
328	68	A 2/c 2/a 2/a:1	A 2 2 -1ab	-cba	a	16	yes	no	no
329	68	A 2/c 2/a 2/a:2	-A 2ab 2b	-cba, cell choice 2	a	16	yes	no	no
330	68	B 2/b 2/c 2/b:1	B 2 2 -1ab	bca	b	16	yes	no	no
331	68	B 2/b 2/c 2/b:2	-B 2ab 2b	bca, cell choice 2	b	16	yes	no	no
332	68	B 2/b 2/a 2/b:1	B 2 2 -1ab	a-cb	b	16	yes	no	no
333	68	B 2/b 2/a 2/b:2	-B 2b 2ab	a-cb, cell choice 2	b	16	yes	no	no
334	69	F 2/m 2/m 2/m	-F 2 2	cell choice 1	face	32	yes	no	no
335	70	F 2/d 2/d 2/d:1	F 2 2 -1d	cell choice 1	face	32	yes	no	no
336	70	F 2/d 2/d 2/d:2	-F 2uv 2vw	cell choice 2	face	32	yes	no	no
337	71	I 2/m 2/m 2/m	-I 2 2	cell choice 1	body	16	yes	no	no
338	72	I 2/b 2/a 2/m	-I 2 2c	abc	body	16	yes	no	no
339	72	I 2/m 2/c 2/b	-I 2a 2	cab	body	16	yes	no	no
340	72	I 2/c 2/m 2/a	-I 2b 2b	bca	body	16	yes	no	no
341	73	I 21/b 21/c 21/a	-I 2b 2c	abc	body	16	yes	no	no
342	73	I 21/c 21/a 21/b	-I 2a 2b	ba-c	body	16	yes	no	no
343	74	I 21/m 21/m 21/a	-I 2b 2	abc	body	16	yes	no	no
344	74	I 21/m 21/m 21/b	-I 2a 2a	ba-c	body	16	yes	no	no
345	74	I 21/b 21/m 21/m	-I 2c 2c	cab	body	16	yes	no	no
346	74	I 21/c 21/m 21/m	-I 2 2b	-cba	body	16	yes	no	no
347	74	I 21/m 21/c 21/m	-I 2 2a	bca	body	16	yes	no	no
348	74	I 21/m 21/a 21/m	-I 2c 2	a-cb	body	16	yes	no	no

Table 2.3: Orthorhombic spacegroup information.

tetragonal									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantiomeric
349	75	P 4	P 4	cell choice 1	primitive	4	no	yes	no
350	76	P 41	P 4w	cell choice 1	primitive	4	no	yes	yes
351	77	P 42	P 4c	cell choice 1	primitive	4	no	yes	no
352	78	P 43	P 4cw	cell choice 1	primitive	4	no	yes	yes
353	79	I 4	I 4	cell choice 1	body	8	no	yes	no
354	80	I 41	I 4bw	cell choice 1	body	8	no	yes	no
355	81	P -4	P -4	cell choice 1	primitive	4	no	no	no
356	82	I -4	I -4	cell choice 1	body	8	no	no	no
357	83	P 4/m	-P 4	cell choice 1	primitive	8	yes	no	no
358	84	P 42/m	-P 4c	cell choice 1	primitive	8	yes	no	no
359	85	P 4/n:1	P 4ab -1ab	cell choice 1	primitive	8	yes	no	no
360	85	P 4/n:2	-P 4a	cell choice 2	primitive	8	yes	no	no
361	86	P 42/n:1	P 4n -1n	cell choice 1	primitive	8	yes	no	no
362	86	P 42/n:2	-P 4bc	cell choice 2	primitive	8	yes	no	no
363	87	I 4/m	-I 4	cell choice 1	body	16	yes	no	no
364	88	I 41/a:1	I 4bw -1bw	cell choice 1	body	16	yes	no	no
365	88	I 41/a:2	-I 4ad	cell choice 2	body	16	yes	no	no
366	89	P 4 2 2	P 4 2	cell choice 1	primitive	8	no	yes	no
367	90	P 4 21 2	P 4ab 2ab	cell choice 1	primitive	8	no	yes	no
368	91	P 41 2 2	P 4w 2c	cell choice 1	primitive	8	no	yes	yes

369	92	P 41 21 2	P 4abw 2nw	cell choice 1	primitive	8	no	yes	yes
370	93	P 42 2 2	P 4c 2	cell choice 1	primitive	8	no	yes	no
371	94	P 42 21 2	P 4n 2n	cell choice 1	primitive	8	no	yes	no
372	95	P 43 2 2	P 4cw 2c	cell choice 1	primitive	8	no	yes	yes
373	96	P 43 21 2	P 4nw 2abw	cell choice 1	primitive	8	no	yes	yes
374	97	I 4 2 2	I 4 2	cell choice 1	body	16	no	yes	no
375	98	I 41 2 2	I 4bw 2bw	cell choice 1	body	16	no	yes	no
376	99	P 4 m m	P 4 -2	cell choice 1	primitive	8	no	no	no
377	100	P 4 b n	P 4 -2ab	cell choice 1	primitive	8	no	no	no
378	101	P 42 c m	P 4c -2c	cell choice 1	primitive	8	no	no	no
379	102	P 42 n m	P 4n -2n	cell choice 1	primitive	8	no	no	no
380	103	P 4 c c	P 4 -2c	cell choice 1	primitive	8	no	no	no
381	104	P 4 n c	P 4 -2n	cell choice 1	primitive	8	no	no	no
382	105	P 42 m c	P 4c -2	cell choice 1	primitive	8	no	no	no
383	106	P 42 b c	P 4c -2ab	cell choice 1	primitive	8	no	no	no
384	107	I 4 m m	I 4 -2	cell choice 1	body	16	no	no	no
385	108	I 4 c m	I 4 -2c	cell choice 1	body	16	no	no	no
386	109	I 41 m d	I 4bw -2	cell choice 1	body	16	no	no	no
387	110	I 41 c d	I 4bw -2c	cell choice 1	body	16	no	no	no
388	111	P -4 2 m	P -4 2	cell choice 1	primitive	8	no	no	no
389	112	P -4 2 c	P -4 2c	cell choice 1	primitive	8	no	no	no
390	113	P -4 21 m	P -4 2ab	cell choice 1	primitive	8	no	no	no
391	114	P -4 21 c	P -4 2n	cell choice 1	primitive	8	no	no	no
392	115	P -4 m 2	P -4 -2	cell choice 1	primitive	8	no	no	no
393	116	P -4 c 2	P -4 -2c	cell choice 1	primitive	8	no	no	no
394	117	P -4 b 2	P -4 -2ab	cell choice 1	primitive	8	no	no	no
395	118	P -4 n 2	P -4 -2n	cell choice 1	primitive	8	no	no	no
396	119	I -4 m 2	I -4 -2	cell choice 1	body	16	no	no	no
397	120	I -4 c 2	I -4 -2c	cell choice 1	body	16	no	no	no
398	121	I -4 2 m	I -4 2	cell choice 1	body	16	no	no	no
399	122	I -4 2 d	I -4 2bw	cell choice 1	body	16	no	no	no
400	123	P 4/m 2/m 2/m	-P 4 2	cell choice 1	primitive	16	yes	no	no
401	124	P 4/m 2/c 2/c	-P 4 2c	cell choice 1	primitive	16	yes	no	no
402	125	P 4/n 2/b 2/m:1	P 4 2 -1ab	cell choice 1	primitive	16	yes	no	no
403	125	P 4/n 2/b 2/m:2	-P 4a 2b	cell choice 2	primitive	16	yes	no	no
404	126	P 4/n 2/n 2/c:1	P 4 2 -1n	cell choice 1	primitive	16	yes	no	no
405	126	P 4/n 2/n 2/c:2	-P 4a 2bc	cell choice 2	primitive	16	yes	no	no
406	127	P 4/m 21/b 2/m	-P 4 2ab	cell choice 1	primitive	16	yes	no	no
407	128	P 4/m 21/n 2/c	-P 4 2n	cell choice 1	primitive	16	yes	no	no
408	129	P 4/n 21/m 2/m:1	P 4ab 2ab -1ab	cell choice 1	primitive	16	yes	no	no
409	129	P 4/n 21/m 2/m:2	-P 4a 2a	cell choice 2	primitive	16	yes	no	no
410	130	P 4/n 21/c 2/c:1	P 4ab 2n -1ab	cell choice 1	primitive	16	yes	no	no
411	130	P 4/n 21/c 2/c:2	-P 4a 2ac	cell choice 2	primitive	16	yes	no	no
412	131	P 42/m 2/m 2/c	-P 4c 2	cell choice 1	primitive	16	yes	no	no
413	132	P 42/m 2/c 2/m	-P 4c 2c	cell choice 1	primitive	16	yes	no	no
414	133	P 42/n 2/b 2/c:1	P 4n 2c -1n	cell choice 1	primitive	16	yes	no	no
415	133	P 42/n 2/b 2/c:2	-P 4ac 2b	cell choice 2	primitive	16	yes	no	no
416	134	P 42/n 2/n 2/m:1	P 4n 2 -1n	cell choice 1	primitive	16	yes	no	no
417	134	P 42/n 2/n 2/m:2	-P 4ac 2bc	cell choice 2	primitive	16	yes	no	no
418	135	P 42/m 21/b 2/c	-P 4c 2ab	cell choice 1	primitive	16	yes	no	no
419	136	P 42/m 21/n 2/m	-P 4n 2n	cell choice 1	primitive	16	yes	no	no
420	137	P 42/n 21/m 2/c:1	P 4n 2n -1n	cell choice 1	primitive	16	yes	no	no
421	137	P 42/n 21/m 2/c:2	-P 4ac 2a	cell choice 2	primitive	16	yes	no	no
422	138	P 42/n 21/c 2/m:1	P 4n 2ab -1n	cell choice 1	primitive	16	yes	no	no
423	138	P 42/n 21/c 2/m:2	-P 4ac 2ac	cell choice 2	primitive	16	yes	no	no
424	139	I 4/m 2/m 2/m	-I 4 2	cell choice 1	primitive	32	yes	no	no

425	140	I 4/m 2/c 2/m	-I 4 2c	cell choice 1	primitive	32	yes	no	no
426	141	I 41/a 2/m 2/d:1	I 4bw 2bw -1bw	cell choice 1	body	32	yes	no	no
427	141	I 41/a 2/m 2/d:2	-I 4bd 2	cell choice 2	body	32	yes	no	no
428	142	I 41/a 2/c 2/d:1	I 4bw 2aw -1bw	cell choice 1	body	32	yes	no	no
429	142	I 41/a 2/c 2/d:2	-I 4bd 2c	cell choice 2	body	32	yes	no	no

Table 2.4: Tetragonal spacegroup information.

trigonal									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
430	143	P 3	P 3	cell choice 1	primitive	3	no	yes	no
431	144	P 31	P 31	cell choice 1	primitive	3	no	yes	yes
432	145	P 32	P 32	cell choice 1	primitive	3	no	yes	no
433	146	R 3:H	R 3	hexagonal	rhombohedral	9	no	yes	no
434	146	R 3:R	P 3*	Rhombohedral	primitive	3	no	yes	no
435	147	P -3	-P 3	cell choice 1	primitive	6	yes	no	no
436	148	R -3:H	-R 3	hexagonal	rhombohedral	18	yes	no	no
437	148	R -3:R	-P 3*	Rhombohedral	primitive	6	yes	no	no
438	149	P 3 1 2	P 3 2	cell choice 1	primitive	6	no	yes	no
439	150	P 3 2 1	P 3 2"	cell choice 1	primitive	6	no	yes	no
440	151	P 31 1 2	P 31 2 (0 0 4)	cell choice 1	primitive	6	no	yes	yes
441	152	P 31 2 1	P 31 2"	cell choice 1	primitive	6	no	yes	yes
442	153	P 32 1 2	P 32 2 (0 0 2)	cell choice 1	primitive	6	no	yes	yes
443	154	P 32 2 1	P 32 2"	cell choice 1	primitive	6	no	yes	yes
444	155	R 3 2:H	R 3 2"	hexagonal	rhombohedral	18	no	yes	no
445	155	R 3 2:R	P 3* 2	Rhombohedral	primitive	6	no	yes	no
446	156	P 3 m 1	P 3 -2"	cell choice 1	primitive	6	no	no	no
447	157	P 3 1 m	P 3 -2	cell choice 1	primitive	6	no	no	no
448	158	P 3 c 1	P 3 -2" c	cell choice 1	primitive	6	no	no	no
449	159	P 3 1 c	P 3 -2c	cell choice 1	primitive	6	no	no	no
450	160	R 3 m:H	R 3 -2"	hexagonal	rhombohedral	18	no	no	no
451	160	R 3 m:R	P 3* -2	Rhombohedral	primitive	6	no	no	no
452	161	R 3 c:H	R 3 -2" c	hexagonal	rhombohedral	18	no	no	no
453	161	R 3 c:R	P 3* -2n	Rhombohedral	primitive	6	no	no	no
454	162	P -3 1 2/m	-P 3 2	cell choice 1	primitive	12	yes	no	no
455	163	P -3 1 2/c	-P 3 2c	cell choice 1	primitive	12	yes	no	no
456	164	P -3 2/m 1	-P 3 2"	cell choice 1	primitive	12	yes	no	no
457	165	P -3 2/c 1	-P 3 2" c	cell choice 1	primitive	12	yes	no	no
458	166	R -3 2/m:H	-R 3 2"	hexagonal	rhombohedral	36	yes	no	no
459	166	R -3 2/m:R	-P 3* 2	Rhombohedral	primitive	12	yes	no	no
460	167	R -3 2/c:H	-R 3 2" c	hexagonal	rhombohedral	36	yes	no	no
461	167	R -3 2/c:R	-P 3* 2n	Rhombohedral	primitive	12	yes	no	no

Table 2.5: Trigonal spacegroup information.

hexagonal									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
462	168	P 6	P 6	cell choice 1	primitive	6	no	yes	no
463	169	P 61	P 61	cell choice 1	primitive	6	no	yes	yes
464	170	P 65	P 65	cell choice 1	primitive	6	no	yes	yes
465	171	P 62	P 62	cell choice 1	primitive	6	no	yes	yes
466	172	P 64	P 64	cell choice 1	primitive	6	no	yes	yes

467	173	P 63	P 6c	cell choice 1	primitive	6	no	yes	no
468	174	P -6	P -6	cell choice 1	primitive	6	no	no	no
469	175	P 6/m	-P 6	cell choice 1	primitive	6	yes	no	no
470	176	P 63/m	-P 6c	cell choice 1	primitive	12	yes	no	no
471	177	P 6 2 2	P 6 2	cell choice 1	primitive	12	no	yes	no
472	178	P 61 2 2	P 61 2 (0 0 5)	cell choice 1	primitive	12	no	yes	yes
473	179	P 65 2 2	P 65 2 (0 0 1)	cell choice 1	primitive	12	no	yes	yes
474	180	P 62 2 2	P 62 2 (0 0 4)	cell choice 1	primitive	12	no	yes	yes
475	181	P 64 2 2	P 64 2 (0 0 2)	cell choice 1	primitive	12	no	yes	yes
476	182	P 63 2 2	P 6c 2c	cell choice 1	primitive	12	no	yes	no
477	183	P 6 m m	P 6 -2	cell choice 1	primitive	12	no	no	no
478	184	P 6 c c	P 6 -2c	cell choice 1	primitive	12	no	no	no
479	185	P 63 c m	P 6c -2	cell choice 1	primitive	12	no	no	no
480	186	P 63 m c	P 6c -2c	cell choice 1	primitive	12	no	no	no
481	187	P -6 m 2	P -6 2	cell choice 1	primitive	12	no	no	no
482	188	P -6 c 2	P -6c 2	cell choice 1	primitive	12	no	no	no
483	189	P -6 2 m	P -6 -2	cell choice 1	primitive	12	no	no	no
484	190	P -6 2 c	P -6c -2c	cell choice 1	primitive	12	no	no	no
485	191	P 6/m 2/m 2/m	-P 6 2	cell choice 1	primitive	24	yes	no	no
486	192	P 6/m 2/c 2/c	-P 6 2c	cell choice 1	primitive	24	yes	no	no
487	193	P 63/m 2/c 2/m	-P 6c 2	cell choice 1	primitive	24	yes	no	no
488	194	P 63/m 2/m 2/c	-P 6c 2c	cell choice 1	primitive	24	yes	no	no

Table 2.6: Hexagonal spacegroup information.

cubic									
id	Int. Nr.	long Hermann-Mauguin name	Hall name	cell choice	centered	#	Chiral	Centric	Enantio-morphic
489	195	P 2 3	P 2 2 3	cell choice 1	primitive	12	no	yes	no
490	196	F 2 3	F 2 2 3	cell choice 1	face	48	no	yes	no
491	197	I 2 3	I 2 2 3	cell choice 1	body	24	no	yes	no
492	198	P 21 3	P 2ac 2ab 3	cell choice 1	primitive	12	no	yes	no
493	199	I 21 3	I 2b 2c 3	cell choice 1	body	24	no	yes	no
494	200	P 2/m -3	-P 2 2 3	cell choice 1	primitive	24	yes	no	no
495	201	P 2/n -3:1	P 2 2 3 -1n	cell choice 1	primitive	24	yes	no	no
496	201	P 2/n -3:2	-P 2ab 2bc 3	cell choice 2	primitive	24	yes	no	no
497	202	F 2/m -3	-F 2 2 3	cell choice 1	face	96	yes	no	no
498	203	F 2/d -3:1	F 2 2 3 -1d	cell choice 1	face	96	yes	no	no
499	203	F 2/d -3:2	-F 2uv 2vw 3	cell choice 2	face	96	yes	no	no
500	204	I 2/m -3	-I 2 2 3	cell choice 1	body	48	yes	no	no
501	205	P 21/a -3	-P 2ac 2ab 3	cell choice 1	primitive	24	yes	no	no
502	206	I 21/a -3	-I 2b 2c 3	cell choice 1	body	48	yes	no	no
503	207	P 4 3 2	P 4 2 3	cell choice 1	primitive	24	no	yes	no
504	208	P 42 3 2	P 4n 2 3	cell choice 1	primitive	24	no	yes	no
505	209	F 4 3 2	F 4 2 3	cell choice 1	face	96	no	yes	no
506	210	F 41 3 2	F 4d 2 3	cell choice 1	face	96	no	yes	no
507	211	I 4 3 2	I 4 2 3	cell choice 1	body	48	no	yes	no
508	212	P 43 3 2	P 4acd 2ab 3	cell choice 1	primitive	24	no	yes	yes
509	213	P 41 3 2	P 4bd 2ab 3	cell choice 1	primitive	24	no	yes	yes
510	214	I 41 3 2	I 4bd 2c 3	cell choice 1	body	48	no	yes	no
511	215	P -4 3 m	P -4 2 3	cell choice 1	primitive	24	no	no	no
512	216	F -4 3 m	F -4 2 3	cell choice 1	face	96	no	no	no
513	217	I -4 3 m	I -4 2 3	cell choice 1	body	48	no	no	no
514	218	P -4 3 n	P -4n 2 3	cell choice 1	primitive	24	no	no	no
515	219	F -4 3 c	F -4a 2 3	cell choice 1	face	96	no	no	no
516	220	I -4 3 d	I -4bd 2c 3	cell choice 1	body	48	no	no	no

517	221	P 4/m -3 2/m	-P 4 2 3	cell choice 1	primitive	48	yes	no	no
518	222	P 4/n -3 2/n:1	P 4 2 3 -1n	cell choice 1	primitive	48	yes	no	no
519	222	P 4/n -3 2/n:2	-P 4a 2bc 3	cell choice 2	primitive	48	yes	no	no
520	223	P 42/m -3 2/n	-P 4n 2 3	cell choice 1	primitive	48	yes	no	no
521	224	P 42/n -3 2/m:1	P 4n 2 3 -1n	cell choice 1	primitive	48	yes	no	no
522	224	P 42/n -3 2/m:2	-P 4bc 2bc 3	cell choice 2	primitive	48	yes	no	no
523	225	F 4/m -3 2/m	-F 4 2 3	cell choice 1	face	192	yes	no	no
524	226	F 4/m -3 2/c	-F 4a 2 3	cell choice 1	face	192	yes	no	no
525	227	F 41/d -3 2/m:1	F 4d 2 3 -1d	cell choice 1	face	192	yes	no	no
526	227	F 41/d -3 2/m:2	-F 4vw 2vw 3	cell choice 2	face	192	yes	no	no
527	228	F 41/d -3 2/c	F 4d 2 3 -1ad	cell choice 1	face	192	yes	no	no
528	228	F 41/d -3 2/c	-F 4ud 2vw 3	cell choice 2	face	192	yes	no	no
529	229	I 4/m -3 2/m	-I 4 2 3	cell choice 1	body	96	yes	no	no
530	230	I 41/a -3 2/d	-I 4bd 2c 3	cell choice 1	body	96	yes	no	no

Table 2.7: Cubic spacegroup information.

## Bibliography

- [1] Farha, O.K.; Yazaydin, A.O.; Eryazici, I.; Malliakas, C.D.; Hauser, B.G.; Kanatzidis, M.G.; Nguyen, S.T.; Snurr, R.Q.; Hupp, J.T. *Nature Chem.* **2010**, .
- [2] Qiu, S.L.; Pang, W.Q.; Kessler, H.; J.L. Guth, *Zeolites* **1989**, *9*, 440-444.
- [3] Pluth, J.J.; Smith, J.V.; Bennett, J.M. *Acta Cryst.* **1986**, *C42*, 283-286.
- [4] Calligaris, M.; Nardin, G.; Randaccio, L. *Zeolites* **1983**, *3*, 205-208.
- [5] Pluth, J.J.; Smith, J.V. *American Mineralogist* **1990**, *75*, 501-507.
- [6] Gies, H. *Z. Kristallogr.* **1986**, *175*, 93-104.
- [7] Gard, J.A.; Tait, J.M., in *Refinement of the crystal structure of erionite*, Proceedings of the 3th International Zeolite Conference, edited by Uytterhoeven, J.B. Leuven University Press; Leuven, 1973, pp. 94-99.
- [8] Morris, R.E.; Weigel, S.J.; Henson, N.J.; Bull, L.M.; Janicke, M.T.; Chmelka, B.F.; Cheetham, A.J. *J. Am. Chem. Soc.* **1994**, *116*, 11849-11855.
- [9] Cambor, M.A.; Corma, A.; Lightfoot, P.; Villaescusa, L.A.; Wright, P.A. *Angew. Chem. Int. Ed.* **1997**, *36*, 2659-2661.
- [10] Corma, A.; Rey, F.; Rius, J.; Sabater, M.J.; Valencia, S. *Nature* **2004**, *431*, 287-290.
- [11] Pluth, J.J.; Smith, J.V. *J. Am. Chem. Soc.* **1980**, *102*, 4704-4708.
- [12] Barrer, R.M.; Villiger, H. *Z. Kristal.* **1969**, *128*, 352-370.
- [13] van Koningsveld, H.; van Bekkum, H.; Jansen, J.C. *Acta Cryst.* **1987**, *43*, 127-132.
- [14] van Koningsveld, H.; Jansen, J.C.; van Bekkum, H. *Zeolites* **1990**, *10*, 235-242.
- [15] Olson, D.H.; Kokotailo, G.T.; Lawton, S.L.; Meier, W.M. *J. Phys. Chem.* **1981**, *85*, 2238-2243.
- [16] Pluth, J.J.; Smith, J.V. *J. Phys. Chem.* **1989**, *93*, 6516-6520.
- [17] Marler, B. *Zeolites* **1987**, *7*, 393-397.

# Chapter 3

## Potentials

### 3.1 Functional forms of force fields

The molecular energy can be described as an Taylor expansion in bonds, bends, torsions, etc.

$$\begin{aligned} U = & \sum_{\text{bonds}} U_r(r) + \sum_{\text{bends}} U_\theta(\theta) + \sum_{\text{torsions}} U_\phi(\phi) + \sum_{\text{out-of-plane bends}} U_\chi(\chi) + \sum_{\text{non-bonded}} U_{nb}(r) \\ & + \sum_{\text{bond-bond}} U_{bb'}(r, r') + \sum_{\text{bond-bend}} U_{b\theta'}(r, \theta) + \sum_{\text{bend-bend}} U_{\theta\theta'}(\theta, \theta') \\ & + \sum_{\text{bond-torsion}} U_{r\phi}(r, \phi, r') + \sum_{\text{bend-torsion}} U_{\theta\phi}(\theta, \phi, \theta') + \dots \end{aligned} \quad (3.1)$$

This expansion is believed to capture all the chemical entities we can think of, such as atoms, bonds, angles, etc, and physical properties like equilibrium structures, vibrational spectra, etc. The cross terms are not ad-hoc functions, but arise naturally from this expansion. For example, bonds and bends interact, as the bend angle becomes smaller the bond lengths tend to increase. Their inclusion leads to two advantages: 1) they increase the accuracy of the force field (especially the vibrational frequencies), and 2) they increase the transferability of the diagonal terms  $U_r(r)$ ,  $U_\theta(\theta)$ ,  $U_\phi(\phi)$ ,  $U_\chi(\chi)$ . On top of the terms in Eq. 3.1 one can add ad hoc terms, such as hydrogen bonding, that are not adequately accounted for otherwise.

Eq. 3.1 is historically referred to as an *force field*. The name arose from the lowest order approximation using only springs with *force constants*. Force fields have matured and have become quite accurate and many parameters exists for a wide range of structure. These parameters are crucial and determine the quality of the force field. Unfortunately, deriving high quality parameters remains more than a art rather than a science. However, some progress has been made and in the end of the chapter some algorithms are described how to obtain them.

The terms in Eq. 3.1 consists of a functional form, force constants (a resistance against a change from the optimum value), and a reference value. The functional form is chosen such as to be an accurate description of the true potential energy (either known from experiment or from quantum mechanics), although one can simplify the functional form to decrease computational evaluation time of the energy at the cost of diminished accuracy. This tradeoff has almost vanished for intra-molecular potentials but is still an issue for the non-bonded terms. The reference value is *not* the equilibrium value (except by chance). For example, bond lengths are affected by all other terms in the force field and the more strained a molecule the farther the bond equilibrium length will deviate from its reference value. This means that one can not simply take the equilibrium values from known experiment.

## 3.2 Bonded potentials diagonal terms

### 3.2.1 Bond-stretching potentials

The bond stretching potential describes the change in energy as the bond stretches and contracts. The simplest functional form would be Hook's law:

$$U = \frac{1}{2}k(r - r_0)^2 \quad (3.2)$$

where  $k$  is the force constant and  $r_0$  the reference value for the bond. This form is computationally very fast, but not very realistic. It is well known that it is easier to stretch a bond than it is to compress a bond. The "Morse" potential is an-harmonic and provides a much better description of the energy

$$U = D \left(1 - e^{-\alpha(r - r_0)}\right)^2 \quad (3.3)$$

Expanding around the equilibrium value leads to

$$U = D\alpha^2(r - r_0)^2 \left[1 - \alpha(r - r_0) + \frac{7}{12}\alpha^2(r - r_0)^2 \dots\right] \quad (3.4)$$

The first term is the harmonic potential (with  $k = 2D\alpha^2$ ) and for organic structures where distortions from equilibrium are small the difference between the potentials are small. However, for larger deviations the Morse potential provides a significantly better description. The Morse potential provides a restoring force which goes to zero at long distances. For minimizations starting far equilibrium could result in non-convergence. Some force fields solved this problem by using modification of Hook's law. MM2 added a cubic term making the bond an-harmonic. However, this leads to large negative energies for poor initial geometries with large distortions. MM3 added the quartic term to solve this. Note the  $7/12$  terms in the MM2/3 functional forms originate from the Taylor expansion of the Morse potential, and the cubic and quartic terms are chosen to mimic the Morse potentials for moderate distortions. Dinur and Hagler proposed a functional form based on inverse bond lengths which follows the true potential energy compared to QM over an even wider range

$$U = U_0 + C_2 \left(\frac{1}{r} - \frac{1}{r_0}\right)^2 + C_3 \left(\frac{1}{r} - \frac{1}{r_0}\right)^3 \quad (3.5)$$

The implemented bond-potentials:

- HARMONIC\_BOND

$$U = \frac{1}{2}p_0(r - p_1)^2 \quad (3.6)$$

2 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_1$  in Å.

- CORE\_SHELL\_SPRING

$$U = \frac{1}{2}p_0r^2 \quad (3.7)$$

1 argument:  $p_0/k_B$  in units of K/Å<sup>2</sup>.

- MORSE\_BOND

$$U = p_0 \left[ \left(1 - e^{-p_1(r - p_2)}\right)^2 - 1 \right] \quad (3.8)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  in Å<sup>-1</sup>, and  $p_2$  in Å.

- LJ\_12\_6\_BOND

$$U = \frac{p_0}{r^{12}} - \frac{p_1}{r^6} \quad (3.9)$$

2 arguments:  $p_0/k_B$  in units of K Å<sup>12</sup>, and  $p_1/k_B$  in units of K Å<sup>6</sup>.

- LENNARD\_JONES\_BOND

$$U = 4p_0 \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] \quad (3.10)$$

2 arguments:  $p_0/k_B$  in units of K,  $p_1$  in Å.

- BUCKINGHAM\_BOND

$$U = p_0 e^{-p_1 r} - \frac{p_2}{r^6} \quad (3.11)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup>, and  $p_2/k_B$  in K Å<sup>6</sup>.

- RESTRAINED\_HARMONIC\_BOND

$$U = \begin{cases} \frac{1}{2}p_0(r - p_1)^2 & |r - p_1| \leq p_2 \\ \frac{1}{2}p_0p_2^2 + p_0p_2(|r - p_1| - p_2) & |r - p_1| > p_2 \end{cases} \quad (3.12)$$

3 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_1$  in Å, and  $p_2$  in Å.

- QUARTIC\_BOND

$$U = \frac{1}{2}p_0(r - p_1)^2 + \frac{1}{3}p_2(r - p_1)^3 + \frac{1}{4}p_3(r - p_1)^4 \quad (3.13)$$

4 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_1$  in Å,  $p_2/k_B$  in K/Å<sup>3</sup>, and  $p_3/k_B$  in K/Å<sup>4</sup>.

- CFF\_QUARTIC\_BOND

$$U = p_0(r - p_1)^2 + p_2(r - p_1)^3 + p_3(r - p_1)^4 \quad (3.14)$$

4 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_1$  in Å,  $p_2/k_B$  in K/Å<sup>3</sup>, and  $p_3/k_B$  in K/Å<sup>4</sup>.

- MM3\_BOND

$$U = p_0(r - p_1)^2 \left( 1 - 2.55(r - p_1) + \frac{7}{12}2.55^2(r - p_1)^2 \right) \quad (3.15)$$

2 arguments:  $p_0$  in units of mdyne/Å molecule,  $p_1$  in Å.

- RIGID\_BOND

Use for connections between rigid units.

- FIXED\_BOND

Use for bonds constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Monte-Carlo, Molecular Dynamics, and minimization.

- MEASURE\_BOND

A histogram of the bond-distance can be computed.

### 3.2.2 Urey-Bradley potentials

The Urey-Bradley potential is sometimes used to account for the repulsion between two atoms bound to a common atom. In more modern force field they are replaced by bond/bend cross potentials. Urey-Bradley are essentially just bonds between 1-3 nearest neighbor atoms and the same range of potentials is offered as for 1-2 bonds in RASPA.

- HARMONIC\_UREYBRADLEY

$$U = \frac{1}{2}p_0(r - p_1)^2 \quad (3.16)$$

2 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_1$  in Å.

- MORSE\_UREYBRADLEY

$$U = p_0 \left[ \left( 1 - e^{-p_1(r-p_2)} \right)^2 - 1 \right] \quad (3.17)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  in  $\text{\AA}^{-1}$ , and  $p_2$  in  $\text{\AA}$ .

- LJ\_12\_6\_UREYBRADLEY

$$U = \frac{p_0}{r^{12}} - \frac{p_1}{r^6} \quad (3.18)$$

2 arguments:  $p_0/k_B$  in units of  $\text{K\AA}^{12}$ , and  $p_1/k_B$  in units of  $\text{K\AA}^6$ .

- LENNARD\_JONES\_UREYBRADLEY

$$U = 4p_0 \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] \quad (3.19)$$

2 arguments:  $p_0/k_B$  in units of K,  $p_1$  in  $\text{\AA}$ .

- BUCKINGHAM\_UREYBRADLEY

$$U = p_0 e^{-p_1 r} - \frac{p_2}{r^6} \quad (3.20)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  in  $\text{\AA}^{-1}$ , and  $p_2/k_B$  in  $\text{K\AA}^6$ .

- RESTRAINED\_HARMONIC\_UREYBRADLEY

$$U = \begin{cases} \frac{1}{2} p_0 (r - p_1)^2 & |r - p_1| \leq p_2 \\ \frac{1}{2} p_0 p_2^2 + p_0 p_2 (|r - p_1| - p_2) & |r - p_1| > p_2 \end{cases} \quad (3.21)$$

3 arguments:  $p_0/k_B$  in units of  $\text{K/\AA}^2$ ,  $p_1$  in  $\text{\AA}$ , and  $p_2$  in  $\text{\AA}$ .

- QUARTIC\_UREYBRADLEY

$$U = \frac{1}{2} p_0 (r - p_1)^2 + \frac{1}{3} p_2 (r - p_1)^3 + \frac{1}{4} p_3 (r - p_1)^4 \quad (3.22)$$

4 arguments:  $p_0/k_B$  in units of  $\text{K/\AA}^2$ ,  $p_1$  in  $\text{\AA}$ ,  $p_2/k_B$  in  $\text{K/\AA}^3$ , and  $p_3/k_B$  in  $\text{K/\AA}^4$ .

- CFF\_QUARTIC\_UREYBRADLEY

$$U = p_0 (r - p_1)^2 + p_2 (r - p_1)^3 + p_3 (r - p_1)^4 \quad (3.23)$$

4 arguments:  $p_0/k_B$  in units of  $\text{K/\AA}^2$ ,  $p_1$  in  $\text{\AA}$ ,  $p_2/k_B$  in  $\text{K/\AA}^3$ , and  $p_3/k_B$  in  $\text{K/\AA}^4$ .

- MM3\_UREYBRADLEY

$$U = p_0 (r - p_1)^2 \left( 1 - 2.55 (r - p_1) + \frac{7}{12} 2.55^2 (r - p_1)^2 \right) \quad (3.24)$$

2 arguments:  $p_0$  in units of mdyne/ $\text{\AA}$  molecule,  $p_1$  in  $\text{\AA}$ .

- RIGID\_UREYBRADLEY

Use for connections between rigid units.

- FIXED\_UREYBRADLEY

Use for bonds constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Monte-Carlo, Molecular Dynamics, and minimization.

- MEASURE\_UREYBRADLEY

A histogram of the Urey-Bradley distance can be computed.

### 3.2.3 Bending potential

The simplest approach for an angle potential is the harmonic potential

$$U = \frac{1}{2}k(\theta - \theta_0)^2 \quad (3.25)$$

Angles are much softer than bonds, especially in zeolites where a Si-O-Si angle ranges between 135 and 180 degrees. A problem with all polynomial representations of angles is that angles of 180 degrees results in singular point (unless the reference angle is 180 degrees). The case of 0 degree is not possible due to repulsion of the i and k atoms in the i-j-k bend. The singularity is due to the fact that the force expression of such a polynomial contains a factor  $1/\sin(\theta)$ . A common solution is to use a trigonometric function

$$U = \frac{1}{2}k[\cos(\theta) - \cos(\theta_0)]^2 \quad (3.26)$$

Note that close to the maximum these potentials have no restoring force, but for small distortions this is not a problem. The MM force fields use higher order terms. A six power term was needed to describe the highly bent bicyclo[1.1.1]pentane. Cubic terms and higher become desirable when the bending is more than 10-15 degrees. MM3 angle bending has been divided into in-plane and out-of-plane bending for planar trigonal centers.

- HARMONIC\_BEND,CORE\_SHELL\_BEND

$$U = \frac{1}{2}p_0(\theta_{ijk} - p_1)^2 \quad (3.27)$$

2 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup> and  $p_1$  in degrees.

- QUARTIC\_BEND

$$U = \frac{1}{2}p_0(\theta_{ijk} - p_1)^2 + \frac{1}{3}p_2(\theta_{ijk} - p_1)^3 + \frac{1}{4}p_3(\theta_{ijk} - p_1)^4 \quad (3.28)$$

4 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup>,  $p_1$  in degrees,  $p_2/k_B$  in K/rad<sup>3</sup>, and  $p_3/k_B$  in K/rad<sup>4</sup>.

- CFF\_QUARTIC\_BEND

$$U = p_0(\theta_{ijk} - p_1)^2 + p_2(\theta_{ijk} - p_1)^3 + p_3(\theta_{ijk} - p_1)^4 \quad (3.29)$$

4 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup>,  $p_1$  in degrees,  $p_2/k_B$  in K/rad<sup>3</sup>, and  $p_3/k_B$  in K/rad<sup>4</sup>.

- HARMONIC\_COSINE\_BEND

$$U = \frac{1}{2}p_0(\cos\theta_{ijk} - \cos p_1)^2 \quad (3.30)$$

2 arguments:  $p_0/k_B$  in units of K and  $p_1$  in degrees.

- COSINE\_BEND

$$U = p_0(1 + \cos(p_1\theta_{ijk} - p_2)) \quad (3.31)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  dimensionless, and  $p_2$  in degrees.

- MM3\_BEND

$$\begin{aligned} U = \frac{1}{2}p_0(\theta_{ijk} - p_1)^2 & \left( 1 - 0.014(\theta_{ijk} - p_1) + 5.6 \times 10^{-5}(\theta_{ijk} - p_1)^2 - 7 \times 10^{-7}(\theta_{ijk} - p_1)^3 \right. \\ & \left. + 2.2 \times 10^{-8}(\theta_{ijk} - p_1)^4 \right) \end{aligned} \quad (3.32)$$

2 arguments:  $p_0$  in units of mdyne Å/rad<sup>2</sup>,  $p_1$  in degrees.




Figure 3.1: The definition of the Wilson inversion-bend angle  $\chi$ . On the left a positive Wilson angle, and on the right a negative Wilson angle.

- MM3\_IN\_PLANE\_BEND

$$U = \frac{1}{2}p_0 (\theta_{ijk} - p_1)^2 \left( 1 - 0.014(\theta_{ijk} - p_1) + 5.6 \times 10^{-5}(\theta_{ijk} - p_1)^2 - 7 \times 10^{-7}(\theta_{ijk} - p_1)^3 + 2.2 \times 10^{-8}(\theta_{ijk} - p_1)^4 \right) \quad (3.33)$$

2 arguments:  $p_0$  in units of mdyne Å/rad<sup>2</sup>,  $p_1$  in degrees. The bend is 'in-plane' and only applicable to bends in a defined planar trigonal centers. The bend is dependend on the fourth atom of the trigonal center.

- FIXED\_BEND

Use for bend-angle constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Molecular Dynamics and minimization. Does not work (yet) in Monte-Carlo.

- MEASURE\_BEND

A histogram of the bend angle can be computed.

### 3.2.4 Wilson inversion-bend potential

Common planar molecule that contain a double bond or sp<sup>2</sup> hybridization form planar groups with trigonal centers. For example: the carbon and nitrogen centers in formamide, and the carbon centers in benzene. The mode of motion is different from bond stretching, bending, and internal rotation. The associated harmonic potential is

$$U = \frac{1}{2}k(\chi)^2 \quad (3.34)$$

with  $\chi$  the out-of-plane angle. Two possible definitions are in use

1. the distance of the central atom from the plane defined by the other three atoms (pyramid height),
2. the average angle between any bond that extends from the central atom and the plane defined by the other two bonds.

Note that an alternative to the out-of-plane angle is the *improper torsion* using

$$U = \frac{1}{2}k(1 - \cos 2\chi) \quad (3.35)$$

The out-of-plane potential can also be used for non-planar structure, for example in united-atom for chiral centers to avoid inversion of the chiral center. Another example of its use is coordination complexes where




Figure 3.2: The definition of the dihedral angle  $\phi$ : the angle between the planes formed by atoms a-b-c and b-c-d. On the left a positive dihedral angle, and on the right a negative dihedral angle.

now the plane of the ligands need no longer be defined exactly. In square planar complexes it is necessary to define an average plane through the ligands (usually the least-square plane). Note that the definition include one central atom which is listed as the second in  $a - b - c - d$ :  $a$ ,  $c$ , and  $d$  are bonded to the central atom  $b$ . The inversion angle potential is the average of the three possible inversion angle terms.

- HARMONIC\_INVERSION

$$U = \frac{1}{2} p_0 (\chi_{ijk} - p_1)^2 \quad (3.36)$$

2 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup> and  $p_1$  in degrees.

- HARMONIC\_COSINE\_INVERSION

$$U = \frac{1}{2} p_0 (\cos(\chi_{ijk}) - \cos(p_1))^2 \quad (3.37)$$

2 arguments:  $p_0/k_B$  in units of K and  $p_1$  in degrees.

- PLANAR\_INVERSION

$$U = p_0 (1 - \cos(\chi)) \quad (3.38)$$

1 argument:  $p_0/k_B$  in units of K.

- MM3\_INVERSION

$$\begin{aligned} U = \frac{1}{2} p_0 (\chi - p_1)^2 & \left( 1 - 0.014(\chi - p_1) + 5.6 \times 10^{-5} (\chi - p_1)^2 - 7 \times 10^{-7} (\chi - p_1)^3 \right. \\ & \left. + 2.2 \times 10^{-8} (\chi - p_1)^4 \right) \end{aligned} \quad (3.39)$$

2 arguments:  $p_0$  in units of mdyne Å/rad<sup>2</sup>,  $p_1$  in degrees.

- FIXED\_INVERSION\_BEND

Use for inversion bend-angle constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Molecular Dynamics and minimization. Does not work (yet) in Monte-Carlo.

### 3.2.5 Torsion potential

Intramolecular rotations about bonds do not occur freely. A possible description with a physical interpretation is the three-term Fourier expansion

$$U = \frac{V_1}{2} [1 + \cos \phi] + \frac{V_2}{2} [1 - \cos 2\phi] + \frac{V_3}{2} [1 + \cos 3\phi] \quad (3.40)$$

1. the 1 fold-term has been attributed to residual dipole-dipole interactions, to Van der Waal interactions, or to any other direct interaction between atoms not accounted for otherwise,
2. the 2-fold arises from conjugation or hyper conjugation, being geometrically related to p orbitals,
3. and the 3-fold term has a steric (or bonding/anti-bonding) origin.

The values for 4-fold or higher are small and it is not known whether these are essential to include. It may be that Van der Waals and dipole interactions already take care of these effects. Torsions are even softer than bond angles. All possible values can be found in structures. Therefore, the energy function must be valid over the entire range, the function must be periodic, and for reasons of symmetry have stationary points at 0 and 180 degrees. The periodicity is the number of minima for the potential, usually 3 for an  $sp^3$ - $sp^3$  bond and 2 for a conjugate bond.

The definition of a torsion includes two central and two terminal atoms. The term "torsional" means an internal rigid rotation and "dihedral" means a rotation of two vicinal bonds about a middle bond.

- HARMONIC\_DIHEDRAL

$$U = \frac{1}{2} p_0 (\phi_{ijkl} - p_1)^2 \quad (3.41)$$

2 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup>,  $p_1$  in degrees.

- HARMONIC\_COSINE\_DIHEDRAL

$$U = \frac{1}{2} p_0 [\cos(\phi_{ijkl}) - \cos(p_1)]^2 \quad (3.42)$$

2 arguments:  $p_0/k_B$  in units of K,  $p_1$  in degrees.

- THREE\_COSINE\_DIHEDRAL

$$U = \frac{1}{2} p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2} p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2} p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.43)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K

- MM3\_DIHEDRAL

$$U = \frac{1}{2} p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2} p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2} p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.44)$$

3 arguments:  $p_0, p_1, p_2$  in units of kcal/mol.

- CFF\_DIHEDRAL

$$U = p_0 [1 - \cos(\phi_{ijkl})] + p_1 [1 - \cos(2\phi_{ijkl})] + p_2 [1 - \cos(3\phi_{ijkl})] \quad (3.45)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- CFF\_DIHEDRAL2

$$U = p_0 [1 + \cos(\phi_{ijkl})] + p_1 [1 + \cos(2\phi_{ijkl})] + p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.46)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- SIX\_COSINE DIHEDRAL

The Ryckaert-Bellemans potentials is often used for alkanes, the use implies exclusion of VDW-interactions between the first and last atoms of the dihedral, and  $\phi' = \phi - \pi$  is defined according to the polymer convention  $\phi'(\text{trans}) = 0$ .

$$U = \sum_{n=0}^5 p_n \cos^n (\phi'_{ijkl}) \quad (3.47)$$

$$= p_0 + p_1 \cos (\phi'_{ijkl}) + p_2 \cos^2 (\phi'_{ijkl}) + p_3 \cos^3 (\phi'_{ijkl}) + p_4 \cos^4 (\phi'_{ijkl}) + p_5 \cos^5 (\phi'_{ijkl}) \quad (3.48)$$

6 arguments:  $p_0/k_B, \dots, p_5/k_B$  in units of K. Rewritten in terms of  $\phi$  the potential reads

$$U = p_0 - p_1 \cos (\phi_{ijkl}) + p_2 \cos^2 (\phi_{ijkl}) - p_3 \cos^3 (\phi_{ijkl}) + p_4 \cos^4 (\phi_{ijkl}) - p_5 \cos^5 (\phi_{ijkl}) \quad (3.49)$$

- TRAPPE DIHEDRAL

$$U = p_0 + p_1 [1 + \cos (\phi_{ijkl})] + p_2 [1 - \cos (2\phi_{ijkl})] + p_3 [1 + \cos (3\phi_{ijkl})] \quad (3.50)$$

4 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B$  in units of K.

- CVFF DIHEDRAL

$$U = p_0 [1 + \cos (p_1 \phi_{ijkl} - p_2)] \quad (3.51)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  dimensionless, and  $p_2$  in degrees.

- OPLS DIHEDRAL

$$U = \frac{1}{2}p_0 + \frac{1}{2}p_1 [1 + \cos (\phi_{ijkl})] + \frac{1}{2}p_2 [1 - \cos (2\phi_{ijkl})] + \frac{1}{2}p_3 [1 + \cos (3\phi_{ijkl})] \quad (3.52)$$

4 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B$  in units of K.

- FOURIER\_SERIES DIHEDRAL

The general form of a Fourier expansion is:

$$U = \sum_{n=1}^6 [a_n \cos (n\phi) + b_n \sin (n\phi)] \quad (3.53)$$

This form uses equilibrium angles of 0 for  $n = 1, 3, 5$  and 180 for  $n = 2, 4, 6$

$$\begin{aligned} U = & \frac{1}{2}p_0 [1 + \cos \phi] + \frac{1}{2}p_1 [1 - \cos (2\phi)] + \frac{1}{2}p_2 [1 + \cos (3\phi)] + \\ & \frac{1}{2}p_3 [1 - \cos (4\phi)] + \frac{1}{2}p_4 [1 + \cos (5\phi)] + \frac{1}{2}p_5 [1 - \cos (6\phi)] \end{aligned} \quad (3.54)$$

6 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B, p_4/k_B, p_5/k_B$  in units of K.

- FOURIER\_SERIES DIHEDRAL\_2

The general form of a Fourier expansion is:

$$U = \sum_{n=1}^6 [a_n \cos (n\phi) + b_n \sin (n\phi)] \quad (3.55)$$

This form uses equilibrium angles of 0 for  $n = 1, 3, 4, 5, 6$  and 180 for  $n = 2$

$$\begin{aligned} U = & \frac{1}{2}p_0 [1 + \cos \phi] + \frac{1}{2}p_1 [1 - \cos (2\phi)] + \frac{1}{2}p_2 [1 + \cos (3\phi)] + \\ & \frac{1}{2}p_3 [1 + \cos (4\phi)] + \frac{1}{2}p_4 [1 + \cos (5\phi)] + \frac{1}{2}p_5 [1 + \cos (6\phi)] \end{aligned} \quad (3.56)$$

6 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B, p_4/k_B, p_5/k_B$  in units of K.

- **FIXED\_DIHEDRAL**

Use for dihedral-angle constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Molecular Dynamics and minimization. Does not work (yet) in Monte-Carlo.

The following identities are convenient when dealing with torsions:

$$\begin{aligned}\cos 1x &= \cos x \\ \cos 2x &= -1 + 2 \cos^2 x \\ \cos 3x &= -3 \cos x + 4 \cos^3 x \\ \cos 4x &= 1 - 8 \cos^2 x + 8 \cos^4 x \\ \cos 5x &= 5 \cos x - 20 \cos^3 x + 16 \cos^5 x \\ \cos 6x &= -1 + 18 \cos^2 x - 48 \cos^4 x + 32 \cos^6 x\end{aligned}\tag{3.57}$$

$$\begin{aligned}\sin 1x &= \sin x \\ \sin 2x &= (\sin x)(2 \cos x) \\ \sin 3x &= (\sin x)(-1 + 4 \cos^2 x) \\ \sin 4x &= (\sin x)(-4 \cos x + 8 \cos^3 x) \\ \sin 5x &= (\sin x)(1 - 12 \cos^2 x + 16 \cos^4 x) \\ \sin 6x &= (\sin x)(6 \cos x - 32 \cos^3 x + 32 \cos^5 x)\end{aligned}\tag{3.58}$$

### 3.2.6 Improper torsion potential

The improper torsion is an alternative for the out-of-plane angle, and a possible definition is

$$U = \frac{1}{2}k(1 - \cos 2\chi)\tag{3.59}$$

It is termed "improper torsion" because it simply treats the four atoms in the plane as if they were bonded in the same way as in a true torsional angle. Note that the definition include one central atom which is listed as the second in  $a - b - c - d$ :  $a$ ,  $c$ , and  $d$  are bonded to the central atom  $b$ . Improper torsions are often used to keep sp<sup>2</sup> atoms planar and sp<sup>3</sup> atoms in a tetrahedral geometry.

The CHARMM convention is to list the central atom first, while there are no rules how to order the other three atoms. Hence, six possibilities exist for the definition of an improper torsion. The AMBER convention is that the out-of-plane atom is listed in the third position and the order of the other atoms is determined alphabetically by atom type, and by the atom number (i.e. the order in the molecule) when atom types are identical.

- **HARMONIC\_IMPROPER\_DIHEDRAL**

$$U = \frac{1}{2}p_0(\phi_{ijkl} - p_2)^2\tag{3.60}$$

2 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup>,  $p_1$  in degrees.

- **HARMONIC\_COSINE\_IMPROPER\_DIHEDRAL**

$$U = \frac{1}{2}p_0[\cos(\phi_{ijkl}) - \cos(p_1)]^2\tag{3.61}$$

2 arguments:  $p_0/k_B$  in units of K,  $p_1$  in degrees.




Figure 3.3: The most common (CVFF, DLPOLY) definition of the improper dihedral angle  $\phi$ : the angle between the planes formed by atoms 'a-c-d' and 'c-d-b'. On the left a positive improper dihedral angle, and on the right a negative improper dihedral angle. The atoms need to be listed in the order 'a-c-d-b'. Note that an exchange of atoms 'c' and 'd' leads to a change of sign, but *not* in magnitude.




Figure 3.4: A second definition of the improper dihedral angle (CHARMM, AMBER). The central atom is 'c', and the improper torsion is enter as 'a-b-c-d'. However, an exchange of terminal atoms leads to a change in magntitude and the improper torsion needs to be symmetrized by adding two additional improper torsions 'b-d-c-a' and 'd-a-c-b' and rescaling the force constant by a factor of 1/3.

- THREE\_COSINE\_IMPROPER\_DIHEDRAL

$$U = \frac{1}{2}p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2}p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2}p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.62)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- MM3\_IMPROPER\_DIHEDRAL

$$U = \frac{1}{2}p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2}p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2}p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.63)$$

3 arguments:  $p_0, p_1, p_2$  in units of kcal/mol.

- CFF\_IMPROPER\_DIHEDRAL

$$U = p_0 [1 - \cos(\phi_{ijkl})] + p_1 [1 - \cos(2\phi_{ijkl})] + p_2 [1 - \cos(3\phi_{ijkl})] \quad (3.64)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- CFF\_IMPROPER\_DIHEDRAL2

$$U = p_0 [1 + \cos(\phi_{ijkl})] + p_1 [1 + \cos(2\phi_{ijkl})] + p_2 [1 + \cos(3\phi_{ijkl})] \quad (3.65)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- **SIX\_COSINE\_IMPROPER\_DIHEDRAL**

The Ryckaert-Bellemans potentials is often used for alkanes, the use implies exclusion of VDW-interactions between the first and last atoms of the dihedral, and  $\phi' = \phi - \pi$  is defined according to the polymer convention  $\phi'(\text{trans}) = 0$ .

$$U = \sum_{n=0}^5 p_n \cos^n (\phi'_{ijkl}) \quad (3.66)$$

$$= p_0 + p_1 \cos (\phi'_{ijkl}) + p_2 \cos^2 (\phi'_{ijkl}) + p_3 \cos^3 (\phi'_{ijkl}) + p_4 \cos^4 (\phi'_{ijkl}) + p_5 \cos^5 (\phi'_{ijkl}) \quad (3.67)$$

6 arguments:  $p_0/k_B, \dots, p_5/k_B$  in units of K. Rewritten in terms of  $\phi$  the potential reads

$$U = p_0 - p_1 \cos (\phi_{ijkl}) + p_2 \cos^2 (\phi_{ijkl}) - p_3 \cos^3 (\phi_{ijkl}) + p_4 \cos^4 (\phi_{ijkl}) - p_5 \cos^5 (\phi_{ijkl}) \quad (3.68)$$

- **TRAPPE\_IMPROPER\_DIHEDRAL**

$$U = p_0 + p_1 [1 + \cos (\phi_{ijkl})] + p_2 [1 - \cos (2\phi_{ijkl})] + p_3 [1 + \cos (3\phi_{ijkl})] \quad (3.69)$$

4 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B$  in units of K.

- **CVFF\_IMPROPER\_DIHEDRAL**

$$U = p_0 [1 + \cos (p_1 \phi_{ijkl} - p_2)] \quad (3.70)$$

3 arguments:  $p_0/k_B$  in units of K,  $p_1$  dimensionless, and  $p_2$  in degrees.

- **OPLS\_IMPROPER\_DIHEDRAL**

$$U = \frac{1}{2}p_0 + \frac{1}{2}p_1 [1 + \cos (\phi_{ijkl})] + \frac{1}{2}p_2 [1 - \cos (2\phi_{ijkl})] + \frac{1}{2}p_3 [1 + \cos (3\phi_{ijkl})] \quad (3.71)$$

4 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B$  in units of K.

- **FOURIER\_SERIES\_IMPROPER\_DIHEDRAL**

The general form of a Fourier expansion is:

$$U = \sum_{n=1}^6 [a_n \cos (n\phi) + b_n \sin (n\phi)] \quad (3.72)$$

This form uses equilibrium angles of 0 for  $n = 1, 3, 5$  and 180 for  $n = 2, 4, 6$

$$\begin{aligned} U = & \frac{1}{2}p_0 [1 + \cos \phi] + \frac{1}{2}p_1 [1 - \cos (2\phi)] + \frac{1}{2}p_2 [1 + \cos (3\phi)] + \\ & \frac{1}{2}p_3 [1 - \cos (4\phi)] + \frac{1}{2}p_4 [1 + \cos (5\phi)] + \frac{1}{2}p_5 [1 - \cos (6\phi)] \end{aligned} \quad (3.73)$$

6 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B, p_4/k_B, p_5/k_B$  in units of K.

- **FOURIER\_SERIES\_IMPROPER\_DIHEDRAL\_2**

The general form of a Fourier expansion is:

$$U = \sum_{n=1}^6 [a_n \cos (n\phi) + b_n \sin (n\phi)] \quad (3.74)$$

This form uses equilibrium angles of 0 for  $n = 1, 3, 4, 5, 6$  and 180 for  $n = 2$

$$U = \frac{1}{2}p_0 [1 + \cos \phi] + \frac{1}{2}p_1 [1 - \cos (2\phi)] + \frac{1}{2}p_2 [1 + \cos (3\phi)] + \frac{1}{2}p_3 [1 + \cos (4\phi)] + \frac{1}{2}p_4 [1 + \cos (5\phi)] + \frac{1}{2}p_5 [1 + \cos (6\phi)] \quad (3.75)$$

6 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B, p_3/k_B, p_4/k_B, p_5/k_B$  in units of K.

- **FIXED\_IMPROPER\_DIHEDRAL**

Use for improper-dihedral-angle constraint using the 'SHAKE' and 'RATTLE'-algorithm. Applies to Molecular Dynamics and minimization. Does not work (yet) in Monte-Carlo.

### 3.3 Non-bonded potentials

#### 3.3.1 Van der Waals potentials

The general expression for Van der Waals potentials when using a cutoff distance is

$$U_{ij}^{\text{VDW}} = \begin{cases} U_{ij}(r_{ij}) & \text{if } r_{ij} \leq r_c \\ 0 & \text{otherwise} \end{cases} \quad (3.76)$$

- **NONE**

$$U = 0 \quad (3.77)$$

zero parameters.

**LENNARD\_JONES**

- **LENNARD\_JONES\_SMOOTHED3**  
**LENNARD\_JONES\_SMOOTHED5**

$$U = 4p_0 \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] \quad (3.78)$$

2 parameters:  $p_0/k_B$  in units of K, and  $p_1$  in Å.

**FEYNMAN\_HIBBS.LENNARD\_JONES**

- **FEYNMAN\_HIBBS.LENNARD\_JONES\_SMOOTHED3**  
**FEYNMAN\_HIBBS.LENNARD\_JONES\_SMOOTHED5**

$$U = 4p_0 \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] + \frac{\hbar^2}{24p_2 k_B T} 4p_0 \left[ 132 \left( \frac{p_1}{r} \right)^{12} - 30 \left( \frac{p_1}{r} \right)^6 \right] \frac{1}{r^2} \quad (3.79)$$

3 parameters:  $p_0/k_B$  in units of K,  $p_1$  in Å, and  $p_2$  is the reduced mass in unified atomic mass units.

**FEYNMAN\_HIBBS2.LENNARD\_JONES2**

- **FEYNMAN\_HIBBS.LENNARD\_JONES2\_SMOOTHED3**  
**FEYNMAN\_HIBBS.LENNARD\_JONES2\_SMOOTHED5**

$$U = 4p_0 \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] + 4p_0 \frac{1}{T} \left[ 132 \left( \frac{p_1}{r} \right)^{12} - 30 \left( \frac{p_1}{r} \right)^6 \right] \frac{p_2}{r^2} \quad (3.80)$$

3 parameters:  $p_0/k_B$  in units of K,  $p_1$  in Å, and  $p_2$  in units of Å<sup>2</sup>.

- **LENNARD\_JONES\_SHIFTED\_FORCE**

$$U = 4p_0 \left\{ \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] - \left[ \left( \frac{p_1}{r_c} \right)^{12} - \left( \frac{p_1}{r_c} \right)^6 \right] + \left[ 12 \left( \frac{p_1}{r_c} \right)^{12} - 6 \left( \frac{p_1}{r_c} \right)^6 \right] \frac{(r - r_c)}{r_c} \right\} \quad (3.81)$$

2 parameters:  $p_0/k_B$  in units of K, and  $p_1$  in Å.

- LENNARD\_JONES\_SHIFTED\_FORCE2

$$4p_0 \left\{ \left[ \left( \frac{p_1}{r} \right)^{12} - \left( \frac{p_1}{r} \right)^6 \right] + \left[ 6 \left( \frac{p_1}{r_c} \right)^{12} - 3 \left( \frac{p_1}{r_c} \right)^6 \right] \frac{r^2}{r_c^2} + 7 \left( \frac{p_1}{r_c} \right)^{12} + 4 \left( \frac{p_1}{r_c} \right)^6 \right\} \quad (3.82)$$

2 parameters:  $p_0/k_B$  in units of K, and  $p_1$  in Å.

- POTENTIAL\_12\_6
- POTENTIAL\_12\_6\_SMOOTHED3
- POTENTIAL\_12\_6\_SMOOTHED5

$$U = \frac{p_0}{r^{12}} - \frac{p_1}{r^6} \quad (3.83)$$

2 parameters:  $p_0/k_B$  in units of K Å<sup>12</sup>, and  $p_1/k_B$  in units of K Å<sup>6</sup>.

- POTENTIAL\_12\_6\_2\_0
- POTENTIAL\_12\_6\_2\_0\_SMOOTHED3
- POTENTIAL\_12\_6\_2\_0\_SMOOTHED5

$$U = \frac{p_0}{r^{12}} + \frac{p_1}{r^6} + \frac{p_2}{r^2} + p_3 \quad (3.84)$$

4 parameters:  $p_0/k_B$  in units of K Å<sup>12</sup>,  $p_1/k_B$  in units of K Å<sup>6</sup>,  $p_2/k_B$  in units of K Å<sup>2</sup>, and  $p_3$  in units of K.

- MORSE
- MORSE\_SMOOTHED3
- MORSE\_SMOOTHED5

$$U = p_0 \left[ (1 - e^{-p_1*(r-p_2)})^2 - 1 \right] \quad (3.85)$$

3 parameters:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup> and  $p_2$  in units of Å.

- MORSE2
- MORSE2\_SMOOTHED3
- MORSE2\_SMOOTHED5

$$U = p_0 \left[ e^{p_1*(1-r/p_2)} - 2e^{(p_1/2)*(1-r/p_2)} \right] \quad (3.86)$$

3 parameters:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup> and  $p_2$  in units of Å.

- MORSE3
- MORSE3\_SMOOTHED3
- MORSE3\_SMOOTHED5

$$U = p_0 \left[ \left( 1 - e^{\left( \frac{-\ln 2}{2^{1/6}-1} \right) \left( \frac{r}{p_2} - 2^{1/6} \right)} \right)^2 - 1 \right] \quad (3.87)$$

2 parameters:  $p_0/k_B$  in units of K  $p_2$  in units of Å. This form of the Morse potential resembles the Lennard-Jones potential.

- CFF\_9\_6
- CFF\_9\_6\_SMOOTHED3
- CFF\_9\_6\_SMOOTHED5

$$U = \frac{p_0}{r^9} - \frac{p_1}{r^6} \quad (3.88)$$

2 parameters:  $p_0/k_B$  in units of K Å<sup>9</sup>, and  $p_1/k_B$  in units of K Å<sup>6</sup>.

CFF\_EPS\_SIGMA

- CFF\_EPS\_SIGMA\_SMOOTHED3  
CFF\_EPS\_SIGMA\_SMOOTHED5

$$U_{ij} = p_0 \left[ 2 \left( \frac{p_1}{r} \right)^9 - 3 \left( \frac{p_1}{r} \right)^6 \right] \quad (3.89)$$

2 parameters:  $p_0/k_B$  in units of K, and  $p_1$  in Å.

BUCKINGHAM

- BUCKINGHAM\_SMOOTHED3  
BUCKINGHAM\_SMOOTHED5

$$U = p_0 e^{-p_1 r} - \frac{p_2}{r^6} \quad (3.90)$$

3 parameters:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup>, and  $p_2$  in K Å<sup>6</sup>. Warning: in literature sometimes  $\rho = \frac{1}{p_1}$  is given,  $\rho$  is usually around 0.3-0.4 Å,  $p_1$  is usually around 2-4 Å<sup>-1</sup>.

BUCKINGHAM2

- BUCKINGHAM2\_SMOOTHED3  
BUCKINGHAM2\_SMOOTHED5

$$U = \begin{cases} 10^{10} & r < p_3 \\ p_0 e^{-p_1 r} - \frac{p_2}{r^6} & \text{otherwise} \end{cases} \quad (3.91)$$

4 parameters:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup>,  $p_2$  in K Å<sup>6</sup>, and  $p_3$  in [Å]. Warning: in literature sometimes  $\rho = \frac{1}{p_1}$  is given,  $\rho$  is usually around 0.3-0.4 Å,  $p_1$  is usually around 2-4 Å<sup>-1</sup>.

MM3\_VDW

- MM3\_VDW\_SMOOTHED3  
MM3\_VDW\_SMOOTHED5

$$U_{ij} = \begin{cases} \sqrt{p_0^i p_0^j} \left[ 1.84 \times 10^5 e^{-\frac{12}{P}} - 2.25 P^6 \right] & \text{if } P \geq 3.02 \\ \sqrt{p_0^i p_0^j} 192.27 P^2 & \text{if } P < 3.02 \end{cases} \quad (3.92)$$

with  $P = \frac{p_1^i + p_1^j}{r_{ij}}$  and where  $p_1^i$  and  $p_1^j$  are the VDW radii of atoms  $i$  and  $j$ , and  $r_{ij}$  the separation distance in Å between atoms  $i$  and  $j$ .

2 arguments:  $p_0$  in units of kcal/mol,  $p_1$  in units of Å.

MATSUOKA\_CLEMENTI\_YOSHIMINE

- MATSUOKA\_CLEMENTI\_YOSHIMINE\_SMOOTHED3  
MATSUOKA\_CLEMENTI\_YOSHIMINE\_SMOOTHED5

$$U = p_0 e^{-p_1 r_{ij}} + p_2 e^{-p_3 r_{ij}} \quad (3.93)$$

4 arguments:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup>,  $p_2/k_B$  in units of K, and  $p_3$  in units of Å<sup>-1</sup>.

GENERIC

- GENERIC\_SMOOTHED3  
GENERIC\_SMOOTHED5

$$U = p_0 e^{-p_1 r} - \frac{p_2}{r^4} - \frac{p_3}{r^6} - \frac{p_4}{r^8} - \frac{p_5}{r^{10}} \quad (3.94)$$

6 arguments:  $p_0/k_B$  in units of K,  $p_1$  in units of Å<sup>-1</sup>,  $p_2/k_B$  in units of K Å<sup>4</sup>,  $p_3/k_B$  in units of K Å<sup>6</sup>,  $p_4/k_B$  in units of K Å<sup>8</sup>, and  $p_5/k_B$  in units of K Å<sup>10</sup>.

- PELLENQ\_NICHOLSON
- PELLENQ\_NICHOLSON\_SMOOTHED3  
PELLENQ\_NICHOLSON\_SMOOTHED5

$$U = p_0 e^{-p_1 r} - f_6 \frac{p_2}{r^6} - f_8 \frac{p_3}{r^8} - f_{10} \frac{p_4}{r^{10}} \quad (3.95)$$

with

$$f_{2n} = 1 - \sum_{k=0}^{2n} \frac{(p_1 r_{ij})^k}{k!} e^{-p_1 r_{ij}} \quad (3.96)$$

5 arguments:  $p_0/k_B$  in units of K,  $p_1$  in units of  $\text{\AA}^{-1}$ ,  $p_2/k_B$  in units of  $\text{K\AA}^6$ ,  $p_3/k_B$  in units of  $\text{K\AA}^8$ , and  $p_4/k_B$  in units of  $\text{K\AA}^{10}$ .

#### HYDRATED\_ION\_WATER

- HYDRATED\_ION\_WATER\_SMOOTHED3  
HYDRATED\_ION\_WATER\_SMOOTHED5

$$U = p_0 e^{-p_1 r} - \frac{p_2}{r^4} - \frac{p_3}{r^6} - \frac{p_4}{r^{12}} \quad (3.97)$$

5 arguments:  $p_0/k_B$  in units of K,  $p_1$  in units of  $\text{\AA}^{-1}$ ,  $p_2/k_B$  in units of  $\text{K\AA}^4$ ,  $p_3/k_B$  in units of  $\text{K\AA}^6$ , and  $p_4/k_B$  in units of  $\text{K\AA}^{12}$ .

#### MIE

- MIE\_SMOOTHED3  
MIE\_SMOOTHED5
- The Mie-potential [1]

$$U = \left( \frac{p_0}{r^{p_1}} - \frac{p_2}{r^{p_3}} \right) \quad (3.98)$$

4 arguments:  $p_0/k_B$  in units of  $\text{K\AA}^{p_1}$ ,  $p_1$  dimensionless,  $p_2/k_B$  in units of  $\text{K\AA}^{p_3}$ , and  $p_3$  dimensionless.

#### BORN\_HUGGINS\_MEYER

- BORN\_HUGGINS\_MEYER\_SMOOTHED3  
BORN\_HUGGINS\_MEYER\_SMOOTHED5

$$U_{ij} = p_0 e^{p_1(p_2 - r_{ij})} - \frac{p_3}{r_{ij}^6} - \frac{p_4}{r_{ij}^8} \quad (3.99)$$

5 arguments:  $p_0/k_B$  in units of K,  $p_1$  dimensionless,  $p_2$  in units of  $\text{\AA}$ ,  $p_3/k_B$  in units of  $\text{K\AA}^6$ , and  $p_4/k_B$  in units of  $\text{K\AA}^8$ .

#### HYDROGEN

- HYDROGEN\_SMOOTHED3  
HYDROGEN\_SMOOTHED5

$$U = \frac{p_0}{r^{12}} - \frac{p_1}{r^{10}} \quad (3.100)$$

2 arguments:  $p_0/k_B$  in units of  $\text{K\AA}^{12}$ , and  $p_1/k_B$  in units of  $\text{K\AA}^{10}$ .

### 3.3.2 Tail corrections

energy

$$U^{\text{Tail}} = \frac{2\pi}{V} \sum_a \sum_b N_a N_b \left[ \int_{r_c}^{\infty} r^2 U(r) dr \right] \quad (3.101)$$

potential	$\int_{r_c}^{\infty} r^2 U(r) dr$
LENNARD_JONES	$\frac{4}{3} p_0 p_1^3 \left[ \frac{1}{3} \left( \frac{p_1}{r} \right)^9 - \left( \frac{p_1}{r} \right)^3 \right]$
LENNARD_JONES_SHIFTED_FORCE	-

**pressure**

$$P^{\text{Tail}} = - \sum_a \sum_b \frac{2\pi}{3V} N_a N_b \left[ \int_{r_c}^{\infty} r^2 r \frac{\partial U(r)}{\partial r} dr \right] \quad (3.102)$$

$$= \sum_a \sum_b \left( \frac{2\pi}{3V} r_c^3 N_a N_b U(r_c) + U^{\text{Tail}} \right) \quad (3.103)$$

**chemical potential**

$$\beta \mu^{\text{Tail}} = 2U^{\text{Tail}} \quad (3.104)$$

### 3.3.3 Electrostatics

**Charge-charge interaction**

- Ewald

The potential energy for a system of charges in a periodic system can be written as

$$U = U^{\text{real}} + U^{\text{rec}} \quad (3.105)$$

where

$$\begin{aligned} U^{\text{real}} &= \sum_{i < j} q_i q_j \frac{\text{erfc}(\alpha r_{ij})}{r_{ij}} \\ U^{\text{rec}} &= \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{1}{k^2} e^{-\frac{k^2}{4\alpha^2}} \left( \left| \sum_{i=1}^N q_i \cos(\mathbf{k} \cdot \mathbf{r}_i) \right|^2 + \left| \sum_{i=1}^N q_i \sin(\mathbf{k} \cdot \mathbf{r}_i) \right|^2 \right) - \sum_i \frac{\alpha}{\sqrt{\pi}} q_i^2 \end{aligned} \quad (3.106)$$

where  $q_i$  and  $q_j$  are the charges of particle  $i$  and  $j$ , respectively,  $\mathbf{r}_i$  the position of atom  $i$ ,  $V$  the volume of the cell,  $\alpha$  a damping factor,  $k$  the wavelength, and 'erfc' the error function complement. The expression gives the *exact* solution for charges in a periodic system up to arbitrary precision. One part is computed in 'real' space, and the long-range part is more conveniently computed in Fourier space.

- CoulombTruncated

$$U = \begin{cases} \sum_{i < j} \frac{1}{4\pi\epsilon} \frac{q_i q_j}{r_{ij}} & \text{if } r_{ij} \leq r_c \\ 0 & \text{otherwise} \end{cases} \quad (3.107)$$

- CoulombShifted

$$U = \begin{cases} \sum_{i < j} \frac{q_i q_j}{4\pi\epsilon} \left( \frac{1}{r_{ij}} - \frac{1}{r_c} \right) & \text{if } r_{ij} \leq r_c \\ 0 & \text{otherwise} \end{cases} \quad (3.108)$$

- CoulombSmoothed

- Wolf

**Charge-dipole interaction**

- Ewald

- CoulombTruncated

$$U = \begin{cases} \sum_{i,j} \frac{1}{4\pi\epsilon} \frac{-q_i}{r_{ij}^2} (\boldsymbol{\mu}_j \cdot \mathbf{r}_{ij}) & \text{if } r_{ij} \leq r_c \\ 0 & \text{otherwise} \end{cases} \quad (3.109)$$

## Dipole-dipole interaction

- Ewald

- CoulombTruncated

$$U = \begin{cases} \sum_{i,j} \frac{1}{4\pi\epsilon} \frac{1}{r_{ij}^3} \left[ \boldsymbol{\mu}_i \cdot \boldsymbol{\mu}_j - 3 \frac{(\boldsymbol{\mu}_i \cdot \mathbf{r}_{ij})(\mathbf{r}_{ij} \cdot \boldsymbol{\mu}_j)}{r_{ij}^2} \right] & \text{if } r_{ij} \leq r_c \\ 0 & \text{otherwise} \end{cases} \quad (3.110)$$

## 3.4 Bonded potentials cross terms

### 3.4.1 Bond-bond potential

- CFF\_BOND\_BOND\_CROSS,CVFF\_BOND\_BOND\_CROSS

$$U = p_0 (r - p_1) (r' - p_2) \quad (3.111)$$

3 arguments:  $p_0/k_B$  in units of K/Å<sup>2</sup>,  $p_0$  and  $p_1$  in Å.

### 3.4.2 Bond-bend potential

- CFF\_BOND\_BEND\_CROSS,CVFF\_BOND\_BEND\_CROSS

$$U = (\theta - p_0) [p_1 (r - p_2) + p_3 (r' - p_4)] \quad (3.112)$$

5 arguments:  $p_0$  in degrees,  $p_1/k_B$  in units of K/Å/rad,  $p_2$  in Å,  $p_3/k_B$  in units of K/Å/rad,  $p_4$  in Å.

- MM3\_BOND\_BEND\_CROSS

$$U = p_0 [(r - p_1) + (r' - p_2)] (\theta - p_3) \quad (3.113)$$

4 arguments:  $p_0$  in mdyne/rad,  $p_1$  and  $p_2$  in Å, and  $p_3$  in degrees.

- TRUNCATED\_HARMONIC

$$U = \frac{1}{2} p_0 (\theta - p_1)^2 e^{-\frac{r_{ij}^8 + r_{ik}^8}{p_2^8}} \quad (3.114)$$

3 arguments:  $p_0/k_B$  in K/rad<sup>2</sup>,  $p_1$  in degrees, and  $p_2$  in units of Å.

- SCREENED\_HARMONIC

$$U = \frac{1}{2} p_0 (\theta - p_1)^2 e^{-\left(\frac{r_{ij}}{p_2} + \frac{r_{ik}}{p_3}\right)} \quad (3.115)$$

4 arguments:  $p_0$  in K/rad<sup>2</sup>,  $p_1$  in degrees,  $p_2$  and  $p_3$  in units of Å.

- SCREENED\_VESSAL

$$U = \frac{p_0}{8(\theta_{ijk} - \pi)^2} \left[ (p_1 - \pi)^2 - (\theta_{ijk} - \pi)^2 \right]^2 e^{-\left(\frac{r_{ij}}{p_2} + \frac{r_{ik}}{p_3}\right)} \quad (3.116)$$

4 arguments:  $p_0$  in K/rad<sup>2</sup>,  $p_1$  in degrees,  $p_2$  and  $p_3$  in units of Å.

- TRUNCATED\_VESSAL

$$U = p_0 \left[ \theta_{ijk}^{p_2} (\theta_{ijk} - p_1)^2 (\theta_{ijk} + p_1 - 2\pi)^2 - \frac{p_2}{2} \pi^{p_2-1} (\theta_{ijk} - p_1)^2 (\pi - p_1)^3 e^{-\frac{r_{ij}^8 + r_{ik}^8}{p_3^8}} \right] \quad (3.117)$$

4 arguments:  $p_0$  in K/rad<sup>4+p2</sup>,  $p_1$  in degrees,  $p_2$  dimensionless, and  $p_3$  in Å.

### 3.4.3 Bend-bend potential

- CFF\_BEND\_BEND\_CROSS,CVFF\_BEND\_BEND\_CROSS

$$U = p_0 (\theta - p_1) (\theta' - p_2) \quad (3.118)$$

3 arguments:  $p_0$  in units of K/rad<sup>2</sup>,  $p_1$  and  $p_2$  in units of degrees.

- MM3\_BEND\_BEND\_CROSS

$$U = -p_0 (\theta - p_1) (\theta' - p_2) \quad (3.119)$$

3 arguments:  $p_0$  in units of mdyne/rad<sup>2</sup>,  $p_1$  and  $p_2$  in units of degrees.

### 3.4.4 Bond-torsion potential

The bond-torsions potential correlates the torsion  $i - j - k - l$  with the central bond  $j - k$ , or with the two terminating bonds.

- MM3\_BOND\_TORSION\_CROSS

The MM3 bond-torsion potential correlates the torsion  $i - j - k - l$  with the central bond  $j - k$

$$U = \frac{1}{2}p_0 (r - p_3) (1 + \cos \phi) + \frac{1}{2}p_1 (r - p_3) (1 + \cos 2\phi) + \frac{1}{2}p_2 (r - p_3) (1 + \cos 3\phi) \quad (3.120)$$

4 arguments:  $p_0, p_1, p_2$  in units of kcal/mol,  $p_3$  the reference length of the central bond in Å.

### 3.4.5 Bend-torsion potential

- CFF\_BEND\_TORSION\_CROSS,CVFF\_BEND\_TORSION\_CROSS

$$U = p_0 (\theta - p_1) (\theta' - p_2) \cos \phi \quad (3.121)$$

3 arguments:  $p_0$  in units of K/rad<sup>3</sup>,  $p_1$  and  $p_2$  in units of degrees.

- SMOOTHED\_DIHEDRAL

$$U = p_0 (1 + \cos(p_1 \phi_{ijkl} - p_2)) S(\theta_{ijk}) S(\theta_{jkl}) \quad (3.122)$$

3 arguments:  $p_0/k_B$  in units of K/rad<sup>2</sup>,  $p_1$  dimensionless, and  $p_2$  in degrees.

- SMOOTHED\_THREE\_COSINE\_DIHEDRAL

$$U = \left\{ \frac{1}{2}p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2}p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2}p_2 [1 + \cos(3\phi_{ijkl})] \right\} S(\theta_{ijk}) S(\theta_{jkl}) \quad (3.123)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- SMOOTHED\_CFF\_DIHEDRAL

$$U = \{p_0 [1 - \cos(\phi_{ijkl})] + p_1 [1 - \cos(2\phi_{ijkl})] + p_2 [1 - \cos(3\phi_{ijkl})]\} S(\theta_{ijk}) S(\theta_{jkl}) \quad (3.124)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K.

- SMOOTHED\_CFF\_DIHEDRAL2

$$U = \{p_0 [1 + \cos(\phi_{ijkl})] + p_1 [1 + \cos(2\phi_{ijkl})] + p_2 [1 + \cos(3\phi_{ijkl})]\} S(\theta_{ijk}) S(\theta_{jkl}) \quad (3.125)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K/rad.

- NICHOLAS\_DIHEDRAL

$$U = \left\{ \frac{1}{2}p_0 [1 + \cos(\phi_{ijkl})] + \frac{1}{2}p_1 [1 - \cos(2\phi_{ijkl})] + \frac{1}{2}p_2 [1 + \cos(3\phi_{ijkl})] \right\} S(\theta_{ijk}) \quad (3.126)$$

3 arguments:  $p_0/k_B, p_1/k_B, p_2/k_B$  in units of K/rad.

- SMOOTHED\_CFF\_BEND\_TORSION\_CROSS

$$U = S(\theta_1) [p_0 * (\text{Theta}_1 - p_1) * (\theta_2 - p_2) \cos(\phi)] S(\theta_2) \quad (3.127)$$

3 arguments:  $p_0/k_B$  in units K/rad<sup>3</sup>,  $p_1$  and  $p_2$  in units of degrees.

The smoothing function  $S(\theta)$  is defined as

$$S(\theta) = \begin{cases} 1 & \theta < \theta_{\text{on}} \\ (\theta_{\text{off}} - \theta)^2 \frac{\theta_{\text{off}} + 2\theta - 3\theta_{\text{on}}}{(\theta_{\text{off}} - \theta_{\text{on}})^3} & \theta \geq \theta_{\text{on}} \end{cases} \quad (3.128)$$

with  $\theta_{\text{on}} = 170^\circ$  and  $\theta_{\text{off}} = 180^\circ$ .

## 3.5 Polarization

### 3.5.1 Introduction

The total potential energy of a set of atoms possessing charges, dipoles, and anisotropic polarizabilities can be expressed as

$$U = U_{\text{bonded}} + U_{\text{nonbonded}} + U_{\text{polarization}} \quad (3.129)$$

The non-bonded energy arising from charges and dipoles is given by

$$U_{\text{nonbonded}} = \frac{1}{2} \sum_{i,j \neq i} q_i T_{ij} q_j + q_i T_{ij}^\alpha \mu_{j,\alpha} - \mu_{i,\alpha} T_{ij}^\alpha q_j + \mu_{i,\alpha} T^{\alpha\beta} \mu_{j,\beta} \quad (3.130)$$

and the polarization energy (the work to form induces dipoles) by

$$U_{\text{polarization}} = \frac{1}{2} \sum_i \mu_{i,\alpha}^{\text{ind}} \alpha_{i,\alpha\beta}^{-1} \mu_{i,\beta}^{\text{ind}} \quad (3.131)$$

where  $q_i$  is the charge of atom  $i$ ,  $\mu_{i,\alpha}$  the  $\alpha$ -component of the total dipole moment of atom  $i$ ,  $\mu_{i,\alpha}^{\text{ind}}$  the  $\alpha$ -component of the induced dipole moment of atom  $i$ , and  $\alpha_{i,\alpha\beta}$  the  $\alpha\beta$ -component of the polarizability tensor of atom  $i$ . Throughout, the Einstein summation convention is used for Greek letter indices. The total dipole moment of atom  $i$  is given by

$$\mu_{i,\alpha} = \mu_{i,\alpha}^{\text{stat}} + \mu_{i,\alpha}^{\text{ind}} \quad (3.132)$$

The interaction tensors are defined as

$$\mathcal{T}_{ij} = \frac{1}{r_{ij}} \quad (3.133)$$

$$\mathcal{T}_{ij}^\alpha = \nabla_\alpha \mathcal{T}_{ij} = -\frac{r_{ij,\alpha}}{r_{ij}^3} \quad (3.134)$$

$$\mathcal{T}_{ij}^{\alpha\beta} = \nabla_\alpha \nabla_\beta \mathcal{T}_{ij} = \frac{3r_{ij,\alpha} r_{ij,\beta}}{r_{ij}^5} - \frac{\delta_{\alpha\beta}}{r_{ij}^3} \quad (3.135)$$

$$\mathcal{T}_{ij}^{\alpha\beta\gamma} = \nabla_\alpha \nabla_\beta \nabla_\gamma \mathcal{T}_{ij} = -\frac{15r_{ij,\alpha} r_{ij,\beta} r_{ij,\gamma}}{r_{ij}^7} + \frac{3(\delta_{\alpha\beta} r_{ij,\gamma} + \delta_{\alpha\gamma} r_{ij,\beta} + \delta_{\beta\gamma} r_{ij,\alpha})}{r_{ij}^5} \quad (3.136)$$

Here,  $r_{ij,\alpha} = r_{j\alpha} - r_{i\alpha}$ , where  $r_{i\alpha}$  is the  $\alpha$ -component of the position vector atom  $i$ .  
The electrostatic potential, field, and gradient are given by

$$\phi_i \equiv \phi(\mathbf{r}_i) = \sum_{j \neq i} (T_{ij} q_j + T_{ij}^\alpha \mu_{j,\alpha}) \quad (3.137)$$

$$E_{i,\alpha} \equiv E_\alpha(\mathbf{r}_i) = \sum_{j \neq i} (T_{ij}^\alpha q_j + T_{ij}^{\alpha\beta} \mu_{j,\beta}) \quad (3.138)$$

$$E_{i,\alpha\beta} \equiv E_{\alpha\beta}(\mathbf{r}_i) = - \sum_{j \neq i} (T_{ij}^{\alpha\beta} q_j + T_{ij}^{\alpha\beta\gamma} \mu_{j,\gamma}) \quad (3.139)$$

The induced dipole moments are determined by a minimization of  $U$  with respect to  $\mu_{i,\alpha}^{\text{ind}}$ ,

$$\mu_{i,\alpha}^{\text{ind}} = \alpha_{i,\alpha\beta} E_{i,\beta} \quad (3.140)$$

### 3.5.2 Computational scheme

The potential energy can be separated into two contributions: an electrostatic and an induction term:

$$U = U_{\text{elec}} + U_{\text{ind}} \quad (3.141)$$

where the electrostatic energy and induction energy are given by

$$U_{\text{elec}} = \frac{1}{2} \sum_i (q_i \phi_i^{\text{stat}} - \mu_{i,\alpha}^{\text{stat}} E_{i,\alpha}^{\text{stat}}) \quad (3.142)$$

$$U_{\text{ind}} = - \frac{1}{2} \sum_i \mu_{i,\alpha}^{\text{ind}} E_{i,\alpha}^{\text{stat}} \quad (3.143)$$

with

$$\phi_i^{\text{stat}} = \sum_{ij} (T_{ij} q_j + T_{ij}^\alpha \mu_{j,\alpha}^{\text{stat}}) \quad (3.144)$$

$$E_{i,\alpha}^{\text{stat}} = \sum_{j \neq i} (T_{ij}^\alpha q_j + T_{ij}^{\alpha\beta} \mu_{j,\beta}^{\text{stat}}) \quad (3.145)$$

The force on atom  $i$  arising from atom  $j$  is

$$f_{ij,\alpha} = q_i T_{ij}^\alpha q_j + q_i T_{ij}^{\alpha\beta} \mu_{j,\beta} - \mu_{i,\beta} T_{ij}^{\beta\alpha} q_j - \mu_{i,\beta} T_{ij}^{\beta\alpha\gamma} \mu_{j,\gamma} \quad (3.146)$$

and the total force on atom  $i$  can be expressed as

$$f_{i,\alpha} = \sum_{j \neq i} q_i E_{i,\alpha} + \mu_{i,\beta} E_{i,\beta\alpha} \quad (3.147)$$

where  $E_{i,\alpha}$  is the  $\alpha$ -component of the electrostatic field, and  $E_{i,\beta\alpha}$  the  $\alpha\beta$ -component of the field gradient on atom  $i$  arising from the other charges and total dipole moments, i.e. Eq 3.138 and 3.139.

The procedure to compute the electrostatic and induction energy is

1. Compute the electrostatic field  $E_{i,\alpha}^{\text{stat}}$  according to

$$E_{i,\alpha}^{\text{stat}} = \sum_{j \neq i} (T_{ij}^\alpha q_j + T_{ij}^{\alpha\beta} \mu_{j,\beta}^{\text{stat}}) \quad (3.148)$$

2. Calculate the induced dipole moments  $\mu_{i,\alpha}^{\text{ind}}$  by self-consistent iterations using

$$\begin{aligned}\mu_{i,\alpha}^{\text{ind}} &= \alpha_{i,\alpha\beta} E_{i,\beta} \\ E_{i,\alpha} &= E_{i,\alpha}^{\text{stat}} + E_{i,\alpha}^{\text{ind}} \quad \text{with } E_{i,\alpha}^{\text{ind}} = \sum_{j \neq i} T_{ij}^{\alpha\beta} \mu_{j,\beta}^{\text{ind}}\end{aligned}\tag{3.149}$$

3. Calculate the electrostatic energy and induction energy using

$$\begin{aligned}U_{\text{elec}} &= \frac{1}{2} \sum_i (q_i \phi_i^{\text{stat}} - \mu_{i,\alpha}^{\text{stat}} E_{i,\alpha}^{\text{stat}}) \\ U_{\text{ind}} &= -\frac{1}{2} \sum_i \mu_{i,\alpha}^{\text{ind}} E_{i,\alpha}^{\text{stat}}\end{aligned}\tag{3.150}$$

4. Calculate the electrostatic field gradient  $E_{i,\alpha}^{\alpha\beta}$  from charge and total dipole moments using

$$E_{i,\alpha}^{\alpha\beta} = - \sum_{j \neq i} (T_{ij}^{\alpha\beta} q_j + T_{ij}^{\alpha\beta\gamma} \mu_{j,\gamma})\tag{3.151}$$

5. Calculate the forces according to

$$f_{i,\alpha} = \sum_{j \neq i} q_i E_{i,\alpha} + \mu_{i,\beta} E_{i,\beta\alpha}\tag{3.152}$$

## Bibliography

- [1] G. Mie. Kinetic theory of one-atom substance. *Ann. Physik*, 11:657, 1903.

# Chapter 4

## Examples

### 4.1 Introduction

Often the best way of learning a program is to look at various examples. Note these examples are just for that purpose and real simulation runs should be much longer, both in initialization time as well as production run time.

Tip: VMD is capable of showing pdb-files with several frames. This is the way RASPA produces movies. Standard VMD does not show the box itself but some extension scripts have been written. To show the unit cell in VMD you can input into the console:

```
draw pbctools.tcl
```

make sure the 'pbctools.tcl' and 'pbctools.tcl' are in the current directory, they are located in the 'utils' directory of RASPA. For NPT simulations the box is properly updated.

The output-files begin with some essential data about the program: the version number, whether a 64-bits or 32-bits executable is run, the used compiler, when the output-file was generated and on which node and system.

```
Compiler and run-time data
=====
RASPA 0.99-480
Compiled as a 64-bits application
Compiler: gcc 4.2.1 (Based on Apple Inc. build 5658) (LLVM build 2336.1.00)
Compile Date = Dec 9 2011, Compile Time = 17:21:59

Fri Dec 9 18:03:35 2011
Simulation started on Friday, December 09.
The start time was 06:03 PM.

Cpu data: x86_64
Cpu Model: MacBookPro6,2
Host name: Davids-MacBook-Pro.local
OS release: 11.2.0
OS type: Darwin
OS version: 11C74
```

## 4.2 Basic examples

### Example 1: Monte Carlo of methane in a box

A Monte Carlo run of 100 methane molecules in a  $30 \times 30 \times 30$  Å box. After 1000 cycles of initialization the production run is started. A movie is written and every 10th configuration is appended to the movie. The movie is stored in 'Movies/System\_0', and can be viewed with VMD: 'vmd AllComponents.pdb'.

```

SimulationType          MonteCarlo
NumberOfCycles         10000
NumberOfInitializationCycles 1000
PrintEvery             100

Forcefield              GarciaPerez2006

Box 0
BoxLengths 30 30 30
ExternalTemperature 300.0
Movies yes
WriteMoviesEvery 100

Component 0 MoleculeName      methane
           MoleculeDefinition   TraPPE
           TranslationProbability 1.0
           CreateNumberOfMolecules 100

```

In RASPA, the cycle is define as  $\max(20, N)$  steps, where  $N$  is the number of molecules in the system. In every cycle, each of the molecules has on average been used for a Monte Carlo move (accepted or rejected). There is a minimum of 20 steps to avoid that low-density systems or not sampled well. The definition of a cycle is less dependent on the system size. The number of Monte carlo steps is roughly the the number of cycles times the average number of molecules.

In the output file the simulation writes an important check to the file

```

Energy-drift status
=====
Adorbate/Adsorbate energy-drift:          -8.33097e-10
  Adorbate/Adsorbate VDW energy-drift:    -8.33097e-10
=====
Total energy-drift: -9.84073e-10

```

In Monte Carlo only difference in energies are computed. These differences are continuously added to keep track of the current energies (from which average energies etc. are computed). Obviously, the current energy that is kept track off during the simulation should be equal to a full recalculation of the energies. The difference between the two signals an error. If the drift is higher than say  $1e-3$  or  $1e-4$  the results of the simulation are in error. This could be due to an error in one of the Monte Carlo moves or because the force field is "wrong" (a typical error is when one forgets to define required potentials).

The performance of Monte Carlo moves is monitored. Translation moves are usually scaled to achieve an acceptance rate of 50%. Here, the move reached its upper limit of 1 Å because of the low density of the system.

```

Performance of the translation move:
=====
Component 0 [methane]
  total      329740.000000 330574.000000 329274.000000
  successfull 283299.000000 283739.000000 283304.000000
  accepted   0.859159 0.858322 0.860390
  displacement 1.000000 1.000000 1.000000

```

Averages are computed along with an error bar. The error is computed by dividing the simulation in 5 blocks and calculating the standard deviation. The errors in RASPA are computed as the 95% confidence interval.

```

Total energy:
=====
  Block[ 0]      -18003.83404 [K]
  Block[ 1]      -18179.35286 [K]
  Block[ 2]      -18342.84622 [K]
  Block[ 3]      -18951.71107 [K]
  Block[ 4]      -18345.26757 [K]
=====
Average       -18364.60235 [K] +/-      638.75814 [K]

```

### Example 2: Monte Carlo of CO<sub>2</sub> in a box and N<sub>2</sub> in another box (two independent simulations)

RASPA has a build-in structure of being able to simulate several systems at the same time. This has applications in Gibbs-ensembles and (hyper) parallel tempering for example. However, this capability can

also be used for independent systems. The first box is  $30 \times 30 \times 30$  Å with  $90^\circ$  angles, containing 50 N<sub>2</sub> and 25 CO<sub>2</sub> and molecules and moved around by translation, rotation and reinsertion. The second box is monoclinic and of size  $25 \times 25 \times 25$  with  $\beta = 120^\circ$ ,  $\alpha = \gamma = 90^\circ$  containing 25 N<sub>2</sub> and 50 CO<sub>2</sub> molecules. The first system is at 300K, the second at 500K.

```

SimulationType MonteCarlo
NumberOfCycles 10000
NumberOfInitializationCycles 1000
PrintEvery 100

Forcefield GarciaPerez2006

Box 0
BoxLengths 25 25 25
ExternalTemperature 300.0
Movies yes
WriteMoviesEvery 10

Box 1
BoxLengths 30 30 30
BoxAngles 1
90 120 90
ExternalTemperature 500.0
Movies yes
WriteMoviesEvery 10

Component 0 MoleculeName N2
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
RegrowProbability 1.0
CreateNumberOfMolecules 50 25

Component 1 MoleculeName CO2
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
RegrowProbability 1.0
CreateNumberOfMolecules 25 50

```

One thing to note is that system-dependent statements apply to the *current* box, following 'Box [int]'. The initialization of the systems with molecules is done using the 'CreateNumberOfMolecules' which applies similarly to the *current* component specified using 'component [int]'. The list of integers represent the initial amount of molecules for each system. Note that when the 'BoxAngles' line is omitted,  $\alpha = \beta = \gamma = 90^\circ$  is assumed as the default.

### Example 3: Monte Carlo of a binary mixture in a box

A Monte Carlo run of 50 propane and 50 butane molecules in a  $30 \times 30 \times 30$  Å box. The MC moves are translation, rotation, and full reinsertion. After 1000 steps of initialization the production run is started. A movie is written and every 10th configuration is appended to the movie. The movie is stored in 'Movies/System[0]', and can be viewed with VMD: 'vmd AllComponents.pdb'.

```

SimulationType MonteCarlo
NumberOfCycles 10000
NumberOfInitializationCycles 2000
PrintEvery 100

Forcefield GarciaPerez2006

Box 0
BoxLengths 30 30 30
ExternalTemperature 300.0
Movies yes
WriteMoviesEvery 10

Component 0 MoleculeName propane
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 50

Component 1 MoleculeName butane
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 50

```

### Example 4: Monte Carlo of CO<sub>2</sub> and N<sub>2</sub> in two independent boxes

An example of a binary mixture of CO<sub>2</sub> and N<sub>2</sub> in two independent boxes. Box one contains 100 CO<sub>2</sub> molecules at 300 Kelvin, box two (monoclinic shape) contains 100 N<sub>2</sub> molecules at 500 Kelvin. The movies

for box one are appended every 10 cycles, the movie for box two every 5 cycles. Three types of Monte Carlo moves are used: translation, rotation, and reinsertion. The force field used is the TraPPE force field.

```

SimulationType MonteCarlo
NumberOfCycles 10000
NumberOfInitializationCycles 1000
PrintEvery 100

Forcefield GarciaPerez2006

Box 0
BoxLengths 25 25 25
ExternalTemperature 300.0
Movies yes
WriteMoviesEvery 10

Box 1
BoxLengths 30 30 30
BoxAngles 90 120 90
ExternalTemperature 500.0
Movies yes
WriteMoviesEvery 5

Component 0 MoleculeName CO2
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 100 0

Component 1 MoleculeName N2
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 0 100

```

### Example 5: Molecular dynamics of methane in a box measuring the mean-square displacement

A molecular dynamics run of 100 methane molecules in a  $25 \times 25 \times 25$  Å box at 300 K. The simulations starts with 1000 InitializationSteps using Monte Carlo, the only MC moves are translation and reinsertion. After 1000 steps of initialization the equilibration run is started. Here, the atoms are assigned a velocities, and during the equilibration run the distribution should attain the Maxwell-Boltzmann distribution. After the initialization and equilibration runs, the production is started. The mean-square displacement is measured and written to 'MSDOrderN/System[0]' for both self-and collective diffusion (the slope of the mean square displacement is related to the diffusion coefficients). They can be plotted with 'gnuplot'. In contrast to Monte Carlo where the ensemble basically follows from the used MC moves, the ensemble for molecular dynamics needs to be explicitly specified using the 'Ensemble' keyword.

```

SimulationType MolecularDynamics
NumberOfCycles 1000000
NumberOfInitializationCycles 1000
NumberOfEquilibrationCycles 10000
PrintEvery 1000

Ensemble NVT
TimeStep 0.0005

Forcefield GarciaPerez2006

Box 0
BoxLengths 25 25 25
ExternalTemperature 300.0
ComputeMSD yes
PrintMSDEvery 5000

Component 0 MoleculeName methane
MoleculeDefinition TraPPE
TranslationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 100

```

In MD, it is important to have good energy-conservation. This is monitored

```
Conserved energy: 5370.3533927734 Energy drifts: 0.0036722592 0.0022292104
```

The first number is the conserved quantity, the second the current relative energy drift, and the last number is the average energy drift. The latter two numbers need to be small, usually smaller than say  $10^{-3}$ . Here, the number is large which is due to the type of used potential in the TraPPE forcefield: unshifted, truncated Lennard-Jones with tail-corrections. Shifted potentials show much better energy conservation because they remove the discontinuity of the force at the cutoff boundary.

Using NVE, temperature control is difficult, the average temperature was  $253.19590 \pm 0.77596$ .

## Example 6: Adsorption isotherm of methane in MFI

Adsorption isotherms can be easily obtained by specifying a list of (increasing) pressures which will be subsequently run. If no 'FugacityCoefficient' keyword is specified these pressure are converted to fugacity using the Peng-Robinson equation of state. Important: it is essential to specify the 'ideal gas Rosenbluth weight' for a component. This value needs to be computed separately and depends only on temperature (see auxiliary examples). This value is the reference state of the ideal gas. It is convenient to specify it in advance, otherwise the correct pressure needs to deduced afterwards and is different from the specified input. For mixtures this becomes cumbersome when the ideal gas Rosenbluth weight of the components is different. It is also convenient to specify the 'void fraction' of the materials (probed with helium) in advance (see auxiliary examples). If you do, the excess adsorption is automatically computed properly. At high pressures and temperatures the excess adsorption can be substantially lower than absolute adsorption. In this example,  $2 \times 2 \times 2$  unit cells are required to meet the required that all of the perpendicular cell lengths are larger than twice the cutoff distance. The default cutoff of 12 Å means the perpendicular lengths should be larger than 24 Å.

```

SimulationType MonteCarlo
NumberOfCycles 25000
NumberOfInitializationCycles 2000
PrintEvery 1000

ContinueAfterCrash no
WriteBinaryRestartFileEvery 2000

Forcefield GarciaPerez2006
RemoveAtomNumberCodeFromLabel yes

Framework 0
FrameworkName MFI_SI
UnitCells 2 2 2
HeliumVoidFraction 0.29
ExternalTemperature 300.0
ExternalPressure 10000.0 20000.0 30000.0 40000.0

ComputeNumberOfMoleculesHistogram yes
WriteNumberOfMoleculesHistogramEvery 5000
NumberOfMoleculesHistogramSize 1100
NumberOfMoleculesRange 80

ComputeEnergyHistogram yes
WriteEnergyHistogramEvery 5000
EnergyHistogramSize 400
EnergyHistogramLowerLimit -110000
EnergyHistogramUpperLimit -20000

Component 0 MoleculeName methane
    MoleculeDefinition TraPPE
    TranslationProbability 0.5
    ReinsertionProbability 0.5
    SwapProbability 1.0
    CreateNumberOfMolecules 0

```

The output-file shows the performance of the various Monte Carlo moves. For adsorption, a good check is that the acceptance ratio of the 'swap addition' and the 'swap deletion' should be close.

```

Performance of the swap addition move:
=====
Component [methane] total tried: 126109.000000 succesfull growth: 109688.000000 (86.978725 [%]) accepted: 39558.000000 (31.368102 [%])

Performance of the swap deletion move:
=====
Component [methane] total tried: 126236.000000 succesfull growth: 126236.000000 (100.000000 [%]) accepted: 39550.000000 (31.330207 [%])

Performance of the regrow move:
=====
Component [methane] total tried: 125396.000000 succesfull growth: 109358.000000 87.210118 [%] accepted: 24240.000000 (19.330760 [%])

```

Adsorption results are displayed in various units for both absolute and excess adsorption.

```

Component 0 [methane]
-----
Block[ 0] 15.09680      [-]
Block[ 1] 15.14960      [-]
Block[ 2] 15.22480      [-]
Block[ 3] 15.34960      [-]
Block[ 4] 15.11780      [-]

-----  

Average          15.1877200000 +/-   0.1837137948 [-]
Average loading absolute [molecules/unit cell] 1.8984650000 +/-   0.0229642244 [-]
Average loading absolute [mol/kg framework] 0.3291321873 +/-   0.0039812508 [-]
Average loading absolute [milligram/gram framework] 5.2800899500 +/-   0.0638690575 [-]

Block[ 0] 15.09680      [-]
Block[ 1] 15.14960      [-]
Block[ 2] 15.22480      [-]
Block[ 3] 15.34960      [-]
Block[ 4] 15.11780      [-]

```

```

-----+
Average 15.0681524193 +/- 0.1837137948 [-]
Average loading excess [molecules/unit cell] 1.8835190524 +/- 0.0229642244 [-]
Average loading excess [mol/kg framework] 0.3265410453 +/- 0.0039812508 [-]
Average loading excess [milligram/gram framework] 5.23885216579 +/- 0.0638690575 [-]

```

### Example 7: Henry coefficient of *n*-hexane in mono-clinic ERI

The monoclinic version of erionite (ERI) is named 'ERI\_mono', the orthorhombic version is 'ERI'. The monoclinic version needs at least  $3 \times 3 \times 3$  unit cells to be larger than twice the cutoff, while the orthorhombic needs  $2 \times 2 \times 2$  (the unit cell shapes and size are different). To compute the Henry coefficient of hexane in erionite two simulations need to be performed. First the ideal Rosenbluth gas value needs to be computed at the desired temperature (see Auxiliary examples). This value needs to be filled in first. Next the simulation is started and the Henry coefficient is listed in the output.

```

SimulationType MonteCarlo
NumberOfCycles 20000
NumberOfInitializationCycles 0
PrintEvery 1000
PrintPropertiesEvery 1000

Forcefield GarciaPerez2006

Framework 0
FrameworkName ERI_SI
RemoveAtomNumberCodeFromLabel yes
UnitCells 3 3 3
ExternalTemperature 573.0

Component 0 MoleculeName hexane
MoleculeDefinition TraPPE
IdealRosenbluthValue 0.00312147
WidomProbability 1.0
CreateNumberOfMolecules 0

```

The average Widom Rosenbluth weight and Henry coefficient are printed:

```

Average Widom Rosenbluth factor:
=====
Block[ 0] 1.00749 [-]
Block[ 1] 0.996774 [-]
Block[ 2] 1.00742 [-]
Block[ 3] 0.995992 [-]
Block[ 4] 1.01246 [-]

[hexane] Average Widom: 1.00403 +/- 0.013018 [-]

Average Henry coefficient:
=====
Block[ 0] 1.35128e-07 [mol/kg/Pa]
Block[ 1] 1.33692e-07 [mol/kg/Pa]
Block[ 2] 1.3512e-07 [mol/kg/Pa]
Block[ 3] 1.33587e-07 [mol/kg/Pa]
Block[ 4] 1.35796e-07 [mol/kg/Pa]

[hexane] Average Henry coefficient: 1.34664e-07 +/- 1.746e-09 [mol/kg/Pa]

```

### Example 8: Henry coefficient of *n*-pentane to *n*-nonane in MFI

By using multiple components several Henry coefficients can be computed simultaneously. The Widom insertion probe move never actually inserts the molecules, it just compute the energy at randomly chosen insertion positions. Note that the ideal gas Rosenbluth weights decrease with chain length.

```

SimulationType MonteCarlo
NumberOfCycles 10000
NumberOfInitializationCycles 0
PrintEvery 1000
PrintPropertiesEvery 1000

Forcefield GarciaPerez2006

Framework 0
FrameworkName MFI_SI
RemoveAtomNumberCodeFromLabel yes
UnitCells 2 2 2
ExternalTemperature 573.0

Component 0 MoleculeName C5
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 0.064
WidomProbability 1.0
CreateNumberOfMolecules 0

Component 1 MoleculeName C6
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 0.0164423
WidomProbability 1.0
CreateNumberOfMolecules 0

```

```

Component 2 MoleculeName      C7
MoleculeDefinition      TraPPE
IdealGasRosenbluthWeight  0.00425143
WidomProbability        1.0
CreateNumberOfMolecules   0

Component 3 MoleculeName      C8
MoleculeDefinition      TraPPE
IdealGasRosenbluthWeight  0.0011068
WidomProbability        1.0
CreateNumberOfMolecules   0

Component 4 MoleculeName      C9
MoleculeDefinition      TraPPE
IdealGasRosenbluthWeight  0.000289648
WidomProbability        1.0
CreateNumberOfMolecules   0

```

The resulting Henry coefficients are:

```

Average Henry coefficient:
=====
[CG] Average Henry coefficient: 2.98386e-06 +/- 4.36063e-08 [mol/kg/Pa]
[CG] Average Henry coefficient: 5.90253e-06 +/- 7.2814e-08 [mol/kg/Pa]
[C7] Average Henry coefficient: 1.17407e-05 +/- 2.48425e-07 [mol/kg/Pa]
[C8] Average Henry coefficient: 2.27983e-05 +/- 7.43978e-07 [mol/kg/Pa]
[C9] Average Henry coefficient: 4.28771e-05 +/- 1.62704e-06 [mol/kg/Pa]

```

### Example 9: Computing the radial distribution function of a methane/ethane-mixture

The radial distribution function (RDF) is a good indication of the status of the fluid: solid, liquid or gas. Here we compute the RDF for an equimolar mixture of (united atom) methane and ethane in the NVT ensemble using MD.

```

SimulationType      MolecularDynamics
NumberofCycles     1000000
NumberofInitializationCycles 1000
NumberofEquilibrationCycles 5000
PrintEvery         1000

ContinueAfterCrash no
WriteBinaryRestartFileEvery 5000

Ensemble          NVT

Forcefield         GarciaPerez2006

Box 0
BoxLengths 24.0 24.0 24.0
ComputeRDF yes
WriteRDFEvery 10000
RDFHistogramSize 300
RDFRange 12.0

ExternalTemperature 300.0

Component 0 MoleculeName      methane
MoleculeDefinition      TraPPE
TranslationProbability  0.5
RotationProbability    0.5
ReinsertionProbability 1.0
CreateNumberOfMolecules 50

Component 1 MoleculeName      ethane
MoleculeDefinition      TraPPE
TranslationProbability  0.5
RotationProbability    0.5
ReinsertionProbability 1.0
CreateNumberOfMolecules 50

```

The RDF is a static property and equally well be computed using MC:

```

SimulationType      MonteCarlo
NumberofCycles     50000
NumberofInitializationCycles 10000
PrintEvery         1000

ContinueAfterCrash no
WriteBinaryRestartFileEvery 5000

Forcefield         GarciaPerez2006

Box 0
BoxLengths 24.0 24.0 24.0
ComputeRDF yes
WriteRDFEvery 10000
RDFHistogramSize 300
RDFRange 12.0

ExternalTemperature 300.0

Component 0 MoleculeName      methane
MoleculeDefinition      TraPPE
TranslationProbability  0.5

```




Figure 4.1: The radial distribution function of (a) methane/ethane at 300K, (b) water at 298K.

```

RotationProbability      0.5
ReinsertionProbability   1.0
CreateNumberOfMolecules 50

Component 1 MoleculeName      ethane
MoleculeDefinition       TraPPE
TranslationProbability    0.5
RotationProbability       0.5
ReinsertionProbability    1.0
CreateNumberOfMolecules 50
  
```

### Example 10: Computing the radial distribution function of liquid water

RASPA computes the RDF for all (pseudo-)atoms pairs, unless you specified 'no' to the 'PrintToPDB'-field of the 'pseudo\_atoms' file. For example, the  $L$ -atoms of water should not be printed to movie-files, and there would be little point generating the RDF for interactions with these 'dummy' interaction sites.

```

SimulationType           MolecularDynamics
NumberOfCycles          25000
NumberOfInitializationCycles 100
NumberOfEquilibrationCycles 5000
PrintEvery               100
RestartFile              yes

Ensemble                 NVT

Forcefield                Local

Box 0
Boxlengths 24.83 24.83 24.83
ComputeRDF yes
WriteRDFEvery 1000
ExternalTemperature 298.0

Component 0 MoleculeName      Tip5p
MoleculeDefinition       Water
TranslationProbability    0.5
RotationProbability       0.5
ReinsertionProbability    1.0
CreateNumberOfMolecules 0
  
```

We used the NVT ensemble and therefore we have good temperature control

```

Temperature:            303.395 (avg. 298.306), Translational (avg. 298.291), Rotational (avg. 298.321)
Temperature Adsorbates: 303.395 (avg. 298.306), Translational (avg. 298.291), Rotational (avg. 298.321)
  
```

Shifted potentials are used and the relative energy conservation is 0.0000159 (excellent).

### Example 11: measuring bond/bend/dihedral angle distributions

The distributions of the bond-lengths, bend-angles, and torsion-angles can be computed in MD using:

```

SimulationType           MD
NumberOfCycles          1000000000
NumberOfInitializationCycles 5000
NumberOfEquilibrationCycles 10000
PrintEvery               1000
RestartFile              no

Ensemble NVT
  
```




Figure 4.2: The bond, bend, and dihedral angle distributions, comparison between Monte-Carlo and Molecular Dynamics.

```

ContinueAfterCrash      no
WriteBinaryRestartFileEvery   5000

Forcefield           GarciaPerez2006

Box 0
Boxlengths 25 25 25
ExternalTemperature 298.0
ExternalPressure 0.0
ComputeMoleculeProperties yes

component 0 MoleculeName      2-methylbutane
          StartingBead    0
          FugacityCoefficient 1.0
          MoleculeDefinition TraPPE
          TranslationProbability 1.0
          RotationProbability 1.0
          RegrowProbability 1.0
          CBMCProbability 1.0
          CreateNumberOfMolecules 32

```

The distributions of the bond-lengths, bend-angles, and torsion-angles can be computed in MC using:

```

SimulationType      MonteCarlo
NumberofCycles     5000000
NumberofInitializationCycles 10000
PrintEvery         5000
RestartFile        no

ContinueAfterCrash  no
WriteBinaryRestartFileEvery 50000

Forcefield           GarciaPerez2006

Box 0
Boxlengths 25 25 25
ExternalTemperature 298.0
ExternalPressure 0.0
ComputeMoleculeProperties yes

component 0 MoleculeName      2-methylbutane
          StartingBead    0
          FugacityCoefficient 1.0
          MoleculeDefinition TraPPE
          TranslationProbability 1.0
          RotationProbability 1.0
          RegrowProbability 1.0
          CBMCProbability 1.0
          CreateNumberOfMolecules 32

```

The relevant files in 'MolecularProperties/System\_0' are plotted in Figure 4.2. In the Monte-Carlo scheme the configurations are generated using configurational-bias Monte-Carlo. As can be seen, the distributions are identical to what MD would give.

### 4.3 Non-basic examples

#### Example 1: Adsorption of a binary CO<sub>2</sub>/CH<sub>4</sub> (1:3) mixture in IRMOF-1

Appreciable adsorption in MOF materials occurs at higher pressure than zeolites, usually in the range up to 10 bar. At these high pressures absolute and excess adsorption are different, and excess adsorption eventually even goes down. This is due to the fact that excess adsorption is relative to what would have been in the free pore volume at these conditions. So one can compress the outside fluid but eventually the pores are filled up. At that maximum absolute loading the excess adsorption will go down.

```

SimulationType MonteCarlo
NumberOfCycles 50000
NumberOfInitializationCycles 5000
PrintEvery 5000

Forcefield IRMOF-1

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1
HeliumVoidFraction 0.81
ExternalTemperature 300.0
ExternalPressure 1000000.0

Component 0 MoleculeName CO2
MoleculeDefinition TraPPE
MolFraction 0.25
TranslationProbability 0.5
ReinsertionProbability 0.5
IdentityChangeProbability 1.0
NumberOfIdentityChanges 2
IdentityChangesList 0 1
SwapProbability 1.0
CreateNumberOfMolecules 0

Component 1 MoleculeName methane
MoleculeDefinition TraPPE
MolFraction 0.75
TranslationProbability 0.5
ReinsertionProbability 0.5
IdentityChangeProbability 1.0
NumberOfIdentityChanges 2
IdentityChangesList 0 1
SwapProbability 1.0
CreateNumberOfMolecules 0

```

To compute the excess adsorption the void fraction of a structure needs to be specified using 'HeliumVoidFraction [real]'. RASPA automatically uses an equation of state (default: Peng-Robinson) to compute the fugacities from the pressure and mol-fraction as is done here for a mixture of CO<sub>2</sub> and CH<sub>4</sub>. It also computes the amount of excess molecules from this equation of state.

```

Component 0 [CO2] (Adsorbate molecule)

Critical temparure [K]: 304.128200
Critical pressure [Pa]: 7377300.000000
Acentric factor [-]: 0.223940

Fluid is a vapour

MolFraction: 0.2500000000 [-]
Compressibility: 0.9714389725 [-]

Density of the bulk fluid phase: 18.1624884819 [kg/m^3]

Amount of excess molecules: 0.8675190741 [-]

Partial pressure: 250000.0000000000 [Pa]
1875.0000000000 [Torr]
2.5000000000 [bar]
2.4673081668 [atm]

Fugacity coefficient: 0.9503504709 [-]

Partial fugacity: 237587.6177345715 [Pa]
1781.9071330093 [Torr]
2.3758761773 [bar]
2.3448074783 [atm]

Component 1 [methane] (Adsorbate molecule)

Critical temparure [K]: 190.564000
Critical pressure [Pa]: 4599200.000000
Acentric factor [-]: 0.011420

Fluid is a vapour

MolFraction: 0.7500000000 [-]
Compressibility: 0.9714389725 [-]

Density of the bulk fluid phase: 6.6206386115 [kg/m^3]

Amount of excess molecules: 2.6025572224 [-]

Partial pressure: 750000.0000000001 [Pa]
5625.0000000000 [Torr]
7.5000000000 [bar]
7.4019245004 [atm]

Fugacity coefficient: 0.9790119494 [-]

Partial fugacity: 734258.9620174330 [Pa]
5506.9422151307 [Torr]
7.3425896202 [bar]
7.2465725341 [atm]

```

Also noteworthy is the use of the identity-change move for mixtures. A molecule of a certain type can be changed at the same position into a molecule of another type. It is specified per component as a list of

other components that are allowed for this move. The identity-change move is highly recommended at high loadings.

At each 'PrintEvery' steps the loadings are shown in a variety of units for both excess and absolute adsorption:

```
Loadings per component:
-----
Component 0 (CO2), current number of molecules: 13 (avg. 14.49236), density: 55.11440 (avg. 61.44138) [kg/m^3]
    absolute adsorption: 13.00000 (avg. 14.49236) [mol/ucl], 2.1108713277 (avg. 2.3531933869) [mol/kg], 92.8983916953 (avg. 103.5628643601) [mg/g]
        47.3130187448 (avg. 52.7444195030) [cm^3 STP/g], 28.0696606694 (avg. 31.2919783377) [cm^3 STP/cm^3]
    excess adsorption: 12.1324809259 (avg. 13.6248441639) [mol/ucl], 1.9700081631 (avg. 2.2123302223) [mol/kg], 86.6990742528 (avg. 97.3635469175) [mg/g]
        44.1557151897 (avg. 49.5871159479) [cm^3 STP/g], 26.1965094359 (avg. 29.4188271042) [cm^3 STP/cm^3]
Component 1 (methane), current number of molecules: 15 (avg. 19.09103), density: 23.18129 (avg. 29.50365) [kg/m^3]
    absolute adsorption: 15.00000 (avg. 19.09103) [mol/ucl], 2.4356207627 (avg. 3.099010314) [mol/kg], 39.0733486608 (avg. 49.7300382994) [mg/g]
        54.5919447055 (avg. 69.4811065367) [cm^3 STP/g], 32.3880700031 (avg. 41.2214467636) [cm^3 STP/cm^3]
    excess adsorption: 12.39744227776 (avg. 16.4884754138) [mol/ucl], 2.0130312689 (avg. 2.6773115375) [mol/kg], 32.2939736100 (avg. 42.9506632486) [mg/g]
        45.1200340402 (avg. 60.0091958713) [cm^3 STP/g], 26.7686163026 (avg. 35.6019930632) [cm^3 STP/cm^3]
```

and at the end error bars are computed for all properties:

```
Component 0 [CO2]
-----
Block[ 0] 14.94810      [-]
Block[ 1] 14.80090      [-]
Block[ 2] 14.57590      [-]
Block[ 3] 14.66720      [-]
Block[ 4] 14.78100      [-]

Average          14.7546200000 +/- 0.2527533944 [-]

Average loading absolute [molecules/unit cell] 14.7546200000 +/- 0.2527533944 [-]
Average loading absolute [mol/kg framework] 2.3957772545 +/- 0.0410407610 [-]
Average loading absolute [milligram/gram framework] 105.4369590827 +/- 1.8061833722 [-]

Average loading excess [molecules/unit cell] 13.8871009259 +/- 0.2527533944 [-]
Average loading excess [mol/kg framework] 2.2549140899 +/- 0.0410407610 [-]
Average loading excess [milligram/gram framework] 99.2376416402 +/- 1.8061833722 [-]

Component 1 [methane]
-----
Block[ 0] 19.36240      [-]
Block[ 1] 19.29190      [-]
Block[ 2] 19.25920      [-]
Block[ 3] 19.17110      [-]
Block[ 4] 19.58860      [-]

Average          19.3346400000 +/- 0.2821765858 [-]

Average loading absolute [molecules/unit cell] 19.3346400000 +/- 0.2821765858 [-]
Average loading absolute [mol/kg framework] 3.1394567082 +/- 0.0458183434 [-]
Average loading absolute [milligram/gram framework] 50.3646086634 +/- 0.7350389415 [-]

Average loading excess [molecules/unit cell] 16.7320827776 +/- 0.2821765858 [-]
Average loading excess [mol/kg framework] 2.7168672144 +/- 0.0458183434 [-]
Average loading excess [milligram/gram framework] 43.5852336126 +/- 0.7350389415 [-]
```

## Example 2: NPT Monte Carlo of propane

The density of propane at 250K and 10 bar is about 559.53 kg/m<sup>3</sup> (NIST database). In this example the density is computed using Monte Carlo in the NPT-ensemble. Given the pressure  $P$ , the temperature  $T$ , and the amount of molecules  $N$ , the density is computed.

```
SimulationType      MonteCarlo
NumberOfCycles     25000
NumberOfInitializationCycles 5000
PrintEvery         100

Forcefield         TraPPE

Box 0
BoxLengths 30 30 30
ExternalTemperature 250.0
ExternalPressure 1000000.0

VolumeChangeProbability 0.25

Component 0 MoleculeName      propane
MoleculeDefinition   TraPPE
TranslationProbability 0.5
ReinsertionProbability 0.5
CreateNumberOfMolecules 256
```

The TraPPE model for propane gives for our simulation of 25000 cycles  $560.84704 \pm 1.72392$  kg/m<sup>3</sup>.

### Example 3: NPT molecular dynamics of water

A molecular dynamics simulation of water in the NPT-ensemble (constant amount of particles  $N$ , constant average pressure  $P$ , and constant average temperature  $T$ ). Many water models are defined, but most are defined with simple Coulombic potentials using cutoffs of 9 Å. None are optimized with the Ewald-summation except for the recalibrated Tip5p-Ew model. Unfortunately, that model is defined using a cutoff always equal to half the box size, while RASPA uses a fixed cutoff (default: 12 Angstrom). A fixed cutoff is more realistic, but requires the shortest perpendicular width to be twice the cutoff, thus here larger than 24 Å. All this results in having to simulate more than 512 water molecules. The tip5p models use 5 fixed charges placed in the water geometry, so for each step 2560 charge sites needs to be computed with Ewald. Conclusion: liquid water is computationally expensive to compute when done properly.

In MD-NPT the average pressure  $\langle P \rangle$  and average temperature  $\langle T \rangle$  are imposed. The instantaneous values, especially for the pressure, are different. RASPA uses the Nose-Hoover chain method, and NPT-MD methods of Martyna and Tuckermann.

Several options are introduced here: "TimeStep [real]" to set the time step. For rigid molecules the time step can be a bit larger because the high frequency movement is removed (the O-H is around 3000 cm<sup>-1</sup>). The cutoff can be set with 'CutOff [real]'. The method to compute charge interactions is set with 'ChargeMethod [Ewald|None]', although Ewald is the default. The precision can specified using 'EwaldPrecision [real]' from which the Ewald parameters  $\kappa$  and the amount of wave vectors is inferred. The initial positions of the water are read from file ('RestartFile yes'), the file is located in directory 'RestartInitial/System[int]'.

The experimental density of water at 300K and 1 bar is about 996.56 kg/m<sup>3</sup> (NIST database).

```

SimulationType           MolecularDynamics
NumberOfCycles          100000
NumberOfInitializationCycles 0
NumberOfEquilibrationCycles 1000
PrintEvery               100
RestartFile              yes

Ensemble                 NPT
TimeStep                  0.001

ChargeMethod              Ewald
CutOff                    10.0
Forcefield                Tip5p-Ew
EwaldPrecision             1e-6

Box 0
BoxLengths   23 23 23
ExternalTemperature 300.0
ExternalPressure 1.0e5
ComputeMSD yes
PrintMSDEvery 5000

Component 0 MoleculeName      Tip5p
StartingBead            0
MoleculeDefinition       Water
TranslationProbability    1.0
RotationProbability       1.0
ReinsertionProbability    1.0
CreateNumberOfMolecules   0

```

The output shows some details of intermediate status during the run: the time run, the current box and average box, etc. The total linear momentum is conserved and zero (the center of mass movement of the system is removed at initialization). For this relatively short run, the average pressure of 1.26 bar is already quite close to the applied 1 bar. Also, the temperature of the water, and of the simulation cell (it is a degree of freedom and has therefore an associated temperature) can also been seen to converge to the applied value of 300K. Energy conservation is adequate with a 0.001 ps time step.

```

Current cycle: 95000 out of 100000
=====
time run: 95.000000 [ps] 0.095000 [ns]
Current Box: 22.88107 0.00000 0.00000 [Å] Average Box: 22.94823 0.00000 0.00000 [Å]
          0.00000 22.88107 0.00000 [Å] 0.00000 22.94823 0.00000 [Å]
          0.00000 0.00000 22.88107 [Å] 0.00000 0.00000 22.94823 [Å]
Box-lengths: 22.88107 22.88107 22.88107 [Å] Average: 22.94823 22.94823 22.94823 [Å]
Box-angles: 90.00000 90.00000 90.00000 [degrees] Average: 90.00000 90.00000 90.00000 [degrees]
Volume: 11979.23948 [Å^3] Average Volume: 12085.81298 [Å^3]

Total linear momentum: -0.0000000000 [Å] -0.0000000000 [Å] -0.0000000000 [Å]

Current Stress: 125.84215 -65.75840 -32.61888 [MPa] Average Stress: -452.24145 -1504.12491 -1645.75996 [kPa]
          -1.68779 82.82297 -38.61796 [MPa] -1516.94287 633.97239 -278.59183 [kPa]
          32.19790 -105.05104 40.04903 [MPa] -1636.72478 -285.77225 197.27149 [kPa]
Average pressure: 126.3341475732 [kPa] 1.2468210962 [atm] 1.2633414757 [bar]

Amount of molecules per component:

```

```

-----
Component 0 (Tip5p), current number of mol/c: 400.00000 (average 400.00000), density: 998.11733 (average 989.50777) kg/m^3
-----
Number of Framework-atoms: 0
Number of Adsorbates: 400
Number of Cations: 0

Conserved energy: -691791.3447338404 Energy drifts: 0.0000617480 0.0001402288
Temperature: 302.061 (avg. 300.407), Translational (avg. 300.481), Rotational (avg. 300.333)
Temperature Adsorbates: 302.061 (avg. 300.407), Translational (avg. 300.481), Rotational (avg. 300.333)
Cell temperature: 265.890 (avg. 300.991)

```

## Example 4: Adsorption of CO<sub>2</sub> in Na-LTA

The Linde Type A structure LTA-4A has 96 aluminum per unit cell. A common 4A sample has 96 charge balancing sodium ions. The ions are small enough to access the sodalite cages, but the bigger methane molecules are exclusively in the big  $\alpha$ -cages and not in the sodalite cages. They need to be artificially blocked. Because the adsorption is dependent on the positions of the ions it is important to start from the crystallographic positions and use *only* translation for the ions. Reinsertion moves may transport the ions to positions in the windows and this is especially important for diffusion (the next example).

```

SimulationType MonteCarlo
NumberOfCycles 25000
NumberOfInitializationCycles 10000
RestartFile no
PrintEvery 1000

Forcefield ElenaSodium2009
ModifyOxgensConnectedToAluminium yes

Framework 0
FrameworkName LTASA
UnitCells 1 1 1
ExternalTemperature 298.0
ExternalPressure 10000.0

Component 0 MoleculeName sodium
MoleculeDefinition TraPPE
TranslationProbability 1.0
RandomTranslationProbability 1.0
ExtraframeworkMolecule yes
CreateNumberOfMolecules 96

Component 1 MoleculeName CO2
MoleculeDefinition TraPPE
BlockPockets yes
BlockPocketsFilename LTA
TranslationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
ExtraframeworkMolecule no
CreateNumberOfMolecules 0

```

## Example 5: Diffusion of CO<sub>2</sub> in Na-LTA

An example of molecular dynamics of an adsorbate (CO<sub>2</sub>) diffusing through the pores of LTA 4A loaded with ions. The ions are read from the restart-file. The mean-square displacement is computed during the run.

```

SimulationType MolecularDynamics
NumberOfCycles 250000
NumberOfInitializationCycles 5000
NumberOfEquilibrationCycles 10000
Restartfile no
PrintEvery 10000

Ensemble NVT

Forcefield ElenaSodium2009
ModifyOxgensConnectedToAluminium yes
TimeStep 0.0005

Framework 0
FrameworkName LTASA
UnitCells 1 1 1
ExternalTemperature 600.0
ComputeMSD yes
PrintMSDEvery 5000

component 0 MoleculeName sodium
MoleculeDefinition TraPPE
TranslationProbability 1.0
ReinsertionProbability 1.0
ExtraframeworkMolecule yes
CreateNumberOfMolecules 96

component 1 MoleculeName CO2
MoleculeDefinition TraPPE

```




Figure 4.3: Gibbs ensemble simulation of CO<sub>2</sub> at 250K. Two simulation boxes are used: one for the gas-branch and one for the liquid branch. The simulation can only be conducted below a certain temperature because otherwise the boxes can swap between gas and liquid. At 250K, the boxes are initialized with an equal amount of molecules, but soon split into gas and liquid. The average densities are straightforward to measure. As shown, the TraPPE model for CO<sub>2</sub> does a good job when compare to experimental data (NIST database).

```

BlockPockets      yes
BlockPocketsFilename LTA
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
ExtraFrameworkMolecule no
CreateNumberOfMolecules 64

```

### Example 6: Diffusion of benzene in rigid IRMOF-1

Benzene (and aromatic molecules in general) are usually kept rigid. RASPA uses quaternions for the description of the orientation of the molecules. The integration schemes of Martyna and Tuckermann are symplectic and conserve energy very well. Even though the molecule is described as a center of mass and a orientation, the forces are still computed atomically. In this example the diffusivity the mean-square displacement of benzene at 298K in IRMOF-1 is computed. The forcefield is 'FlexibleIRMOF-1' which is also perfectly suitable for rigid structures. It has been specifically optimized for iso-reticular metal-organic frameworks.

```

SimulationType      MolecularDynamics
NumberOfCycles     100000000
NumberOfEquilibrationCycles 10000
NumberOfInitializationCycles 1000
PrintEvery          1000
RestartFile        no

Ensemble           NVT

ChargeMethod       Ewald
CutOff             12.0
TimeStep            0.001
Forcefield         FlexibleIRMOF-1
EwaldPrecision    1e-6

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0

Component 0 MoleculeName      benzene
StartingBead        0
MoleculeDefinition  TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 16

```

## Example 7: Gibbs ensemble simulation of CO<sub>2</sub>

The Gibbs ensemble is way of computing coexistence without interfaces. It is one the most used methods to study vapor-liquid and liquid-liquid equilibria, it is not suitable for very dense systems. The conditions for coexistence of two or more phases I, II, ... is that the pressure and temperature of all the phases must be equal, as well as the chemical potential of all the species. The Gibbs ensemble example for the single component CO<sub>2</sub> is listed below. two boxes will be used, one will correspond to the liquid phase, the other one to the gas phase. The 'GibbsVolumeChange' move changes the individual volume leaving the total volume in tact, the 'GibbsSwap' move swaps particles from one box to the other. One of the practical problems is to make sure both boxes remain larger than twice the cutoff length. If not, the program will exit with an error message, and the simulation should be restarted with a bigger volume. Note that RASPA uses orientational biased insertions for small rigid molecules like CO<sub>2</sub>. For this example about 10000-20000 cycles are needed to equilibrate properly.

```

SimulationType MonteCarlo
NumberOfCycles 25000
NumberOfInitializationCycles 20000
RestartFile no
PrintEvery 100

Forcefield TraPPE

Box 0
BoxLengths 30 30 30
BoxAngles 90 90 90
ExternalTemperature 250.0

Box 1
BoxLengths 30 30 30
BoxAngles 90 90 90
ExternalTemperature 250.0

GibbsVolumeChangeProbability 1.0

Component 0 MoleculeName CO2
StartingBead 1
MoleculeDefinitions TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
GibbsSwapProbability 1.0
CreateNumberOfMolecules 150 150

```

## Example 8: Minimization of a flexible framework (fixed volume)

Physically, energy minimization corresponds to an instantaneous freezing of the system; a static structure in which no atom feels a net force corresponds to a temperature of 0 K. In the early 1980's, energy minimization was about all one could afford to do and was dubbed 'molecular mechanics.' Here, a difficult optimization problem: a flexible framework, IRMOF-1, in a periodic unit cell, with many low energy modes. The energy landscape of a framework is very complex. A true minimum is characterized by all positive eigenvalues of the Hessian matrix (the matrix of second derivatives with respect to position). A zero eigenvalue means that moving in the direction of the associated eigenvector does not result in a change in energy. Likewise, a negative en positive eigenvalue means an decrease and increase in energy, respectively. Most of the optimization time is spent on reaching a zero curvature structure, i.e. all positive eigenvalues.

```

SimulationType Minimization
NumberOfCycles 1
Restartfile no

MaximumNumberOfMinimizationSteps 1000
RMSGradientTolerance 1e-8
MaxGradientTolerance 1e-8

Ensemble NVT

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff 12.0
Forcefield FlexibleIRMOF-1-new

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0
Movies yes

FlexibleFramework yes
FrameworkDefinitions FlexibleIRMOF-1-new

```

The minimization needs 119 cycles to optimize IRMOF-1, the last steps are shown here. The convergence is very rapid (quadratic) near the minimum, and the minimum energy can be reached up to arbitrary

precision (the forces on all the atoms are  $1 \times 10^{-8}$  K/Å<sup>2</sup> or smaller). To compute spectra, frequencies and/or eigenmodes a high precision is needed.

```

Starting configuration:
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: 129316.6369463501      0.0000000016      0.0000000015
        0.0000000000      25.8320000000      0.0000000000      0.0000000015      129316.6369463478      0.0000000016
        0.0000000000      0.0000000000      25.8320000000      0.0000000016      0.0000000015      129316.6369463608

Beginning Baker minimization:
-----
Shifting parameter: -144554.0050033366 Lowest eigenvalue: -1933.2972944159
Iteration: 0 Energy: -4210346.0956447003 RMS gradient: 254.796 Max gradient: 16621.7 Number of negative eigenvalues: 30 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: 129316.6369463581      0.0000000016      0.0000000015
        0.0000000000      25.8320000000      0.0000000000      0.0000000016      129316.6369463568      0.0000000016
        0.0000000000      0.0000000000      25.8320000000      0.0000000015      0.0000000016      129316.6369463690
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000

Shifting parameter: -54112.0433245863 Lowest eigenvalue: -942.0341042027
Iteration: 1 Energy: -4291359.2884553447 RMS gradient: 132.811 Max gradient: 9119.05 Number of negative eigenvalues: 30 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: -131739.2152028512      0.0000000004      0.0000000008
        0.0000000000      25.8320000000      0.0000000000      0.0000000004      -131739.2152028439      0.0000000010
        0.0000000000      0.0000000000      25.8320000000      0.0000000008      0.0000000011      -131739.2152028617
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000

Shifting parameter: -6990.6349276451 Lowest eigenvalue: -218.9092550725
Iteration: 2 Energy: -4325503.0430874182 RMS gradient: 38.7491 Max gradient: 2721.14 Number of negative eigenvalues: 36 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: -282155.2863533634      0.0000000002      0.0000000007
        0.0000000000      25.8320000000      0.0000000000      0.0000000002      -282155.2863533496      -0.0000000021
        0.0000000000      0.0000000000      25.8320000000      0.0000000008      -0.0000000021      -282155.2863533603
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000
.....
Iteration: 117 Energy: -4331364.7259515934 RMS gradient: 0.00439145 Max gradient: 0.600384 Number of negative eigenvalues: 0 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: -99169.3483007069      0.0011594244      0.0002624505
        0.0000000000      25.8320000000      0.0000000000      0.0011594245      -99108.1188463482      -0.0004447560
        0.0000000000      0.0000000000      25.8320000000      0.0002624506      -0.0004447560      -99128.0909746577
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000

Iteration: 118 Energy: -4331364.6575047411 RMS gradient: 7.06135e-05 Max gradient: 0.0204619 Number of negative eigenvalues: 0 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: -99208.8822655176      0.0000019593      -0.0000031410
        0.0000000000      25.8320000000      0.0000000000      0.0000019594      -99184.4522480669      -0.0000044895
        0.0000000000      0.0000000000      25.8320000000      -0.0000031410      -0.0000004895      -99202.7818375186
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000

Iteration: 119 Energy: -4331364.6575191850 RMS gradient: 6.91407e-07 Max gradient: 8.83309e-05 Number of negative eigenvalues: 0 Number of zero eigenvalues: 3
Box:    25.8320000000      0.0000000000      0.0000000000      Strain derivative: -99195.5973070737      0.0000000048      -0.0000030267
        0.0000000000      25.8320000000      0.0000000000      0.0000000047      -99195.5991107764      0.0000000006
        0.0000000000      0.0000000000      25.8320000000      -0.0000030267      0.0000000006      -99195.6111130547
Lengths: 25.8320000000      25.8320000000      25.8320000000, Angles: 90.0000000000      90.0000000000      90.0000000000

SUCSES: RMS Gradient tolerance 0.0001 reached (6.91407e-07)
Max Gradient tolerance 0.0001 reached (8.83309e-05)
```

The shifting values are always lower than the lowest eigenvalues, both are negative and approach zero. At iteration 2, the lowest eigenvalues is closer to zero, but still the amount of negative eigenvalues is 6 higher. Also increases in energy can occur. However, eventually the system is driven to all positive eigenvalues (a true energy minimum without saddle points) and the lowest energy. Note that minimization the structure in constant volume results in a finite (non-zero) stress. Minimization taking volume and shape changes into account are usually easier, because the system is less constrained. If one would like to also minimize the cell volume (isotropically) use

Ensemble                          NPT

or use for a change in cell-lengths and cell-angles

Ensemble                          NPTPR

### Example 9: Computing sitings and occupancies of cations in LTA4A

RASPA is capable of computing information on sitings of ions and occupancies. For LTA5A, three sites are experimentally known:

1. Site I is located in the center of the six ring, there are 64 such sites,
2. Site II is located in the eight ring window, with 96 sites,
3. Site III is located opposite to a four ring, with 96 sites.

The force field of Calero et al. is able to provide a very good description of sodium-loaded aluminosilicates. With 'ComputeCationAndAdsorptionSites yes' we say that the adsorption sites and occupancies

should be computed. The information on the sites is read in using 'IonsName LTA' (located in 'structures/zeolites/ions/') and generating the full sitings from the asymmetric data using 'SpaceGroupIons 226' and 'AsymmetricIons yes'. Every ion located more than 3 Å ('CutOffIons 2.0') is regarded as 'unclassified'.

```

SimulationType MonteCarlo
NumberOfCycles 50000
NumberOfInitializationCycles 50000
PrintEvery 100
RestartFile no

ChargeMethod Ewald
CutOff 12.0
Forcefield ElenaSodiumCalcium
EwaldPrecision 1e-6

Framework 0
FrameworkName LTA5A
SpaceGroup 226
Asymmetric yes
UnitCells 1 1 1
CutOffIons 2.0
IonsName LTA
SpaceGroupIons 226
AsymmetricIons yes
ComputeCationAndAdsorptionSites yes
ExternalTemperature 30.0

Component 0 MoleculeName sodium
MoleculeDefinition TraPPE
TranslationProbability 1.0
RandomTranslationProbability 1.0
ExtraFrameworkMolecule yes
CreateNumberOfMolecules 96

```

RASPA creates a directory 'CationAndAdsorptionSites/System[int]' containing the file 'crystallographic.data'. The 'relative occupancy' is the relative probability of finding an ion in such a site. The 'occupancy' can be compared to experiment; it is the amount of ions divided by the amount of sites of that type. For example, 24 ions on average occupy site II, but there are 96 such sites. The occupancy is therefore 0.25.

```

Crystallographic data
=====
Average unclassified: 0.998400

Cation Position 0
-----
Relative Occupation 0.687517 Ions 66.002 Occupation 1.031275
[ 0.099600 0.099600 0.099600] -> [ 0.246348 0.102809 0.101850]
distances [x] 0.563504 [y] 0.556447 [z] 0.560021 [xyz] 0.969945 0.970940 count: 1650106.000000
B-factor:
0.335536 -0.004404 0.324743
-0.004404 0.327723 -0.021652
0.324743 -0.021652 0.325674

Cation Position 1
-----
Relative Occupation 0.250000 Ions 24.000 Occupation 0.250000
[ 0.216500 0.211100 0.000000] -> [ 0.236804 0.221718 0.006002]
distances [x] 0.203540 [y] 0.212305 [z] 0.210883 [xyz] 0.361903 0.387284 count: 600024.000000
B-factor:
0.042283 -0.004565 0.053705
-0.004565 0.039462 0.000200
0.053705 0.000200 0.050990

Cation Position 2
-----
Relative Occupation 0.052083 Ions 5.000 Occupation 0.052083
[ 0.250000 0.106000 0.106000] -> [ 0.250351 0.114502 0.110723]
distances [x] 0.133796 [y] 0.092626 [z] 0.147901 [xyz] 0.219899 0.251636 count: 125005.000000
B-factor:
0.025677 -0.001288 0.012997
-0.001288 0.021022 0.011627
0.012997 0.011627 0.024813

```

Crystallographic studies indicate that these sites are occupied for 97.2, 24.2, and 6.6%, respectively. Molecular simulations with our optimized force field yield occupations of 100, 25.0 and 5.21%, respectively. Less than 1 ion is unclassified.

## Example 10: Computing sitings and occupancies of argon in IRMOF-1

The siting and occupancy analysis can also be performed on regular molecules. Here, both the framework and ions are generated from the asymmetric unit cell using spacegroup 225. These data are from an experimental paper describing both the positions of the framework atoms and of the adsorbing argon (and nitrogen). Dubbeldam et al. used molecular simulation to investigate the sitings computationally.

```

SimulationType MonteCarlo
NumberOfCycles 100000
NumberOfInitializationCycles 50000
PrintEvery 100
RestartFile no

ChargeMethod Ewald
CutOff 12.0
Forcefield FlexibleIRMOF-1
EwaldPrecision 1e-6

Framework 0
FrameworkName IRMOF-1-Ar
SpaceGroup 225
Asymmetric yes
UnitCells 1 1 1
CutOffIons 1.0
IonsName IRMOF-1-Ar
SpaceGroupIons 225
AsymmetricIons yes
ComputeCationAndAdsorptionSites yes
ExternalTemperature 30.0
ExternalPressure 1e4

Component 0 MoleculeName argon
MoleculeDefinition TraPPE
TranslationProbability 1.0
ReinsertionProbability 1.0
ExtraFrameworkMolecule yes
CreateNumberOfMolecules 32

```

For this example of argon in IRMOF-1 at 30K, more than 31 of the 32 molecules are always found in the first site, and only 0.25 molecules are unclassified.

```

Average unclassified: 0.245240
Cation Position 0
-----
Relative Occupation 0.977027 Ions 31.265 Occupation 0.977027
[ 0.341910 0.158090 0.158090] -> [ 0.341643 0.152717 0.148374]
distances [x] 0.155279 [y] 0.154603 [z] 0.155660 [xyz] 0.268782 0.308203 count: 2970192.000000

```

## 4.4 Advanced examples

### Example 1: Adsorption of N<sub>2</sub> in "catenated"-flexible IRMOF-10

Many metal-organic framework come in a catenated form, either interwoven or interpenetrated. Here, we allow the frameworks to move independently of each other. The first of the frameworks is kept fixed, the others are moving. Note that the internal structure of the framework remains unchanged. The whole system is first defined using 'Framework 0', and next two frameworks are defined using 'FrameworkName IRMOF-10'. The first is placed, but the second is shifted with respect to the first one 'ShiftUnitCells [real] [real] [real]' defined in fractional coordinates. It is very useful to compute the histogram of the distance between the centers of mass of the frameworks ('ComputeFrameworkSpacingHistogram yes'). The Monte-Carlo move to displace the framework is 'FrameworkShiftMoveProbability [real]' where [real] is the probability of the move in each *cycle*. Note that Van der Waals interactions and electrostatics between the frameworks are computed but not within each framework.

```

SimulationType MonteCarlo
NumberOfCycles 1000
NumberOfInitializationCycles 1000
PrintEvery 100
RestartFile no

ChargeMethod Ewald
CutOff 12.0
Forcefield FlexibleIRMOF-10
EwaldPrecision 1e-6

Framework 0
FrameworkName IRMOF-10
ShiftUnitCells 0.0 0.0 0.0
FrameworkName IRMOF-10
ShiftUnitCells 0.125 0.125 0.125
UnitCells 1 1 1
FrameworkDefinitions FlexibleIRMOF-10
ComputeFrameworkSpacingHistogram yes
ExternalTemperature 298.0
ExternalPressure 1e4 1e5 1e6 1e7

FlexibleFramework yes
FrameworkShiftMoveProbability 1.0

Component 0 MoleculeName N2

```

```

StartingBead          0
MoleculeDefinition    TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability   1.0
ReinsertionProbability 1.0
SwapProbability        1.0
CreateNumberOfMolecules 0

```

## Example 2: Adsorption of CO<sub>2</sub> in fully-flexible IRMOF-10 with water and DMF solvents presents

A common problem of MOFs is activation. Usually, even after activation, some solvents molecules are still present in the structure. Moreover, during the adsorption experiments it is always possible for water to co-absorb. Here an example to evaluate these solvent effects on the adsorption of CO<sub>2</sub> in IRMOF-10. The amount of water and dimethylformamide (DMF) is constant (no 'swap' move defined), but is moving in the background while CO<sub>2</sub> is swapped in and out.

```

SimulationType      MonteCarlo
NumberofCycles     25000
NumberofInitializationCycles 5000
Restartfile        no
PrintEvery         100

Forcefield          FlexibleIRMOF-16
CutOff             12.0
EwaldPrecision     1e-6

Framework 0
FrameworkName IRMOF-16
ShiftUnitCells 0.0 0.0 0.0
FrameworkName IRMOF-16
ShiftUnitCells 0.16666 0.16666 0.16666
FrameworkDefinitions FlexibleIRMOF-16
ComputeFrameworkSpacingHistogram yes
ExternalTemperature 298.0
ExternalPressure 0.5e5 1e5 2e5 3e5 4e5 5e5 7.5e5 10e5 12e5 14e5 16e5 18e5 20e5

FlexibleFramework yes
FrameworkShiftMoveProbability 1.0

Component 0 MoleculeName      DMF_CS2
           MoleculeDefinition  Rigid
           TranslationProbability 0.5
           RotationProbability   0.5
           ReinsertionProbability 0.25
           ExtraframeworkMolecule yes
           CreateNumberOfMolecules 10

Component 1 MoleculeName      Tip5p
           MoleculeDefinition  Water
           TranslationProbability 0.5
           RotationProbability   0.5
           ReinsertionProbability 0.25
           ExtraframeworkMolecule yes
           CreateNumberOfMolecules 40

Component 2 MoleculeName      CO2
           MoleculeDefinition  TraPPE
           TranslationProbability 0.5
           RotationProbability   0.5
           ReinsertionProbability 0.25
           SwapProbability       1.0
           CreateNumberOfMolecules 0

```

## Example 3: Adsorption of CO<sub>2</sub> in fully-flexible IRMOF-1 ( $\mu$ VT-ensemble)

Flexibility in MOFs is more important than in zeolites. A very efficient move to change the whole framework (and actually also the adsorbates) is have a short NVE MD-run and accept or reject the new configuration. This hybrid MD/MC move can be switched on using 'HybridMCMDMoveProbability [real]', where [real] is the fraction of the move at each cycle.

```

SimulationType      MonteCarlo
NumberofCycles     50000
NumberofInitializationCycles 10000
PrintEvery         100
Restartfile        no

ChargeMethod       Ewald
CutOff            12.0
Forcefield         FlexibleIRMOF-1
EwaldPrecision    1e-6

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1

```

```

HeliumVoidFraction 0.82
FrameworkDefinitions FlexibleIRMOF-1
ExternalTemperature 298.0
ExternalPressure 1e5 1e6 1e7 1e8 1e9 1e10

FlexibleFramework yes

HybridMCMDMoveProbability 1.0

Component 0 MoleculeName CO2
StartingBead 0
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
CreateNumberOfMolecules 0

```

## Example 4: NPT molecular dynamics of flexible IRMOF-1

An NPT-ensemble simulation of a flexible framework IRMOF-1. This type of simulation can be used to compute the average unit cell size at the desired temperature and pressure (and properties like the 'volumetric expansion coefficient' etc). The equilibration, although slow, is very much faster than Monte Carlo. The example show the code for flexible IRMOF-1 at 298K and 1 atm.

```

SimulationType MolecularDynamics
NumberOfCycles 500000
NumberOfEquilibrationCycles 5000
PrintEvery 100
RestartFile yes

Ensemble NPT

Forcefield FlexibleIRMOF-1
CutOff 12.0

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0
ExternalPressure 101325.0

FlexibleFramework yes
FrameworkDefinitions FlexibleIRMOF-1

```

## Example 5: CO<sub>2</sub> adsorption in flexible IRMOF-1 (osmotic ensemble), version I.

Absorption simulations using a flexible framework are very computationally demanding, the current example will probably run about a week. The equilibration is very important and it is best to start with a restart-file obtained from the previous example at the same temperature. The directory 'Restart' produced in the previous example should be copied to 'RestartInitial' and the option 'RestartFile' should be set to 'yes'.

```

SimulationType MonteCarlo
NumberOfCycles 50000
NumberOfInitializationCycles 10000
PrintEvery 100
RestartFile no

ChargeMethod Ewald
CutOff 12.0
Forcefield FlexibleIRMOF-1
EwaldPrecision 1e-6
TimeStep 0.0005

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
HeliumVoidFraction 0.82
ExternalTemperature 298.0
ExternalPressure 1e5 1e6 1e7 1e8 1e9 1e10

FrameworkDefinitions FlexibleIRMOF-1
FlexibleFramework yes

HybridNVEMoveProbability 1.0
NumberOfHybridNVESTeps 5
VolumeChangeProbability 1.0

Component 0 MoleculeName CO2
StartingBead 0
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
CreateNumberOfMolecules 0

```

## Example 6: CO<sub>2</sub> adsorption in flexible IRMOF-1 (osmotic ensemble), version II.

An alternative to using separate MC/MD hybrid and volume moves, is to combine the two into the NPH MC/MD hybrid move. Here, and MD path is computed in the NPH ensemble, and the generated configuration is then accepted or rejected. This move is potentially more efficient because all of the particle have adjusted more gently to the volume changes during the MD path.

```
SimulationType MonteCarlo
NumberOfCycles 50000
NumberOfInitializationCycles 10000
PrintEvery 100
RestartFile no

ChargeMethod Ewald
CutOff 12.0
Forcefield FlexibleIRMOF-1
EwaldPrecision 1e-6
TimeStep 0.0005

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
HeliumVoidFraction 0.82
ExternalTemperature 298.0
ExternalPressure 1e5 1e6 1e7 1e8 1e9 1e10

FrameworkDefinitions FlexibleIRMOF-1
FlexibleFramework yes

HybridNPHMoveProbability 1.0
NumberOfHybridNPHSteps 5

Component 0 MoleculeName CO2
StartingBead 0
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
CreateNumberOfMolecules 0
```

or alternatively, using the NPHPR ensemble with flexible angles,

```
NPTPRCellType RegularUpperTriangle
HybridNPHPRMoveProbability 1.0
NumberOfHybridNPHPRSteps 5
```

or monoclinic

```
NPTPRCellType MonoclinicUpperTriangle
HybridNPHPRMoveProbability 1.0
NumberOfHybridNPHPRSteps 5
```

or anisotropic

```
NPTPRCellType Anisotropic
HybridNPHPRMoveProbability 1.0
NumberOfHybridNPHPRSteps 5
```

## Example 7: methane adsorption in flexible MFI (Demontis model, no VDW/Ewald)

One of the early models for siliceous zeolites was developed by Demontis et al. The model omits the long-range Van der Waals and electrostatic interactions, and uses simple harmonic potentials for stretching and bending. However, as a simple and fast conceptual model it is has proven very powerful to study flexibility.

```
SimulationType MonteCarlo
NumberOfCycles 10000
NumberOfInitializationCycles 1000
PrintEvery 100
RestartFile no

ChargeMethod None
InternalFrameworkLennardJonesInteractions no
CutOff 12.0
DeltaT 0.002
Forcefield ElenaSodiumCalcium
EwaldPrecision 1e-6

Framework 0
FrameworkName MFI
UnitCells 2 2 2
HeliumVoidFraction 1.0
ExternalTemperature 195.0
ExternalPressure 1e4 1e5 1e6 1e7 1e8 1e9
```

```

FlexibleFramework yes
FrameworkDefinitions DemontisModified

HybridNVEMoveProbability 1.0

Component 0 MoleculeName      argon
          StartingBead        0
          MoleculeDefinition   TraPPE
          IdealGasRosenbluthWeight 1.0
          TranslationProbability 1.0
          RotationProbability    1.0
          ReinsertionProbability  1.0
          SwapProbability        1.0
          CreateNumberOfMolecules 0

```

### Example 8: Argon adsorption in flexible MFI (full Nicholas model, cell is allowed to change shape)

The Nicholas model for siliceous zeolites is more accurate than the Demontis model, but also much more expensive to compute. It includes stretching, bending, torsion, Van der Waals and electrostatic interactions. The model is not developed for NPT simulations. However, in this example we do allow the shape of the box to change upon adsorption.

```

SimulationType      MonteCarlo
NumberofCycles     1000
NumberofInitializationCycles 1000
PrintEvery         100
RestartFile        no

ChargeMethod       Ewald
CutOff            12.0
Forcefield         ElenaSodiumCalcium
EwaldPrecision    1e-6

Framework 0
FrameworkName MFI
UnitCells 2 2 2
HeliumVoidFraction 1.0
ExternalTemperature 195.0
ExternalPressure 1e4 1e5 1e6 1e7 1e8 1e9

FlexibleFramework yes
FrameworkDefinitions NicholasModified

HybridMCMDMoveProbability 1.0
BoxShapeChangeProbability 1.0

Component 0 MoleculeName      argon
          StartingBead        0
          MoleculeDefinition   TraPPE
          IdealGasRosenbluthWeight 1.0
          TranslationProbability 1.0
          RotationProbability    1.0
          ReinsertionProbability  1.0
          SwapProbability        1.0
          CreateNumberOfMolecules 0

```

### Example 9: Benzene diffusion in flexible IRMOF-10

Molecules with a phenyl-ring are usually quite rigid. In Monte Carlo rigid units are not a problem, because the MC moves can be developed in such a way that the constraints remain satisfied, i.e. translation and rotation of the whole rigid unit. In molecular dynamics, there are two general approaches. The first is to integrate the molecules atomically and afterwards satisfy the constraints iteratively using for example the shake algorithm. For bigger molecules complications arise, convergence becomes more difficult, and for a planar molecule like benzene additional sites above the molecule are needed. Therefore, the second approach has become more popular. Using quaternions (or Euler angles) one can describe the configurations of the molecule as a center-of-mass position and an orientation. The translation and rotation are integrated and when the forces are needed the atoms positions are computed from the com position and the orientation. The forces are then summed to the center of mass and the torque is computed. Miller et al. have developed an integration algorithm for rigid units (using quaternions) that is symplectic.

All these techniques are combined in the example of diffusion of benzene in IRMOF-10. The integration is performed in the NVT ensemble using the Nose-Hoover thermostats. Three separate NH chains are operating on (i) the translation, (ii) the rotation of the molecules, and (iii) on the framework.

```

SimulationType      MolecularDynamics
NumberofCycles     100000000
NumberofEquilibrationCycles 10000

```

```

NumberOfInitializationCycles 1000
PrintEvery 1000
RestartFile no

Ensemble NVT

ChargeMethod Ewald
CutOff 12.0
TimeStep 0.0005
Forcefield FlexibleIRMOF-10
EwaldPrecision 1e-6

Framework 0
FrameworkName IRMOF-10
UnitCells 1 1 1
ExternalTemperature 298.0

FrameworkDefinitions FlexibleIRMOF-10
FlexibleFramework yes

Component 0 MoleculeName benzene
StartingBead 0
MoleculeDefinition TraPPE
IdealGasRosenbluthWeight 1.0
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
CreateNumberOfMolecules 16

```

## Example 10: Aromatics in Na-Y

The Fuchs group has developed models for aromatics in Na-Y. The xylenes are modeled as rigid. The zeolite Y is taken from the crystallographic position (Fitch structure), and position I and II are filled (48 sodium). The ions are integrated into the framework (framework is charge neutral). The model include the explicit computation of polarization. However, since the aromatics do not polarize each other and the framework is assumed to be rigid, the computation can be immediately terminated after 1 iteration. That is, the polarization energy is directly related the static electric field vector and there is no induction because this electric field does not generate new induced dipoles.

```

SimulationType MonteCarlo
NumberOfCycles 100000
NumberOfInitializationCycles 25000
PrintEvery 1000
Restartfile no

Forcefield AromaticsFuchs2001
ComputePolarization yes
BackPolarization no
OmitAdsorbateAdsorbatePolarization yes
ChargeMethod Ewald
CutOff 12.0

Framework 0
FrameworkName FAU_Fitch
Asymmetric yes
Spacegroup 478
InputFileType cssr
UnitCells 1 1 1
ExternalTemperature 423
ExternalPressure 200

Component 0 MoleculeName m-xylene
StartingBead 0
MoleculeDefinition AromaticsFuchs
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
CreateNumberOfMolecules 0

```

## Example 11: Computing the infra-red spectrum using MD

Spectra reported in the literature using molecular dynamics are usually the Fourier transform of the velocity autocorrelation function (VACF) or the Fourier transform of the charge weighted VACF. RASPA computes the spectra on-the-fly during the simulation, when the option 'ComputeSpectra' is 'yes'. Every 'PrintSpectraEvery' MD-steps the spectra are written. The spectra are written to the directory 'MD\_Spectra'. Here, an example of siliceous LTA.

```

SimulationType MolecularDynamics
NumberOfCycles 250000
NumberOfEquilibrationCycles 10000
PrintEvery 100
RestartFile no

Ensemble NVT

```

```

Forcefield           ElenaSodiumCalcium
TimeStep             0.0005

Framework 0
FrameworkName LTA_SI
UnitCells 1 1 1
ExternalTemperature 600.0
ComputeInfraredSpectra yes
WriteInfraredSpectraEvery 5000

FlexibleFramework yes
FrameworkDefinitions NicholasModified

```

## Example 12: Minimization of monoclinic flexible Hupp-1 (NPT Parinello-Rahman)

The catenated version of Hupp-1 is monoclinic. Because of the minimum-image convention more than  $1 \times 1 \times 1$  unit cells are required. For minimization, RASPA uses imaginary replicas of the unit cell for the computation of the long-range interactions. The amount of replicas is specified with the 'ReplicaUnitCells' keyword. Since the minimization can be done using 1 single unit cell, the computation of eigenvalues/vectors in the mode following technique becomes feasible. Minimization using this technique is fast, i.e. it requires much less iterations than other techniques, is much more accurate, and leads to a true minimum (all positive eigenvalues of the Hessian matrix). In this example, the minimization also includes the cell matrix, but restricted to be remain monoclinic.

```

SimulationType Minimization
NumberOfCycles 1
PrintEvery 10
RestartFile no

Ensemble NPTPR
NPTPRCellType MonoclinicUpperTriangle

RemoveBondNeighboursFromLongRangeInteraction no
RemoveBendNeighboursFromLongRangeInteraction no
RemoveTorsionNeighboursFromLongRangeInteraction no

Remove12NeighboursFromChargeChargeInteraction yes
Remove13NeighboursFromChargeChargeInteraction yes
Remove14NeighboursFromChargeChargeInteraction no

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff 12.0
Forcefield FlexibleHupp-1
EwaldPrecision 1e-8

Framework 0
FrameworkName Hupp-1-mono
Asymmetric yes
SpaceGroup 12
UnitCells 1 1 1
ReplicaUnitCells 3 3 2
ExternalTemperature 298.0
ExternalPressure 1e5
FlexibleFramework yes
FrameworkDefinitions FlexibleHupp-1
Movies yes

```

## Example 13: Minimization of triclinic flexible MnSalen or ZnSalen

The Mn- and Zn-salen are triclinic ( $\alpha$ ,  $\beta$ , and  $\gamma$  angles differ from  $90^\circ$ ). Replicas are used to minimize a single unit cell in NPT-PR. The resulting structure is strain/stress-free. The type of the NPT-PR is 'RegularUpperTriangle' which means the 6 independent strain-elements are included in the optimization. After minimization, the cell is converted to upper-triangular form.

```

SimulationType Minimization
NumberOfCycles 1
PrintEvery 10
RestartFile no

Ensemble NPTPR
NPTPRCellType RegularUpperTriangle

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
TimeStep 0.0005
CutOff 12.0
Forcefield MnSalenMof
EwaldPrecision 1e-8

Remove12NeighboursFromChargeChargeInteraction yes
Remove13NeighboursFromChargeChargeInteraction yes
Remove14NeighboursFromChargeChargeInteraction no

```

```

Framework 0
FrameworkName MnSalenMof
Asymmetric no
SpaceGroup 1
UnitCells 1 1 1
ReplicaUnitCells 3 2 1
ExternalTemperature 298.0
ExternalPressure 0
FlexibleFramework yes
FrameworkDefinitions MnSalenMof
Movies yes

```

### Example 14: Elastic constants of IRMOF-1 (0 Kelvin)

For a minimized structure, one can compute elastic constants at zero Kelvin. Use for example Baker's minimization to obtain a true energy minimum (with *all* positive eigenvalues of the Hessian matrix) and forces smaller than  $10^{-8}$  K/Å. At these conditions, the elastic constants can be computed from the generalized Hessian matrix, which includes the Born-term (second derivative of the energy with respect to strain), and cross terms (derivative of the energy with respect to strain and position). The constants can be computed analytically and therefore with a very high degree of accuracy.

Because a restart-file is written as ascii with insufficient precision, the elastic constants are computed after a minimization. The keyword 'ComputeElasticConstants yes' is added to compute the elastic constants. Starting from a restart file with a converged minimization it usually take 1-3 steps to satisfy the convergence criteria again (mainly for the stress terms).

```

SimulationType Minimization
NumberofCycles 1
PrintEvery 10
RestartFile no

Ensemble NPTPR
NPTPRCellType RegularUpperTriangle

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff 12.0
Forcefield FlexibleIRMOF-1-new
EwaldPrecision 1e-6

ComputeElasticConstants yes

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0
ExternalPressure 0.0

FlexibleFramework yes
FrameworkDefinitions FlexibleIRMOF-1-new

```

The output prints the total elastic constants as well as the individual terms like the relaxation term. In MOFs, the relaxation term is surprisingly large.

Elastic constant [GPa]									
29.30175	0.00000	-0.00000	0.00000	0.98456	0.00000	-0.00000	0.00000	0.98456	
0.00000	11.91216	0.00000	0.98456	-0.00000	-0.00000	0.00000	0.00000	0.00000	0.00000
-0.00000	0.00000	11.91216	-0.00000	-0.00000	0.00000	0.98456	0.00000	-0.00000	
0.00000	0.98456	0.00000	11.91216	-0.00000	0.00000	0.00000	-0.00000	0.00000	0.00000
0.98456	-0.00000	-0.00000	-0.00000	29.30175	-0.00000	0.00000	-0.00000	0.98456	
0.00000	-0.00000	0.00000	0.00000	-0.00000	11.91216	0.00000	0.98456	0.00000	0.00000
-0.00000	-0.00000	0.98456	0.00000	-0.00000	0.00000	11.91216	0.00000	-0.00000	
0.00000	0.00000	0.00000	-0.00000	-0.00000	0.98456	0.00000	11.91216	0.00000	0.00000
0.98456	0.00000	-0.00000	0.00000	0.98456	0.00000	-0.00000	0.00000	29.30175	

Born-term [GPa]									
71.87089	-0.00000	-0.00000	-0.00000	27.81220	-0.00000	-0.00000	-0.00000	27.81220	
-0.00000	20.91401	-0.00000	27.81220	0.00000	0.00000	-0.00000	0.00000	0.00000	0.00000
-0.00000	-0.00000	20.91401	-0.00000	0.00000	0.00000	27.81220	0.00000	0.00000	
-0.00000	27.81220	-0.00000	20.91401	0.00000	0.00000	-0.00000	0.00000	0.00000	0.00000
27.81220	0.00000	0.00000	71.87089	0.00000	0.00000	0.00000	0.00000	27.81220	
-0.00000	0.00000	0.00000	0.00000	20.91401	0.00000	27.81220	0.00000	0.00000	0.00000
-0.00000	-0.00000	27.81220	-0.00000	0.00000	0.00000	20.91401	0.00000	0.00000	0.00000
-0.00000	0.00000	0.00000	0.00000	27.81220	0.00000	20.91401	0.00000	20.91401	0.00000
27.81220	0.00000	0.00000	27.81220	0.00000	0.00000	0.00000	0.00000	71.87089	

Relaxation-term [GPa]									
42.56914	-0.00000	0.00000	-0.00000	26.82765	-0.00000	0.00000	-0.00000	26.82765	
-0.00000	9.00184	-0.00000	26.82765	0.00000	0.00000	-0.00000	-0.00000	-0.00000	
0.00000	-0.00000	9.00184	0.00000	0.00000	-0.00000	26.82765	-0.00000	0.00000	
-0.00000	26.82765	-0.00000	9.00184	0.00000	-0.00000	0.00000	0.00000	-0.00000	
26.82765	0.00000	0.00000	42.56914	0.00000	0.00000	0.00000	0.00000	26.82765	
-0.00000	0.00000	-0.00000	0.00000	9.00184	-0.00000	26.82765	-0.00000	-0.00000	
0.00000	0.00000	26.82765	-0.00000	0.00000	-0.00000	9.00184	-0.00000	0.00000	
-0.00000	-0.00000	-0.00000	0.00000	26.82765	-0.00000	9.00184	-0.00000	-0.00000	
26.82765	-0.00000	0.00000	-0.00000	26.82765	-0.00000	0.00000	-0.00000	42.56914	

A cubic structure has three elastic constants,  $C_{11} = 29.3$  GPa,  $C_{12} = 11.91$  GPa, and  $C_{44} = 0.98$  GPa.

### Example 15: Elastic constants (finite temperature)

At finite temperature the elastic constants can be measured in several ways. Recently, computing the constants from stress-stress correlations has gained popularity, mainly due to its faster convergence. The keyword 'ComputeElasticConstants yes' switches on the computation of the elastic constants. Instead of the usual energy/force routines, it uses routines that also compute the Born-term (second derivative of the energy with respect to strain). Because this computation is very expensive it is convenient to be able to restart it to continue where it left off using the keyword 'WriteBinaryRestartFileEvery 25000'. Restart the simulation from the file written in 'CrashRestart' using 'ContinueAfterCrash yes'.

```

SimulationType      MolecularDynamics
NumberOfCycles     25000000
NumberOfEquilibrationCycles 500000
PrintEvery          5000
RestartFile         no

Ensemble NVT
ComputeElasticConstants yes

ContinueAfterCrash no
WriteBinaryRestartFileEvery 25000

ChargeMethod           Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff                12.0
TimeStep               0.0005
Forcefield             FlexibleIRMOF-1-new
EwaldPrecision         1e-6

Framework 0
FrameworkName IRMOF-1-300K
UnitCells 1 1
FrameworkDefinitions FlexibleIRMOF-1-new
FlexibleFramework yes
ExternalTemperature 300.0
ExternalPressure 0.0

```

### Example 16: Normal mode analysis (frequencies and movies of the modes)

After (true) minimization, it is instructive to view the eigenmodes of the system and analyze the frequencies with which they move. The low frequency modes often involve movement of a large number of atoms. These eigenmodes and eigenvalues represent a new basis, and provide the same information as the Cartesian representation, but physically more intuitive.

```

SimulationType Spectra
PrintEvery      10
RestartFile     yes

Ensemble       NVT

ComputerNormalModes yes
MinimumMode    0
MaximumMode   25
ModeResolution 200

ChargeMethod           Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff                12.0
Forcefield             FlexibleIRMOF-1-new

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1
ExternalTemperature 298.0

FlexibleFramework yes
FrameworkDefinitions FlexibleIRMOF-1-new

```

### Example 17: NPT Parinello-Rahman molecular dynamics (flexible box shape)

An example of molecular dynamics in expended ensembles, here NPT-PR where the box is allowed to change size and shape.

```

SimulationType      MolecularDynamics
NumberOfCycles     100000000
NumberOfEquilibrationCycles 100000
PrintEvery          100
RestartFile         no

```

```

ContinueAfterCrash no
WriteBinaryRestartFileEvery 10000
TimeScaleParameterThermostat 0.15

Ensemble NPTPR
NPTPRCellType RegularUpperTriangle

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
TimeStep 0.0005
CutOff 10.0
Forcefield FlexibleHupp-1
EwaldPrecision 1e-6

Framework 0
FrameworkName Hupp-1-mono
Asymmetric yes
SpaceGroup 12
UnitCells 2 2 2
ExternalTemperature 298.0
ExternalPressure 1e5
FlexibleFramework yes
FrameworkDefinitions FlexibleHupp-1-mono

```

## 4.5 Auxiliary examples

### Example 1: Computing the ideal gas Rosenbluth weight of a molecule

To compare simulation values to experiments a reference state should be chosen. A convenient reference state is the ideal gas. The reference Rosenbluth value can be computed from a simulation of a single chain at the desired temperature. Note that for Rosenbluth weights several chains can be computed simultaneously, since they are computed from Widom insertions where the molecule is never actually inserted in the system.

```

SimulationType MonteCarlo
NumberOfCycles 25000
PrintEvery 1000
PrintPropertiesEvery 1000

Forcefield ElenaSodiumCalcium

Box 0
BoxLengths 30 30 30
ExternalTemperature 298.0

Component 0 MoleculeName pentane
MoleculeDefinition TraPPE
WidomProbability 1.0
CreateNumberOfMolecules 0

Component 1 MoleculeName hexane
MoleculeDefinition TraPPE
WidomProbability 1.0
CreateNumberOfMolecules 0

Component 2 MoleculeName heptane
MoleculeDefinition TraPPE
WidomProbability 1.0
CreateNumberOfMolecules 0

Component 3 MoleculeName octane
MoleculeDefinition TraPPE
WidomProbability 1.0
CreateNumberOfMolecules 0

```

The output contains

```

Henry coefficients
  Component 0:      inf [mol/kg/Pa] (Rosenbluth factor new: 0.0197365763 [-])
  Component 1:      inf [mol/kg/Pa] (Rosenbluth factor new: 0.0029490119 [-])
  Component 2:      inf [mol/kg/Pa] (Rosenbluth factor new: 0.0004454807 [-])
  Component 3:      inf [mol/kg/Pa] (Rosenbluth factor new: 0.0000676949 [-])
Energy <U_gh>_1-<U_L>_0 from Widom
  Component 0: 1573.0718943505 [kJ] ( 13.0792510806 [kJ/mol])
  Component 1: 2106.2865896810 [kJ] ( 17.5126459593 [kJ/mol])
  Component 2: 2638.9423414500 [kJ] ( 21.9413935212 [kJ/mol])
  Component 3: 3164.8618412790 [kJ] ( 26.3141327528 [kJ/mol])

```

which is printed every 'PrintPropertiesEvery' cycles. The "Rosenbluth factor new" are the values of interest. The average and error estimated from block averages is printed at the end of the simulation.

### Example 2: Computing the helium void-fraction of a structure (pore volume)

The void fraction is the empty space of a structure divided by the total volume. In experiment it is measured using helium, because helium does (almost) not adsorb. It would be consistent to also measure this fraction

using helium at room temperature. In practice it is easily computed from Widom particle insertion as the void fraction corresponds to the new Rosenbluth weight. The Rosenbluth weight, and therefore the helium void fraction of IRMOF-1 is approximately 0.8255. The pore volume is the void fraction times the unit cell volume. Note that the values dependent slightly on the cutoff, and shifted vs. truncated potentials.

```

SimulationType      MonteCarlo
NumberOfCycles     500000
PrintEvery         1000
PrintPropertiesEvery 1000

Forcefield          IRMOF-1
CutOff             12.8

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0

Component 0 MoleculeName      helium
           MoleculeDefinition TraPPE
           WidomProbability   1.0
           CreateNumberOfMolecules 0

```

### Example 3: Computing the surface area of IRMOF-1

The geometric surface area can easily be computed by 'rolling an atom over the surface' and measure the surface. In practice, for each framework atom points are generate on a sphere around the framework atom, and the amount of overlap with other framework atoms is determined. The fraction of overlap is multiplied times the area of the sphere. The summation over all framework atoms gives the geometric surface area. This example shows how to compute the surface area of IRMOF-1. 'SurfaceAreaSamplingPointsPerSphere' is the amount of points generated on sphere at a distance dependent on the mixing rule, the probe-atom and the current framework atom type. The more points the higher the accuracy. The simulation usually takes between 5 and 30 minutes.

In this example the structure is probed with hydrogen using the second bead ('H\_com' with  $\sigma = 2.958 \text{ \AA}$ ). The option 'SurfaceAreaProbeDistance Sigma' sets the overlap criteria to  $\sigma$  instead of the default  $\sigma^{1/6}$ .

```

SimulationType      MonteCarlo
NumberOfCycles     1000
PrintEvery         100
PrintPropertiesEvery 100

Forcefield IRMOF-1
CutOff             12.8

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
SurfaceAreaProbeDistance Sigma

Component 0 MoleculeName      hydrogen
           StartingBead    1
           MoleculeDefinition TraPPE
           SurfaceAreaProbability 1.0
           CreateNumberOfMolecules 0

```

The area depends on the probe atom and on whether the well-depth at  $2^{1/6}\sigma$  ( $\approx 1.12246\sigma$ ) is used ('SurfaceAreaProbeDistance Minimum')

```

Surface area: 2082.509853 [m^2/cm^3]
Surface area: 3510.189484 [m^2/g]

```

or  $\sigma$  is used as the distance criteria ('SurfaceAreaProbeDistance Sigma'):

```

Surface area: 2266.243128 [m^2/cm^3]
Surface area: 3819.882429 [m^2/g]

```

### Example 4: Computing the surface area of LTA5A with movable ions

For structures with ions also the surface area of the ions are contributing to the measured surface area. When adsorbates are present also the overlap with these adsorbates is taking into account. This allows for a study of the geometric surface area reduction as a function of loading. The procedure is implemented as a Monte Carlo move and therefore one can also use this move to measure the surface area of a flexible framework and/or when movable cations are present. The example here measures the surface area of LTA5A as probed with a methane molecule. The sodium and calcium are moved during the Monte Carlo run.

```

SimulationType MonteCarlo
NumberOfCycles 5000
NumberOfInitializationCycles 1000
PrintEvery 100
PrintPropertiesEvery 100

Forcefield ElenaSodiumCalcium

Framework 0
FrameworkName LTAS5A
UnitCells 1 1 1
ExternalTemperature 298.0

Component 0 MoleculeName sodium
MoleculeDefinition TraPPE
TranslationProbability 1.0
ReinsertionProbability 1.0
ExtraframeworkMolecule yes
CreateNumberOfMolecules 32

Component 1 MoleculeName calcium
MoleculeDefinition TraPPE
TranslationProbability 1.0
ReinsertionProbability 1.0
ExtraframeworkMolecule yes
CreateNumberOfMolecules 32

Component 2 MoleculeName methane
StartingBead 0
BlockPockets yes
BlockPocketsFileName LTAS5A
MoleculeDefinition TraPPE
SurfaceAreaProbability 1.0
CreateNumberOfMolecules 0

```

## Example 5: Powder diffraction pattern

Powder diffraction is a scientific technique using X-Ray or neutron diffraction on powder or microcrystalline samples for structural characterization of materials. The most widespread use of powder diffraction is in the identification and characterization of crystalline solids, each of which produces a distinctive diffraction pattern. Both the positions (corresponding to lattice spacings) and the relative intensity of the lines are indicative of a particular phase and material, providing a "fingerprint" for comparison. The database of IZA for zeolite has the option to generate the powder diffraction pattern:

<http://izasc.ethz.ch/fmi/xsl/IZA-SC/xrd.xsl>

Here, an example of the powder diffraction pattern for the TON-type zeolite. Only one unit cell is sufficient for the computation (interactions are not needed in the computation, just the position and types of the atoms and the shape and size of the unit cell). The diffraction pattern usually takes a few seconds of computation, and the result is written to 'PowderDiffraction/System[0]/'. It contains two files: 'PeakInformation.dat' and 'Spectrum.dat'.

```

SimulationType MonteCarlo
NumberOfCycles 0

Forcefield ElenaSodiumCalcium

Framework 0
FrameworkName TON
UnitCells 1 1 1

ComputePowderDiffractionPattern yes
DiffractionType Xray
DiffractionRadiationType Copper
WaveLengthType single
TwoThetaMin 1
TwoThetaMax 50
TwoThetaStep 0.02
PeakShape PseudoVoigt
PeakWidthModifierU 0.005

```

The first elements of the file 'PeakInformation.dat' look like:

# 2-theta	d	h	k	l	Mult	Lp	Scat. Factor	Intensity
8.15213	0.09220	[ 1, 1, 0]	4	392.85927	19014.2044440544	100.000000		
10.16550	0.11481	[ 0,-2, 0]	2	282.33302	12381.2641234081	20.9311920		
12.77464	0.14431	[ 2, 0, 0]	2	158.63285	19714.5657150741	20.933178		
16.34589	0.18441	[ 2, 2, 0]	4	96.01738	6730.9888808237	8.651941		
16.55216	0.18672	[ -1,-3, 0]	4	93.58434	739.8429009358	0.926889		
19.42690	0.21888	[ -1,-1,-1]	4	67.33674	3040.0925085839	2.740461		
.....								

So, the elements are the angle  $2\theta$ , the  $d$ -spacing, the Miller indices  $h,k$ , and  $l$ , the multiplicity, the Lorentz-Polarization factor, the scattering factor (including anomalous scattering), and the relative intensity (where the largest intensity is set to 100). The second file 'Spectrum.dat' can be plotted using gnuplot, the first column is  $2\theta$ , the second column the intensity. The shape of the peaks can be influenced with 'PeakShape', and the peak width modifiers 'PeakWidthModifierU', 'PeakWidthModifierV', and 'PeakWidthModifierW'.

## Example 6: Making and using 'grids'

For rigid frameworks one can precompute the energy-grid, because the potential energy field induces by the framework does not evolve in time. For each of the pseudo atoms one can generate a 3D grid where the spacing can be defined. In the example the grid points are 0.1 Å spaced apart ( $a=b=c=25.832\text{\AA}$ ,  $258 \times 258 \times 258 = 17173512$  points). A shorter distance results in more points, more accuracy, but also a bigger grid (more memory is needed). Note that RASPA can handle a 'mixture' of grids and fully computed interactions. The table stores  $U, \frac{\partial U}{\partial x}, \frac{\partial U}{\partial y}, \frac{\partial U}{\partial z}, \frac{\partial^2 U}{\partial x \partial y}, \frac{\partial^2 U}{\partial x \partial z}, \frac{\partial^2 U}{\partial y \partial z}$ , and  $\frac{\partial^3 U}{\partial x \partial y \partial z}$  at each grid point. The interpolation can handle non-orthorhombic cells and can also be used for molecular dynamics (i.e. the force interpolation is consistent with the energy interpolation).

```

SimulationType MakeGrid
Forcefield FlexibleIRMOF-1
Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
NumberOfGrids 2
GridTypes C_coo2 0_coo2
SpacingVDWGrid 0.1
SpacingCoulombGrid 0.1

```

The grids are stored in '/share/raspa/grids/FlexibleIRMOF-1/IRMOF-1/0.100000' and the names are 'IRMOF-1\_C\_coo2\_shifted.grid', 'IRMOF-1\_O\_coo2\_shifted.grid', and 'IRMOF-1\_Electrostatics\_Ewald.grid'. The last grid is the real part of the Ewald summation, i.e.  $\text{erfc}(r)/r$  using a probe charge of +1. They can be used like:

```

SimulationType MonteCarlo
NumberOfCycles 5000
NumberOfInitializationCycles 5000
PrintEvery 100

Forcefield FlexibleIRMOF-1
ChargeMethod Ewald
EwaldPrecision 1e-6

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1 1
ExternalTemperature 298.0
ExternalPressure 5000000.0

NumberOfGrids 2
GridTypes C_coo2 0_coo2
SpacingVDWGrid 0.1
SpacingCoulombGrid 0.1
UseTabularGrid yes

Component 0 MoleculeName CO2
MoleculeDefinition TraPPE
TranslationProbability 1.0
RotationProbability 1.0
ReinsertionProbability 1.0
SwapProbability 1.0
CreateNumberOfMolecules 0

```

In the output file, in the framework section, the used grids are tested one by one. Make sure the relative error is smaller than about 0.001 for the energies. If not, either the wrong grid is used (the current settings for the force field, cutoff etc. are different from what the grid has been made with) or the structure requires a higher interpolation density.

```

PseudoAtom 17 Framework-[C_coo2]
=====
Boltzmann average energy VDW (table) : -156.339634664283
Boltzmann average energy VDW (full) : -156.338313481496
Boltzmann relative error VDW : 0.000053898154
Boltzmann average energy Coulomb (table) : -162.513089693416
Boltzmann average energy Coulomb (full) : -162.5064443012736
Boltzmann relative error Coulomb : 0.000043629350

=====
Boltzmann average Force[x] VDW (table) : 21.192078655566
Boltzmann average Force[x] VDW (full) : 21.189556572402
Boltzmann relative error VDW : 0.000372564644
Boltzmann average Force[x] Coulomb (table) : -18.377488985660
Boltzmann average Force[x] Coulomb (full) : -18.400340940728
Boltzmann relative error Coulomb : 0.001062521118

=====
Boltzmann average Force[y] VDW (table) : 11.482227555716
Boltzmann average Force[y] VDW (full) : 11.471251864189
Boltzmann relative error VDW : 0.000613518208
Boltzmann average Force[y] Coulomb (table) : -9.127387761179
Boltzmann average Force[y] Coulomb (full) : -9.116110451771
Boltzmann relative error Coulomb : 0.001224593420

=====
Boltzmann average Force[z] VDW (table) : 15.563260207643
Boltzmann average Force[z] VDW (full) : 15.564130944046
Boltzmann relative error VDW : 0.000679317971

```

```

Boltzmann average Force[z] Coulomb (table)      : -1.488503262903
Boltzmann average Force[z] Coulomb (full)       : -1.495510408023
Boltzmann relative error Coulomb                : 0.001035710819

PseudoAtom 18 Framework-[_O_co2]
=====
Boltzmann average energy VDW (table)           : -369.207117377703
Boltzmann average energy VDW (full)            : -369.204503904417
Boltzmann relative error VDW                  : 0.00026948884
Boltzmann average energy Coulomb (table)        : 89.951068190157
Boltzmann average energy Coulomb (full)         : 89.951352829264
Boltzmann relative error Coulomb              : 0.000045972426
=====
Boltzmann average Force[x] VDW (table)          : 11.109322726969
Boltzmann average Force[x] VDW (full)           : 11.096831498576
Boltzmann relative error VDW                  : 0.000516197616
Boltzmann average Force[x] Coulomb (table)       : -0.340130177955
Boltzmann average Force[x] Coulomb (full)        : -0.327666240972
Boltzmann relative error Coulomb              : 0.001244675144
=====
Boltzmann average Force[y] VDW (table)          : 30.619340781954
Boltzmann average Force[y] VDW (full)           : 30.622572369859
Boltzmann relative error VDW                  : 0.000543276034
Boltzmann average Force[y] Coulomb (table)        : -0.935937742873
Boltzmann average Force[y] Coulomb (full)         : -0.945257821376
Boltzmann relative error Coulomb              : 0.001176433253
=====
Boltzmann average Force[z] VDW (table)          : 1.461304712846
Boltzmann average Force[z] VDW (full)           : 1.467027686101
Boltzmann relative error VDW                  : 0.000601547053
Boltzmann average Force[z] Coulomb (table)       : 5.220361913531
Boltzmann average Force[z] Coulomb (full)        : 5.220055714404
Boltzmann relative error Coulomb              : 0.001308536873

```

## Example 7: Writing and using binary restart "crash-recovery" files

Usually, and unfortunately sometimes often, computers crash, are rebooted to upgrade software or the "walltime"-limit on the cluster has been reached etc. One can force RASPA to write a "binary-restart-file" from which the program can exactly recover and continued where it left off. The results are identical because the data has been written in binary format and even the random number generator picks up where it left off. One has to add two lines to the 'simulation.input' file:

```

ContinueAfterCrash      no
WriteBinaryRestartFileEvery 1000

```

The second line tells the program to write the file every 1000 cycles. Initially, the 'ContinueAfterCrash' is 'no'. For example, the adsorption of methane in MFI (Basic example 6) should be change to

```

SimulationType      MonteCarlo
NumberofCycles     10000
NumberofInitializationCycles 1000
PrintEvery          100

ContinueAfterCrash  no
WriteBinaryRestartFileEvery 1000

Forcefield          ElenaSodiumCalcium

Framework 0
FrameworkName MFI
UnitCells 2 2 2
HeliumVoidFraction 0.29
ExternalTemperature 300.0
ExternalPressure 10000.0 20000.0 30000.0 40000.0

Component 0 MoleculeName      methane
          MoleculeDefinition TraPPE
          TranslationProbability 0.5
          ReinsertionProbability 0.5
          SwapProbability      1.0
          CreateNumberOfMolecules 0

```

It will write a file 'binary\_restart.dat' in the directory 'CrashRestart'. The size of the file is usually small (a few MB). To restart the code, simply change 'ContinueAfterCrash no' to 'ContinueAfterCrash yes'

```

ContinueAfterCrash      yes
WriteBinaryRestartFileEvery 1000

```

## Example 8: Box properties

Sometimes, codes output the cell matrix, but one would like to know the  $\alpha, \beta, \gamma$  angles, or would like to convert it to upper triangle format.

```

SimulationType      MonteCarlo
NumberofCycles     0

Forcefield          IRMOF-1

```

```

BoxMatrix 0
18.214788779 -0.001276607 0.049876430
-0.001736516 24.701353783 0.048044591
0.073461941 0.052027071 26.736241020

```

The output would give these angles and the convert cell matrix

```

Cell angles (degrees) alpha: 89.776368 beta: 89.662047 gamma: 90.007937
Box[0]:
 18.214937000601 -0.003421665049 0.157700318567
 0.000000000000 24.701408369787 0.104376666947
 0.000000000000 0.000000000000 26.735661875881

perpendicular cell widths:
18.21462 24.70122 26.73566
volume of the cell: 12029.299259186328 (A^3)

```

## 4.6 Validation examples

### Example 1: force field of flexible IRMOF-1

RASPA can numerically check the implemented analytical expressions. Using 'SimulationType Numerical' it will numerically compute the forces from the energy, the stress from the energy derivative with respect to strain, the Hessian from the numerical derivative of the (analytical) forces, etc.

```

SimulationType Numerical

ChargeMethod Ewald
InternalFrameworkLennardJonesInteractions yes
CutOff 12.0
Forcefield FlexibleIRMOF-1-new
EwaldPrecision 1e-6

Framework 0
FrameworkName IRMOF-1
UnitCells 1 1
FrameworkDefinitions FlexibleIRMOF-1-new
FlexibleFramework yes
ExternalTemperature 298.0
ExternalPressure 0.0

```

The output is stored in the directory 'Numerical'. The file 'Forces.dat' contains

```

# Force x,y,z, first line analytical, second line minus numerical value
# The force should be the exact opposite to the numerical derivative
# Framework atoms:
# =====
 0  0 [ Mof_Zn] 2389.4714508332 -2389.4714508335 -2389.4714508335
      2389.4714602890 -2389.4714618412 -2389.4714618412
 0  1 [ Mof_Oa] 0.0000000015 0.0000000009 0.0000000002
      -0.0000000761 -0.0000000761 -0.0000000761
 0  2 [ Mof_Ob] -16621.7144673199 16621.7144673199 -4426.2955310233
      -16621.7144601978 16621.7144594217 -4426.2955330002
 0  3 [ Mof_Ca] 0.0000000010 -0.0000000010 12564.8942142472
      -0.0000001552 -0.0000031044 12564.8942085293
 0  4 [ Mof_Cb] 0.0000000008 -0.0000000006 16084.9904923681
      0.00000023283 0.0000015522 16084.9904959761
 0  5 [ Mof_Cc] -5513.1111635094 5513.1111635094 -12190.0205907734
      -5513.1111542384 5513.1111542384 -12190.0205857431
.....
 0 422 [ Mof_Cg] -12190.0205907775 5513.1111635094 -5513.1111635095
      -12190.0206090262 5513.1111713126 -5513.1111759692
 0 423 [ Mof_H] 879.0456246661 -6994.7019088217 6994.7019088218
      879.0456503630 -6994.7019250443 6994.7019172832
Largest difference: 0.0001492886 0.0001516142 0.0001780060

```

Due to symmetry, some of the forces should be zero. The agreement between numerical and analytical forces is excellent. The file 'StrainDerivativeTensor.dat' contains the derivative of energy with respect to strain (if divided by the volume one obtains the stress-tensor)

```

# energy: -4.21242e+06 [K] -35024.1 [kj/mol] -363.008 [eV]
# stress tensor (analytically)
-7.5020560687 -0.0000000000 -0.0000000000
-0.0000000000 -7.5020560687 -0.0000000000
-0.0000000000 -0.0000000000 -7.5020560687

# strain derivative tensor (analytically)
129316.6369463501 0.0000000016 0.0000000015
0.0000000015 129316.6369463478 0.0000000016
0.0000000016 0.0000000015 129316.6369463608

# strain derivative tensor (numerically)
129316.6177502523 0.0017772739 -0.0006053597
0.0012960906 129316.6239435474 0.0007916242
-0.0017927960 0.0008537124 129316.6258527587

```

The second derivative of the energy with respect to strain is written to the file 'StrainSecondDerivativeTensor.dat'

```
# energy: -4.21242e+06 [K] -35024.1 [kJ/mol] -363.008 [eV]

# strain second derivative tensor (analytically)
 80356775.0198      0.0000      0.0000      0.0000      30174777.1276      -0.0000      0.0000      -0.0000      30174777.1276
      0.0000      22816718.9605     -0.0000      30174777.1276      -0.0000      -0.0000      -0.0000      -0.0000      -0.0000
      0.0000     -0.0000      22816718.9605      -0.0000      -0.0000      -0.0000      30174777.1276      -0.0000      -0.0000
      0.0000      30174777.1276     -0.0000      22816718.9605      -0.0000      -0.0000      -0.0000      -0.0000      -0.0000
 30174777.1276     -0.0000     -0.0000     -0.0000      80356775.0198      -0.0000      -0.0000      -0.0000      30174777.1276
      -0.0000     -0.0000     -0.0000     -0.0000      22816718.9605      -0.0000      -0.0000      30174777.1276      -0.0000
      0.0000     -0.0000      30174777.1276     -0.0000     -0.0000     -0.0000      22816718.9605      -0.0000      -0.0000
      -0.0000     -0.0000     -0.0000     -0.0000      30174777.1276     -0.0000      -0.0000      22816718.9605      -0.0000
 30174777.1276     -0.0000     -0.0000     -0.0000      30174777.1276     -0.0000      -0.0000      -0.0000      80356775.0198

# strain second derivative tensor (numerically from analytical first derivative)
 80356768.7345     -0.0880     -0.0462     -0.1950      30174780.1304     -0.0491      0.8695     -0.0124      30174780.1228
      -0.0896     22816713.9256     -0.0625      30174780.1283     -0.4853     -0.0450     -0.0155      0.0589     -0.0506
      -0.0462     -0.0605      22816713.8055     -0.0544     -0.0534     -0.7676      30174780.1277     -0.0458      0.1841
      -0.1950      30174780.1304     -0.0491     22816713.9263     -0.0348      0.0396      0.0985     -0.0247      0.0170
 30174780.1283     -0.4853     -0.0450     -0.0286      80356765.8023     -0.0001     -0.0206      0.8518      30174780.0874
      -0.0544     -0.0534     -0.7676      0.0449      0.0028     22816712.5191     -0.0163      30174780.0828     -0.2401
      0.8695     -0.0124      30174780.1228     0.0985     -0.0247     -0.0170     22816712.5046     -0.0741     -0.0132
      -0.0155     -0.0589     -0.0506     -0.0206     0.8518      30174780.0874     -0.0839     22816713.8761     0.0249
 30174780.1277     -0.0458     0.1841     -0.0163      30174780.0828     0.2401     -0.0039     0.0271      80356767.6051
```

The second derivative of energy with respect to position is written to the file 'HessianMatrix.dat'

```

element [ 0,  0] analytically: 153081.336214122 numerically: 153081.336260233 difference: -0.0000136110
element [ 0,  1] analytically: -29039.7692839710 numerically: -29039.7692822353 difference: -0.0000017357
element [ 0,  2] analytically: -29039.7692839710 numerically: -29039.7692818260 difference: -0.0000021450
element [ 0,  3] analytically: -62196.8575382852 numerically: -62196.8575638249 difference: -0.0000014575
element [ 0,  4] analytically: 42468.2617792162 numerically: 42468.2617813368 difference: -0.0000021206
element [ 0,  5] analytically: 42468.2617792163 numerically: 42468.2617813176 difference: -0.0000021014
element [ 0,  6] analytically: -8591.9119729929 numerically: -8591.9119817845 difference: 0.0000025546
element [ 0,  7] analytically: 1657.2381758880 numerically: 1657.2381700826 difference: 0.0000058054
element [ 0,  8] analytically: -10462.2229115035 numerically: -10462.2229124288 difference: 0.0000009253
.
.
.
element [1271,1266] analytically: -65951.245989043 numerically: -65951.245949379 difference: -0.0000045065
element [1271,1267] analytically: 107130.5503696070 numerically: 107130.5503646727 difference: 0.0000493535
element [1271,1268] analytically: -143472.332964186 numerically: -143472.332867877 difference: -0.0000096308
element [1271,1269] analytically: 62021.1807794664 numerically: 62021.1807737405 difference: 0.0000057250
element [1271,1270] analytically: -117418.6021011203 numerically: -117418.6020937668 difference: -0.0000073535
element [1271,1271] analytically: 138795.1804072921 numerically: 138795.1803974829 difference: 0.0000098090
# largest difference: -0.0001407942

```

Finally, the derivative of the strain-derivative with respect to position is written to the file 'CrossTerms.dat'

# 0	-71077.0942744063	-81184.5391161912	-81184.5391161913	-83574.0105670245	81184.5391161911	41016.8235549702	-83574.0105670246	41016.8235549702	81184.5391161911
-71077.0943362149	-81184.5391080396	-81184.5391124962	-83574.0104962800	81184.5391075849	41016.8235534229	-83574.0104962800	41016.8235555981	81184.5391135421	
# 1	-81184.5391161912	83574.0105670242	41016.8235549702	81184.5391161911	71077.0942744067	-81184.5391161906	41016.8235549702	-83574.0105670239	-81184.5391161907
-81184.5391119353	83574.0104771200	41016.8235445023	81184.5391080851	71077.0942428401	-81184.5391082518	41016.8235595317	-83574.0104766349	-81184.5391120717	
# 2	-81184.5391161913	41016.8235549702	83574.0105670243	41016.8235549702	-81184.5391161906	-83574.0105670240	81184.5391161911	-81184.5391161907	71077.0942744060
-81184.5391119429	41016.8235488679	83574.0104693589	41016.8235566663	-81184.5391121324	-83574.0104693589	81184.5391076607	-81184.5391081912	71077.0943044432	
# 3	0.0000000039	0.0000000035	0.0000000010	0.0000000043	0.0000000059	-22435.6859250378	0.0000000013	-22435.6859250378	0.0000000062
0.0000111565	-0.0000043732	0.0000070334	0.00000936173	0.0000000606	-22435.6859038461	-0.0000383200	-224353.6859058243	0.0000009701	
# 1270	-75523.8770946184	23026.6334757857	-22892.0977894283	34666.8500027566	-69812.9971667970	76856.2176788618	-35411.3599491967	83850.9197952409	-76856.2179012722
-75523.8768266905	23026.6338757550	-22892.0981259162	34666.8500182329	-69812.9971630503	76856.2178281586	-35411.3600019153	83850.9198236958	-76856.2178674664	
# 1271	75523.8771188713	-22892.0978030707	23026.6334879117	-35411.3599669571	76856.2177031150	-83850.9196425578	34666.8499880272	-76856.2178891456	69812.9973316051
75523.8768230393	-22892.0981239982	23026.6338262785	-35411.3599794076	76856.2178334789	-83850.9196917584	34666.8500302077	-76856.2178665950	69812.9974499655	



# Chapter 5

## The source code

### 5.1 Introduction

### 5.2 Data types

There are several new types, the two most important ones are

- REAL

REAL is a floating point number. It is defined in 'src/constants.h' as

```
#define REAL double
```

but if one needs higher precision one could use

```
#define REAL long double
```

and using the 'qd' library it is even possible to use arbitrary precision.

- VECTOR

An structure with three elements 'x', 'y', and 'z'.

```
typedef struct point
{
    REAL x;
    REAL y;
    REAL z;
} POINT,VECTOR;
```

- REAL\_MATRIX3x3

A  $3 \times 3$  matrix, used as transformations on vectors (like 'strain') and for the three cell-vectors making up the cell matrix. It is defined in 'src/matrix.h'.

```
typedef struct real_matrix3x3
{
    REAL ax;
    REAL ay;
    REAL az;

    REAL bx;
```

```

    REAL by;
    REAL bz;

    REAL cx;
    REAL cy;
    REAL cz;
} REAL_MATRIX3x3;

```

## 5.3 Datastructures

### Box properties and periodic boundaries

For each system, a cell box and other properties are defined in 'src/simulation.h'

```

REAL_MATRIX3x3 *Box;                                // the cell matrix
REAL_MATRIX3x3 *InverseBox;                          // the inverse of the cell matrix
REAL_MATRIX3x3 *ReplicaBox;                         // the cell matrix of the replica system
REAL_MATRIX3x3 *InverseReplicaBox;                  // the inverse of the the cell matrix of the replica system
INT_VECTOR3 *NumberOfReplicaCells;                  // the integere number of replicas in each direction a,b,c
int *TotalNumberOfReplicaCells;                     // the total number of replica cells
VECTOR *ReplicaShift;                             // the shift in a,b,c for each replica cell
int *UseReplicas;                                 // whether or not to use replicas
REAL_MATRIX3x3 *BoxProperties;                     // properties of the cell matrix (i.e. perpendicular lengths)
REAL_MATRIX3x3 *InverseBoxProperties;              // properties of the inverse cell matrix
REAL *Volume;                                     // the volume
REAL *AlphaAngle;                                // the alpha-angle of the cell
REAL *BetaAngle;                                 // the beta-angle of the cell
REAL *GammaAngle;                                // the gamma-angle of the cell
int *BoundaryCondition;                           // the boundary condition (i.e. 'RECTANGULAR' or 'TRICLINIC')

```

These are dynamically allocated arrays and have the same length as the amount of systems present. For example, in a Gibbs simulation two systems are needed, one for the gas-phase and one for the liquid phase. 'Volume[0]' would give the volume of the first cell, and 'Volume[1]' would give the volume of the second cell.

Periodic boundaries are applied after each distance computation calling the function 'ApplyBoundaryCondition' (defined in 'src/potentials.h') It operates on a 'VECTOR' and give the corrected vector back. The system is specified with the global variable 'CurrentSystem'.

```

VECTOR ApplyBoundaryCondition(VECTOR dr)
{
    VECTOR s,t;

    switch(BoundaryCondition[CurrentSystem])
    {
        case FINITE:
            break;
        case RECTANGULAR:
        case CUBIC:
            dr.x-=Box[CurrentSystem].ax*(REAL)NINT(dr.x*InverseBox[CurrentSystem].ax);
            dr.y-=Box[CurrentSystem].by*(REAL)NINT(dr.y*InverseBox[CurrentSystem].by);
            dr.z-=Box[CurrentSystem].cz*(REAL)NINT(dr.z*InverseBox[CurrentSystem].cz);
            break;
        case TRICLINIC:
            // convert from xyz to abc
            s.x=InverseBox[CurrentSystem].ax*dr.x+InverseBox[CurrentSystem].bx*dr.y+InverseBox[CurrentSystem].cx*dr.z;
            s.y=InverseBox[CurrentSystem].ay*dr.x+InverseBox[CurrentSystem].by*dr.y+InverseBox[CurrentSystem].cy*dr.z;
            s.z=InverseBox[CurrentSystem].az*dr.x+InverseBox[CurrentSystem].bz*dr.y+InverseBox[CurrentSystem].cz*dr.z;

            // apply boundary condition
            t.x=s.x-(REAL)NINT(s.x);
            t.y=s.y-(REAL)NINT(s.y);
            t.z=s.z-(REAL)NINT(s.z);
    }
}

```

```

    // convert from abc to xyz
    dr.x=Box[CurrentSystem].ax*t.x+Box[CurrentSystem].bx*t.y+Box[CurrentSystem].cx*t.z;
    dr.y=Box[CurrentSystem].ay*t.x+Box[CurrentSystem].by*t.y+Box[CurrentSystem].cy*t.z;
    dr.z=Box[CurrentSystem].az*t.x+Box[CurrentSystem].bz*t.y+Box[CurrentSystem].cz*t.z;
    break;
default:
    fprintf(stderr,"Error: Unkown boundary condition....\n");
    exit(0);
    break;
}
return dr;
}

```

The function 'NINT' is faster version of 'rint' (or 'floor').

```
#define NINT(x) ((int)((x)>=0.0?((x)+0.5):((x)-0.5)) )
```

A common occurrence of the boundary conditions application is for two positions of atoms 'posA' and 'posB' (of type 'VECTOR')

```

dr.x=posA.x-posB.x;
dr.y=posA.y-posB.y;
dr.z=posA.z-posB.z;
dr=ApplyBoundaryCondition(dr);
rr=SQR(dr.x)+SQR(dr.y)+SQR(dr.z);
r=sqrt(rr);

```

There are functions you can use to transform from Cartesian to fractional coordinates (defined in 'src/potentials.h')

```

VECTOR ConvertFromXYZtoABC(VECTOR t)
{
    VECTOR s;

    s.x=InverseBox[CurrentSystem].ax*t.x+InverseBox[CurrentSystem].bx*t.y+InverseBox[CurrentSystem].cx*t.z;
    s.y=InverseBox[CurrentSystem].ay*t.x+InverseBox[CurrentSystem].by*t.y+InverseBox[CurrentSystem].cy*t.z;
    s.z=InverseBox[CurrentSystem].az*t.x+InverseBox[CurrentSystem].bz*t.y+InverseBox[CurrentSystem].cz*t.z;
    return s;
}

```

and from fractional coordinates to Cartesian

```

VECTOR ConvertFromABCtoXYZ(VECTOR t)
{
    VECTOR dr;

    dr.x=Box[CurrentSystem].ax*t.x+Box[CurrentSystem].bx*t.y+Box[CurrentSystem].cx*t.z;
    dr.y=Box[CurrentSystem].ay*t.x+Box[CurrentSystem].by*t.y+Box[CurrentSystem].cy*t.z;
    dr.z=Box[CurrentSystem].az*t.x+Box[CurrentSystem].bz*t.y+Box[CurrentSystem].cz*t.z;
    return dr;
}

```

## (Pseudo-)atoms

The data structure 'PSEUDO\_ATOM' contains information on atoms, either real atoms or united atoms where several atoms are lumped together (for example: CH3).

```

// Pseudoatoms
typedef struct PseudoAtom
{
    char Name[256];           // the Name of the pseudo-atom ('CH3', 'H', 'O' etc).
    char PrintToPDBName[256];  // the string to print to a pdb-file as name
    int PrintToPDB;            // whether to write this atom to the pdf-file or not
    char ChemicalElement[256]; // the chemical element ('O', 'H', etc)
    int ScatteringType;        // the scattering type (powder diffraction)
    int AnomalousScatteringType; // the anomalous scattering type (powder diffraction)
    REAL TemperatureFactor;    // the temperature factor (powder diffraction)

```

```

REAL Mass;           // the mass of the pseudo-atom
REAL Charge;        // the charge of the pseudo-atom
REAL Polarization; // the polarization of the atom
int HasCharges;    // whether or not the atom has atoms with charges
int IsPolarizable; // whether or not the atom has a induced point dipole
int Interaction;   // whether or not the atom has interactions
REAL Radius;        // the radius (used for calculating Bonds in the zeolite)
int Connectivity;  // the connectivity (used for calculating Bonds/Bends/Torsion in the framework)
} PSEUDO_ATOM;

```

A typical use is, once the type is known, to retrieve the charge for a pseudo-atoms:

```

REAL q;
q=PseudoAtom[type].Charge;

```

Use the following to find out to what pseudoatom a string corresponds to

```

int type;
type=ReturnPseudoAtomNumber("CH4");

```

However, usually the type is a property of each of the atoms of a molecule.

```

int type;
type=Framework[1].Atoms[0][10].Type;

```

and 'type' can then be used to get the mass, charge, polarization, etc. Here, the type is retrieve for atom number 11 (c is starting from 0, unlike Fortran) of the first framework of the second system.

## Framework

Atoms make up a framework, several frameworks can make up 1 system. The definition of a framework atom 'FRAMEWORK\_ATOM' is

```

typedef struct framework_atom
{
    int Type;           // the pseudo-atom type of the atom
    int AssymmetricType; // the 'asymmetric' type

    // MC/MD properties
    POINT Position;      // the position of the atom
    POINT ReferencePosition; // the 'reference' position of the atom

    // MD properties
    VECTOR Velocity;     // the velocity of the atom
    VECTOR ReferenceVelocity; // the 'reference' velocity of the atom
    VECTOR Force;         // the force acting on the atom

    VECTOR ElectricField; // the electricfield vector
    VECTOR ReferenceElectricField; // the 'reference' electricfield vector
    VECTOR InducedElectricField; // the induced electric field
    VECTOR InducedDipole;    // the induced dipole moment on this atom
    int HessianIndex;       // the index in the Hessian matrix for this atom
} FRAMEWORK_ATOM;

```

It contains the properties you'd expect, like type, position, velocity, and force. For polarization, also electric field, induced electric field, and induced dipole are needed. For many applications, one needs to backup the positions and/or velocities. The field 'ReferencePosition' and 'ReferenceVelocity' are useful for that. Also they can be used for some algorithms which need the 'old' values to. An example is the numerical computation of stress. First all positions are copied to the 'ReferencePosition', then the positions 'Position' are generated from the strain at infinite small strain difference and the finite difference scheme is applied. A framework-structure 'FRAMEWORK\_COMPONENT' is defined per system

```
FRAMEWORK_COMPONENT *Framework;
```

with

```
typedef struct FrameworkComponent
{
    char (*Name)[256];                                // the name of the frameworks

    int TotalNumberOfAtoms;                           // the total number of atoms of the frameworks
    int TotalNumberOfUnitCellAtoms;                  // the total number of atoms of the unit cell
    REAL FrameworkDensity;                          // the total density of the frameworks
    REAL FrameworkMass;                            // the total mass of the frameworks

    int NumberOfFrameworks;                         // the number of frameworks
    REAL *FrameworkDensityPerComponent;           // the density per framework
    REAL *FrameworkMassPerComponent;              // the mass per framework

    int *NumberOfAtoms;                            // the number of atoms per framework
    int *NumberOfUnitCellAtoms;                   // the number of unit cell atoms per framework
    FRAMEWORK_ATOM **Atoms;                      // list of framework-atoms per framework
    .....
    .....
} FRAMEWORK_COMPONENT;
```

The structure had the element 'Atoms' which is a list of framework-atoms per framework. So, to get the type of the 11 atom of the first framework of the second system, use

```
int type;
type=Framework[1].Atoms[0][10].Type;
```

Finally, a small example where we print out the positions of all the framework atoms for all frameworks and systems

```
int i,j,f1;
for(i=0;i<NumberOfSystem;i++)
{
    for(f1=0;f1<Framework[i].NumberOfSystems;f1++)
    {
        for(j=0;j<Framework[i].NumberOfAtoms[f1];j++)
        printf("system: %d framework: %d atom: %d -> position: %g %g %g\n",
        i,f1,j,
        Framework[i].Atoms[f1][j].Position.x,
        Framework[i].Atoms[f1][j].Position.y,
        Framework[i].Atoms[f1][j].Position.z);
    }
}
```

## Components

Everything that is independent of a molecule's positions but still a property of molecules is stored in the structure 'COMPONENT'. Here you find the number of atoms for this type of molecule per system, the mass for the component etc. Also computed values for densities of the bulk fluid, compressibility, and the amount of excess molecules are stored. These are computed from the mol fraction, pressure, and critical pressure/temperature and acentric factor. After these properties there are data on the potentials defined for the component: bond, Urey-Bradley, bends, torsions, cross-terms, intra Van der Waals etc. For Monte Carlo the structure contains the probability of all the moves.

```
typedef struct Component
{
```

```

char Name[256];                                // the name of the component ("methane", "C12", "propane" etc).
int NumberOfAtoms;                             // the number of atoms in the component
int StartingBead;                            // the bead of the molecule used for starting the growing process in CBMC
REAL Mass;                                    // the mass of the component
int *NumberOfMolecules;                      // the number of molecules of the component for each system
int *Type;                                     // the pseudo-atom Type of each atom
int *Connectivity;                            // the connectivity of each atom
int HasCharges;                             // whether the molecule contains charges or not
int IsPolarizable;                           // whether the molecule has point dipoles or not
int ExtraFrameworkMolecule;                  // TRUE: Cation, FALSE: Adsorbate
int Swapable;                                 // whether or not the number of molecules is fluctuating (i.e. GCMC)
int Widom;                                    // whether this component is used for Widom insertions

REAL *IdealGasRosenbluthWeight; // the Rosenbluth weight of an ideal-chain per system
REAL *IdealGasTotalEnergy;   // the total energy of an ideal-chain per system

REAL *PartialPressure;                     // the partial pressure of the component per system
REAL *FugacityCoefficient;                // the fugacity coefficient of the component per system
REAL *BulkFluidDensity;                  // the bulkfluid-density of the component per system
REAL *Compressibility;                   // the compressibility of the fluid-fase per system
REAL *MolFraction;                       // the mol-fraction of the component per system
REAL *AmountOfExcessMolecules;          // the amount of excess molecules per system

REAL CriticalTemperature; // the critical temperature of the component
REAL CriticalPressure;   // the critical pressure of the component
REAL AcentricFactor;      // the acentric factor of the component

int NumberOfGroups;                      // the number of groups
GROUP_DEFINITION *Groups;                // the definition of the groups
int *group;                            // to which group an atom belongs
VECTOR *Positions;                     // the positions in the body-fixed frame
.....
.....
int NumberOfBonds;                      // the number of bonds of the component
PAIR *Bonds;                            // the list of bond-pairs
int *BondType;                          // the type of the bond for each bond-pair
REAL (*BondArguments)[MAX_BOND_POTENTIAL_ARGUMENTS]; // the arguments needed for this bond-pair
.....
.....
REAL ProbabilityTranslationMove; // the probability of the translation MC-move for the component
REAL ProbabilityRotationMove;  // the probability of the rotation MC-move for the component
REAL ProbabilityCBMCMove;       // the probability of the partial-regrow MC-move for the component
REAL ProbabilityReinsertionMove; // the probability of the reinsertion MC-move for the component
.....
.....
} COMPONENT;

```

A component consists of 'groups', which is a collection of atoms that are either treated as rigid or as flexible. The component has elements that lists how many of these groups there are, the definition of the group, and the positions of all the atoms in the body-fixed frame. The definition of the group is the structure 'GROUP\_DEFINITION'. Important elements are whether or not the group is rigid, the number of atoms in the group, and the list of atom number present in the groups.

```

typedef struct group_definitions
{
    int Rigid;                                // whether or not the group is rigid
    int Type;                                  // the type, NONLINEAR_MOLECULE, LINEAR_MOLECULE, or POINT_PARTICLE

    REAL Mass;                                 // the mass of the group

    int NumberOfGroupAtoms;                    // the numer of atoms in the group
    int *Atoms;                                // the atoms in the group

    REAL_MATRIX3x3 InertiaTensor;             // the inertia tensor

```

```

VECTOR InertiaVector;           // the inertia vector
VECTOR InverseInertiaVector;   // the inverse of inertia vector

REAL_MATRIX3x3 RotationalMatrix; // the rotational matrix
TRIPLE orientation;           // three atoms A,B,C to compute quaternions
REAL rot_min;

int RotationalDegreesOfFreedom; // the rotational degrees of freedom
} GROUP_DEFINITION;

```

The inertia tensor, vector and rotational matrix etc. are the same for a certain type of molecule. Together with the actually atom positions, the orientations can be computed for all the rigid units (i.e. the quaternions are computed).

## Adsorbate and cations

The definition of an adsorbate atom 'ADSORBATE\_ATOM' is very similar to a framework atom

```

typedef struct adsorbate_atom
{
    int Type;                      // the pseudo-atom type of the atom

    // MC/MD properties
    POINT Position;               // the position of the atom
    POINT ReferencePosition;       // the 'reference' position of the atom

    // MD properties
    VECTOR Velocity;              // the velocity of the atom
    VECTOR ReferenceVelocity;     // the 'reference' velocity of the atom
    VECTOR Force;                 // the force acting on the atom

    VECTOR ElectricField;          // the electricfield vector
    VECTOR ReferenceElectricField; // the 'reference' electricfield vector
    VECTOR InducedElectricField;  // the induced electric field
    VECTOR InducedDipole;         // the induced dipole moment on this atom
    int HessianIndex;             // the index in the Hessian matrix for this atom
} ADSORBATE_ATOM;

```

The definition for cations is identical except it is called 'CATION\_ATOM'. The definition of an adsorbate molecule is

```

typedef struct adsorbate
{
    int Type;                      // the component type of the molecule
    int NumberOfAtoms;              // the number of atoms in the molecule
    GROUP *Groups;                 // data of the rigid groups
    ADSORBATE_ATOM *Atoms;         // list of atoms
} ADSORBATE_MOLECULE;

```

The definition of a cation is called 'CATION\_MOLECULE'. Note that a molecule can consists of atoms, but also can contain rigid units. The atoms are accessible through the 'Atoms' field, and rigid units are accessible through the 'Groups' field. A 'GROUP' consists of

```

typedef struct group
{
    REAL Mass;                     // mass of the rigid unit
    QUATERNION Quaternion;         // orientation of the unit
    QUATERNION QuaternionMomentum; // quaternion momentum
    QUATERNION QuaternionForce;    // quaternion force
    VECTOR Torque;                // torque vector
    VECTOR CenterOfMassPosition;  // the center of mass position
    VECTOR CenterOfMassReferencePosition; // the reference position for the center of mass
    VECTOR CenterOfMassVelocity;   // the center of mass velocity
    VECTOR CenterOfMassForce;     // the center of mass force
    VECTOR AngularVelocity;       // the angular velocity of the rigid unit
} GROUP;

```

which contains elements like position and orientation, and fields for the integration of rigid units, i.e. QuaternionMomentum etc.

Molecules are stored as a list of molecules for each system

```
ADSORBATE_MOLECULE **Adsorbates;
```

To get the type of the 5th atom of the 11th adsorbate of the first system, use

```
int type;
type=Adsorbates[0][10].Atoms[4].Type;
```

As an example, here a function to measure the velocity drift of all the adsorbates in the current system

```
VECTOR MeasureVelocityDrift(void)
{
    int i,k,l,Type,A,f;
    REAL Mass,TotalMass;
    VECTOR com;

    TotalMass=0.0;
    com.x=com.y=com.z=0.0;
    for(i=0;i<NumberOfAdsorbateMolecules[CurrentSystem];i++)
    {
        Type=Adsorbates[CurrentSystem][i].Type;
        for(l=0;l<Components[Type].NumberOfGroups;l++)
        {
            if(Components[Type].Groups[l].Rigid)
            {
                Mass=Components[Type].Groups[l].Mass;
                TotalMass+=Mass;
                com.x+=Mass*Adsorbates[CurrentSystem][i].Groups[l].CenterOfMassVelocity.x;
                com.y+=Mass*Adsorbates[CurrentSystem][i].Groups[l].CenterOfMassVelocity.y;
                com.z+=Mass*Adsorbates[CurrentSystem][i].Groups[l].CenterOfMassVelocity.z;
            }
            else
            {
                for(k=0;k<Components[Type].Groups[l].NumberOfGroupAtoms;k++)
                {
                    A=Components[Type].Groups[l].Atoms[k];
                    Mass=PseudoAtoms[Adsorbates[CurrentSystem][i].Atoms[A].Type].Mass;
                    TotalMass+=Mass;
                    com.x+=Mass*Adsorbates[CurrentSystem][i].Atoms[A].Velocity.x;
                    com.y+=Mass*Adsorbates[CurrentSystem][i].Atoms[A].Velocity.y;
                    com.z+=Mass*Adsorbates[CurrentSystem][i].Atoms[A].Velocity.z;
                }
            }
        }
        com.x/=TotalMass;
        com.y/=TotalMass;
        com.z/=TotalMass;
    }
    return com;
}
```

It loops over all the adsorbate molecules, and asks for the type. The component-type is important to get the number of groups for the current molecule. Then, there is a inner loop over all of the groups of the current molecule. If the group is rigid, then the center of mass velocity is used, otherwise it is flexible and it loops over all the atoms of the flexible group. In general, if something is the same for a type of molecule then it is a property of the component. If it is different for each molecule, it is a property of a molecule.

## 5.4 Modifying

### 5.4.1 Monte Carlo

#### Selecting MC moves

The file 'src/monte\_carlo.c' is the main Monte Carlo simulation routine. The bulk of the code deals with how to select a particular Monte carlo move. Some requirements and conveniences:

- The moves should be chosen in random order
- System move should be chosen much less frequent than particle moves. The particles need to be able to adapt to the new system.
- For  $n$  systems, the amount of steps should be  $n$  times larger.
- For  $n$  times as many molecules, the amount of steps should be  $n$  times larger.
- For multi-component systems one needs more steps.
- For systems at low loadings, the sampling lengths should be increase a bit (i.e. set a minimum amount of inner steps).
- The relative probabilities of particle moves should be taken into account.

A code which achieves all the above is listed here (there are many other ways of doing this). For each MC 'cycle'

```
for(i=0;i<NumberOfSystems;i++)
{
    // choose system at random
    CurrentSystem=(int)(RandomNumber()*(REAL)NumberOfSystems);

    NumberOfSystemMoves=9;
    NumberOfMolecules=NumberOfAdsorbateMolecules[CurrentSystem]+NumberOfCationMolecules[CurrentSystem];
    NumberOfParticleMoves=MAX(MinimumInnerCycles,NumberOfMolecules);
    NumberOfSteps=(NumberOfSystemMoves+NumberOfParticleMoves)*NumberOfComponents;

    // loop over the MC 'steps' per MC 'cycle'
    for(j=0;j<NumberOfSteps;j++)
    {
        // choose any of the MC moves randomly
        ran_int=(int)(RandomNumber()*NumberOfSteps);
        switch(ran_int)
        {
            case 0: if(RandomNumber()<ProbabilityParallelTemperingMove) ParallelTemperingMove(); break;
            case 1: if(RandomNumber()<ProbabilityHybridNVEMove) HybridNVEMove(); break;
            case 2: if(RandomNumber()<ProbabilityHybridNPHMove) HybridNPHMove(); break;
            case 3: if(RandomNumber()<ProbabilityHybridNPHPMove) HybridNPHPMove(); break;
            case 4: if(RandomNumber()<ProbabilityVolumeChangeMove) VolumeMove(); break;
            case 5: if(RandomNumber()<ProbabilityBoxShapeChangeMove) BoxShapeChangeMove(); break;
            case 6: if(RandomNumber()<ProbabilityGibbsVolumeChangeMove) GibbsVolumeMove(); break;
            case 7: if(RandomNumber()<ProbabilityFrameworkChangeMove) FrameworkChangeMove(); break;
            case 8: if(RandomNumber()<ProbabilityFrameworkShiftMove) FrameworkShiftMove(); break;
            default:
                // choose component at random
                CurrentComponent=(int)(RandomNumber()*(REAL)NumberOfComponents);

                // choose the Monte Carlo move at random
                ran=RandomNumber();
                if(ran<Components[CurrentComponent].ProbabilityTranslationMove) TranslationMove();
                else if(ran<Components[CurrentComponent].ProbabilityRandomTranslationMove) RandomTranslationMove();
                else if(ran<Components[CurrentComponent].ProbabilityRotationMove) RotationMove();
                else if(ran<Components[CurrentComponent].ProbabilityCBMCMove) CBMCMove();
```

```

        else if(ran<Components[CurrentComponent].ProbabilityReinsertionMove) ReinsertionMove();
        else if(ran<Components[CurrentComponent].ProbabilityReinsertionInPlaceMove) ReinsertionInPlaceMove();
        else if(ran<Components[CurrentComponent].ProbabilityReinsertionInPlaneMove) ReinsertionInPlaneMove();
        else if(ran<Components[CurrentComponent].ProbabilityIdentityChangeMove) IdentityChangeMove();
        else if(ran<Components[CurrentComponent].ProbabilitySwapMove)
        {
            if(RandomNumber()<0.5) SwapAddMove();
            else SwapRemoveMove();
        }
        else if(ran<Components[CurrentComponent].ProbabilityWidomMove) WidomMove();
        else if(ran<Components[CurrentComponent].ProbabilitySurfaceAreaMove) SurfaceAreaMove();
        else if(ran<Components[CurrentComponent].ProbabilityGibbsSwapChangeMove) GibbsParticleTransferMove();
        else if(ran<Components[CurrentComponent].ProbabilityGibbsIdentityChangeMove) GibbsIdentityChangeMove();
        break;
    }
}
}

```

First is a loop over the amount of systems, and a random system is chosen. Suppose we have 200 single component molecules in this system, then each of the system move is chosen with 1/209 probability (case 0-8), and there is a 200/209 chance to select a particle move (case 9-209). The probability of the particle moves are scaled in such a way that the proper relative occurrence is obeyed (as specified in the input). Note that the swap-move has 50% to be swap insertion and 50% to be swap remove. This is necessary to obey detailed balance. For multi-components more moves are performed.

### Sampling properties during Monte Carlo

The Monte Carlo routine has two parts:

- The initialization part. Here, no properties are computed and MC moves are performed just to reach equilibrium.
- The production run, where properties are computed.

The basic outline of the production run is

```

// initialize sampling-routines at the start of the production run
SampleInfraRedSpectra(INITIALIZE);
SampleMeanSquareDisplacementOrderN(INITIALIZE);
SampleOnsagerMeanSquareDisplacementOrderN(INITIALIZE);
SampleRadialDistributionFunction(INITIALIZE);
SampleFrameworkSpacingHistogram(INITIALIZE);
SamplePositionHistogram(INITIALIZE);
SampleNumberOfMoleculesHistogram(INITIALIZE);
SampleEnergyHistogram(INITIALIZE);
SampleDensityProfile3DVTKGrid(INITIALIZE);
SampleEndToEndDistanceHistogram(INITIALIZE);
SampleMoleculePropertyHistogram(INITIALIZE);
SamplePDBMovies(INITIALIZE);
SampleDcTSTConfigurationFiles(INITIALIZE);
SampleFreeEnergyProfile(INITIALIZE);
SampleCationAndAdsorptionSites(INITIALIZE);

for(CurrentCycle=0;CurrentCycle<NumberOfCycles;CurrentCycle++)
{
    // sample energy average and system/particle properties
    for(CurrentSystem=0;CurrentSystem<NumberOfSystems;CurrentSystem++)
    {
        UpdateEnergyAveragesCurrentSystem();

        SampleRadialDistributionFunction(SAMPLE);
        SampleFrameworkSpacingHistogram(SAMPLE);
        SamplePositionHistogram(SAMPLE);
    }
}

```

```

        SampleNumberOfMoleculesHistogram(SAMPLE);
        SampleEnergyHistogram(SAMPLE);
        SampleDensityProfile3DVTKGrid(SAMPLE);
        SampleEndToEndDistanceHistogram(SAMPLE);
        SampleMoleculePropertyHistogram(SAMPLE);
        SampleFreeEnergyProfile(SAMPLE);
        SampleCationAndAdsorptionSites(SAMPLE);
    }

    // SELECTION OF MC-MOVES (SEE CODE OF THE PREVIOUS SECTION)

    for(CurrentSystem=0;CurrentSystem<NumberofSystems;CurrentSystem++)
    {
        SampleRadialDistributionFunction(PRINT);
        SampleFrameworkSpacingHistogram(PRINT);
        SamplePositionHistogram(PRINT);
        SampleNumberOfMoleculesHistogram(PRINT);
        SampleEnergyHistogram(PRINT);
        SampleDensityProfile3DVTKGrid(PRINT);
        SampleEndToEndDistanceHistogram(PRINT);
        SampleMoleculePropertyHistogram(PRINT);
        SamplePDBMovies(PRINT);
        SampleDCTSTConfigurationFiles(PRINT);
        SampleFreeEnergyProfile(PRINT);
        SampleCationAndAdsorptionSites(PRINT);
    }
}

// finalize output
SampleRadialDistributionFunction(FINALIZE);
SampleFrameworkSpacingHistogram(FINALIZE);
SamplePositionHistogram(FINALIZE);
SampleNumberOfMoleculesHistogram(FINALIZE);
SampleEnergyHistogram(FINALIZE);
SampleDensityProfile3DVTKGrid(FINALIZE);
SampleEndToEndDistanceHistogram(FINALIZE);
SampleMoleculePropertyHistogram(FINALIZE);
SamplePDBMovies(FINALIZE);
SampleDCTSTConfigurationFiles(FINALIZE);
SampleFreeEnergyProfile(FINALIZE);
SampleCationAndAdsorptionSites(FINALIZE);

```

Each of the sampling routine (in 'src/sample.c') has 5 scaling options:

- ALLOCATE to allocate memory needed for the sampling.
- INITIALIZE to initialized the routine if needed.
- SAMPLE to sample the properties.
- PRINT to periodically write the output to file.
- FINALIZE to free the requested memory and clean up.

Adding your own sampling routines requires an additional routine in 'src/sample.c', the definition in 'src/sample.h' and addition to calls to 'src/monte\_carlo.c'.

### 5.4.2 Molecular Dynamics

A molecular dynamics simulation is performed in several steps:

- The proper amount of molecules are created and they are inserted as as no overlaps occurred with the framework or other particles.

- Initialization: during the initialization period an NVT Monte-Carlo (MC) simulation is performed to rapidly achieve an equilibrium molecular arrangement.
- After the initialization period, velocities are assigned, drawn from the Maxwell-Boltzmann distribution at the desired average temperature to all the atoms. The total momentum of the system can be set to zero.
- Equilibration: Next, the system is further equilibrated by performing an NVT MD simulation using a specified ensemble.
- Production run: the simulation is performed in the requested ensemble and properties are measured.

The amount of cycles for each of these steps can be specified. For example, when starting from a restart-file there is no need for the Monte Carlo initialization, and if also the velocities are used from the restart-file then also the MD equilibration could be skipped. Moreover, the equilibration can be done in a different ensemble as the production run. This is most useful for NVE simulations, where the equilibration could be done using NVT. The final temperature of the NVE production run is then quite close the desired temperature (in NVE the temperature is not imposed).

The initialization part is not shown here, as it is very similar to regular Monte Carlo. The basic outline for the equilibration and production run are listed below. The most important lines are the 'Integration();' ones, which evolve the system a single time step. This routine is implemented in 'src/integration.c' and makes use of 'src/thermo\_baro\_stats.c' for temperature and pressure control.

```

// initialize
InitializeEnergiesAllSystems();
InitializeSmallMCStatisticsAllSystems();
InitializeMCMovesStatisticsAllSystems();

// compute initial energy
InitializeNoseHooverAllSystems();
InitializeForcesAllSystems();

// set the current ensemble to the initialization ensemble
for(i=0;i<NumberOfSystems;i++)
    Ensemble[i]=InitEnsemble[i];

InitializeEnergyAveragesAllSystems();

for(CurrentSystem=0;CurrentSystem<NumberOfSystems;CurrentSystem++)
{
    ReferenceEnergy[CurrentSystem]=ConservedEnergy[CurrentSystem];
    Drift[CurrentSystem]=0.0;
}

// Molecular-Dynamics initializing period to achieve a rapid equilibration of the velocities
for(CurrentCycle=0;CurrentCycle<NumberOfEquilibrationCycles;CurrentCycle++)
{
    for(CurrentSystem=0;CurrentSystem<NumberOfSystems;CurrentSystem++)
    {
        // regularly output system status and restart files
        if(CurrentCycle%PrintEvery==0)
        {
            PrintIntervalStatusEquilibration(CurrentCycle,NumberOfEquilibrationCycles,OutputFilePtr[CurrentSystem]);
            PrintRestartFile();
        }

        // evolve the system a full time-step
        Integration();

        // update the current energy-drift
        Drift[CurrentSystem]+=fabs((ConservedEnergy[CurrentSystem]-ReferenceEnergy[CurrentSystem])/
            ReferenceEnergy[CurrentSystem]);
    }
}

```

```

        }

    }

    // initialize sampling-routines at the start of the production run
    for( CurrentSystem=0; CurrentSystem<NumberOfSystems; CurrentSystem++)
    {
        Ensemble[CurrentSystem]=RunEnsemble[CurrentSystem];

        ReferenceEnergy[CurrentSystem]=ConservedEnergy[CurrentSystem];
        Drift[CurrentSystem]=0.0;
    }
    SampleInfraRedSpectra(INITIALIZE);
    SampleEndToEndDistanceHistogram(INITIALIZE);
    SampleMeanSquareDisplacementOrderN(INITIALIZE);
    SampleOnsagerMeanSquareDisplacementOrderN(INITIALIZE);
    SampleEnergyHistogram(INITIALIZE);
    SamplePositionHistogram(INITIALIZE);
    SampleRadialDistributionFunction(INITIALIZE);
    SamplePositionHistogram(INITIALIZE);
    SampleMoleculePropertyHistogram(INITIALIZE);
    SamplePDBMovies(INITIALIZE);
    SampleCationAndAdsorptionSites(INITIALIZE);

    // Molecular-Dynamics production run
    // loop over the amount of production cycles (MD integration steps)
    for( CurrentCycle=0; CurrentCycle<NumberOfCycles; CurrentCycle++)
    {
        // loop over all the systems and handle one by one
        for( CurrentSystem=0; CurrentSystem<NumberOfSystems; CurrentSystem++)
        {
            SampleInfraRedSpectra(SAMPLE);
            SampleEndToEndDistanceHistogram(SAMPLE);
            SampleMeanSquareDisplacementOrderN(SAMPLE);
            SampleOnsagerMeanSquareDisplacementOrderN(SAMPLE);
            SampleEnergyHistogram(SAMPLE);
            SamplePositionHistogram(SAMPLE);
            SampleRadialDistributionFunction(SAMPLE);
            SamplePositionHistogram(SAMPLE);
            SampleMoleculePropertyHistogram(SAMPLE);
            SampleCationAndAdsorptionSites(SAMPLE);

            // update all the average energies
            UpdateEnergyAveragesCurrentSystem();

            if( CurrentCycle%PrintPropertiesEvery==0)
                PrintPropertyStatus(CurrentCycle,NumberOfCycles,OutputFilePtr[CurrentSystem]);

            if( CurrentCycle%PrintEvery==0)
            {
                PrintIntervalStatus(CurrentCycle,NumberOfCycles,OutputFilePtr[CurrentSystem]);
                PrintRestartFile();
            }

            // regulary output radial distribution function
            SampleInfraRedSpectra(PRINT);
            SampleEndToEndDistanceHistogram(PRINT);
            SampleMeanSquareDisplacementOrderN(PRINT);
            SampleOnsagerMeanSquareDisplacementOrderN(PRINT);
            SampleEnergyHistogram(PRINT);
            SamplePositionHistogram(PRINT);
            SampleRadialDistributionFunction(PRINT);
            SamplePositionHistogram(PRINT);
            SampleMoleculePropertyHistogram(PRINT);
            SamplePDBMovies(PRINT);
        }
    }
}

```

```

    SampleCationAndAdsorptionSites(PRINT);

    // evolve the current system a full time step
    Integration();

    // update the current energy-drift
    Drift[CurrentSystem] += fabs((ConservedEnergy[CurrentSystem]-ReferenceEnergy[CurrentSystem])/
        ReferenceEnergy[CurrentSystem]);
}

}

// finalize and clean up
for(CurrentSystem=0;CurrentSystem<NumberOfSystems;CurrentSystem++)
{
    SampleInfraRedSpectra(FINALIZE);
    SampleEndToEndDistanceHistogram(FINALIZE);
    SampleMeanSquareDisplacementOrderN(FINALIZE);
    SampleOnsagerMeanSquareDisplacementOrderN(FINALIZE);
    SampleEnergyHistogram(FINALIZE);
    SamplePositionHistogram(FINALIZE);
    SampleRadialDistributionFunction(FINALIZE);
    SamplePositionHistogram(FINALIZE);
    SampleMoleculePropertyHistogram(FINALIZE);
    SamplePDBMovies(FINALIZE);
    SampleCationAndAdsorptionSites(FINALIZE);
}

```

Adding your own sampling routines requires an additional routine in 'src/sample.c', the definition in 'src/sample.h' and addition to calls to 'src/molecular\_dynamics.c'.

## 5.5 Debugging

### 5.5.1 Linux

There are several debuggers like 'gdb', and memory check utilities available, i.e. valgrind.

### 5.5.2 Mac OSX

Debugging memory error under Max OsX is easy. One can replace the standard library to allocate memory by different ones that check memory allocation and use. It can catch a lot of array out-of-bound error, even for dynamically allocated memory. See

```
man libgmalloc
```

An example, export 'RASPA\_DIR' to the installation directory, start the debugger, load the debugging libraries and start running the code.

```

export RASPA_DIR=${HOME}/RASPA/simulations/
gdb ~/RASPA/simulations/bin/simulate

GNU gdb 6.3.50-20050815 (Apple version gdb-768) (Tue Oct  2 04:07:49 UTC 2007)
Copyright 2004 Free Software Foundation, Inc.
GDB is free software, covered by the GNU General Public License, and you are
welcome to change it and/or distribute copies of it under certain conditions.
Type "show copying" to see the conditions.
There is absolutely no warranty for GDB. Type "show warranty" for details.
This GDB was configured as "i386-apple-darwin"...Reading symbols for shared libraries ... done

```

```
(gdb) set env DYLD_INSERT_LIBRARIES /usr/lib/libgmalloc.dylib
(gdb) r
```

## 5.6 RASPA as a project in Xcode 4

### 5.6.1 Getting started




Figure 5.1: Xcode




Figure 5.2: Xcode




Figure 5.3: Xcode




Figure 5.4: Xcode




Figure 5.5: Xcode




Figure 5.6: Xcode




Figure 5.7: Xcode




Figure 5.8: Xcode




Figure 5.9: Xcode




Figure 5.10: Xcode

## 5.6.2 Fine tuning

Up to now, we can build the project using 'make'. However, Sometimes we would like to build using 'make install'. First, we need to add a second target




Figure 5.11: Xcode

Change "RASPA copy" to "RASPA install"




Figure 5.12: Xcode




Figure 5.13: Xcode




Figure 5.14: Xcode

### 5.6.3 Using 'git' (version control)




Figure 5.15: Xcode




Figure 5.16: Xcode




Figure 5.17: Xcode

## 5.7 Using 'git'

### 5.7.1 Installing git

```
if [ -f /usr/local/git/contrib/completion/git-completion.bash ]; then
    . /usr/local/git/contrib/completion/git-completion.bash
```

```
fi

export PS1='h:\W$(__git_ps1 "(%s)") \u\$ '
```

will give you something like host:git-repository (master) user\$ when cd-ing into the git-repository.

### 5.7.2

remote:

```
mkdir RASPA.git
cd RASPA.git
git --bare init
```

local:

```
rm -rf .git
git init
git config --global user.name "David Dubbeldam"
git config --global user.email D.Dubbeldam@uva.nl
git remote rm RASPA-remote
git config branch.master.remote RASPA-remote && git config branch.master.merge refs/heads/master
git remote add RASPA-remote ssh://raspa@itsaio.science.uva.nl/~raspa/RASPA.git
git add .
git commit -m "Initial commit RASPA-1.0"
git tag v0.99-480
git push RASPA-remote master

git clone ssh://raspa@itsaio.science.uva.nl/~raspa/RASPA.git
cd RASPA
aclocal
autoconf
automake --add-missing
./scripts/make-gcc-local
make
make install
```

See the version with

```
git tag
```

```
git pull
```

Xcode

```
name: RASPA-remote
ssh://itsaio.science.uva.nl/~raspa/RASPA.git
```

## 5.8 Coding conventions

- Local variables are e.g. 'nr\_of\_atoms', global variables are named 'NumberOfAtoms'

## 5.9 Optimization tips

- Part of the clarity is making hunks of code into functions when appropriate. The cost of a function call is extremely small on modern machines, so optimization is NOT a valid excuse for writing ten-page functions.
- Expressions that are not dependent of a loop should be moved outside the loop
- One dimensional array are faster than multi-dimensional arrays
- Avoid referring to global or static variables inside the tightest loops. Global variables are hard to optimize, first copy these to local variables.
- If you already have an array of structs, but find that the critical part of your program is accessing only a small number of fields in each struct, you can split these fields into a separate array so that the unused fields do not get read into the cache unnecessarily.

# Chapter 6

## Troubleshooting

### The numerical value computed from finite differences is not equal to the analytical expression

Using 'SimulationType Numerical' the analytical expression for the force, stress etc. are compared to numerical values from finite differences. If just one (or at the most a few values) are different, then this might be an artifact arising from a finite cutoff in the Van der Waals potential. This can be checked by changing the value of the cutoff by about  $10^{-3}$  Å. This is a very small change, but larger than the displacements used in the finite differences. The problem is that for a finite difference scheme like:

$$f'(x) = \frac{f(x - \Delta) - 8f(x - \frac{1}{2}\Delta) + 8f(x + \frac{1}{2}\Delta) - f(x + \Delta)}{6\Delta} \quad (6.1)$$

it is possible that one of the displacements  $\Delta$  places the particle outside of the cutoff, while the original position was inside (or visa versa). For a force-shifted Van der Waals potential there is no problem, but for shifted potentials, or potentials with a simple truncation, the divergence becomes a problem.

**Excess loading is negative** This usually happens when computing an isotherm and the next pressure is above the vapor pressure. The boundary from gas to liquid adsorption has been crossed and the amount of excess molecules increases by orders of magnitude. There is a reason why experimental gas-phase isotherms are of finite range, they usually stop at the vapor pressure. Also, if the pressure is *very* high the fluid outside the crystal is compressed more and more while the loading inside the crystal remains the same (at maximum loading). Hence, excess adsorption eventually becomes negative.

**Large drift in Monte Carlo energies** This should *not* happen and signals an error in (one of) the Monte Carlo routines. During the Monte Carlo simulations, the running-energies are stored. These are starting energy, and all the added energy *differences*. At the final stage, the energy is recomputed again, and these should match within an error of about  $10^{-5}$  or lower. If you have added your own MC move, check whether you have properly added the energy differences to the running energies.

**Energy is not conserved in molecular dynamics** Usually, this happens because the time step is too large. Also, at initialization, the system can be far from equilibrated and a smaller time step is needed.

**RASPA "hangs" at initialization** Put 'CreateNumberOfMolecules 0' and check if that solves the problem. If so, then you have tried to add too many molecules in the system (i.e. more than actually fit in the system). For systems without a framework, one can also increase the size of the box.

**Segmentation fault** A memory access that is not allowed has occurred. This could happen when the input is incorrect. For example, if it is listed that there are 4 bonds, but you put in 5 lines, then all bonds and what follows next will be read in wrong. This is the most common cause of segmentation faults.

**Mean-square displacement is not linear** There are several known causes:

- Your system is one-dimensional and particles are unable to pass each other. This is known as 'single-file-diffusion' and the mean square displacement is proportional to the square root of time,
- You did not simulate long enough. In some systems it can take up to several nanoseconds before the msd becomes linear in time,
- You forgot to specify interactions between the molecules and they are not interacting.

**Minimization does not converge** A likely cause is that you minimize a system that would like to change angles, but you do not allow it to. In such a system, there is a non-vanishing stress. Try to minimize using NPT-PR with cell type 'Regular' or 'RegularUppertriangle'. Another reason could be that the electrostatics are not computed accurate enough. Increase the precision to 1e-10, using 'EwaldPrecision 1e-10'.

**Output is not written to file** Check with 'df -k' whether the disk is full.

**Molecule can not be grown** Check if the connectivity, i.e. the bonds, are correct.

**Framework flies apart** Check bonds for the framework and whether electrostatics and intra framework Van der Waals interactions are computed.

**Energy during molecular dynamics with a flexible framework is not well conserved** In zeolites, a common problem is that the angle of a Si-O-Si bend can become 180 degrees. This leads to an undefined torsion angle. If this occurs, try to use a smoothing function that slowly switches off the energy and force contributions for these 3 atoms as the angle approaches 180 degrees. See Bend/Torsion cross potentials.

**Amount of detected bonds/bends/torsions etc. for a flexible framework is wrong** For the detection of intra-framework potentials a connectivity table is made, where two atoms are considered bonded when their length is smaller than  $0.56 + r_i + r_j$ , where  $r_i$  and  $r_j$  are the covalent radii for the two atoms. The radii are specified in 'pseudo\_atoms.def'. The most likely cause is a wrong value for the radius. Note that even when starting from a restart-file, the connectivity table is based on the crystal structure.

**Oxgens connected to aluminum type 'O' are not automatically converted to 'Oa'** Use the option

```
ModifyOxgensConnectedToAluminium yes
```

**Strange behaviour when using cations** The problem could be related to CBMC of net-charged molecules. The Rosenbluth weights can become very large or very small, because the energy difference when displacing an ion is large. This can lead to numerical problems for ratios of combination of small/large, for example in the reinsertion move. To see whether this is the cause use 'RandomTranslationProbability' and set 'ReinsertionProbability' to zero.

**Minimization does not converge** Minimization code and algorithms are complex. Due to the harmonic approximation the jumps through the energy landscape can not be too large. A possible remedy therefore is to decrease the maximum step-length (default 0.3) using

```
MaximumStepLength 0.1
```

Another issue is the rotational degrees of freedom of the system. For periodic systems the system is invariant with respect to translation but not with respect to rotation, i.e. the energy changes for rotation of the whole system. In contrast, a molecule in a finite system is invariant with respect to both translation and rotation. However, for periodic systems at low loading with molecules without charges, groups/clusters of molecules can occur that do not have interactions with their images. In effect this has reduced the periodic system to a non-periodic one. If this occurs one can remove the system rotation explicitly with

```
RemoveRotationFromHessian yes
```

Note that this option should *not* be used on a truly periodic system. In addition to these algorithm settings, it is possible that the definition of the molecule and/or framework contains errors.

**Parallel tempering does not work for systems with cations** The problem is physical, it is just that all the energy distributions are more 'spiked' and overlap between the energies of the systems is rare. The only solution is to use more systems (and smaller temperature differences) to increase the acceptance rates of swapped between neighboring system. The same problem happens when one increases the system size.

**Results do not match data from literature** Common reasons include difference in simulation length, system size, cutoff value, tail-corrections vs shifted potentials, and handling of electrostatics. For adsorption, is the crystal structure that you used the same? Another very often made error, is comparing against different units, or an error in the conversion of units.

**Error during compiling** The 'gcc' and 'icc' compilers are tested. These compilers have C extensions that other C compilers could potentially lack.

**Error during linking** Make sure that the 'blas' and 'lapack' libraries are installed on your system.

**The results are different on different machines** We have come across one compiler-error: gcc 4.3.0 using optimizations "-O3" and "-O4" generated wrong code. Gcc 4.3.2 has resolved this bug.

**The program crashes with a 'segmentation fault'** Usually caused by wrong input-files, for example supplying 3 arguments to a torsion when 4 are expected. This causes all input to have strange values. To identify the cause make sure the job will use the same random number sequence are written in the output.

```
RandomSeed [int]
```

This will generate exactly the same sequence of events. Make sure the program is allowed to 'dump a core' (See the unix 'ulimit' command). Also, the executable needs to be compiled with the '-g' option which includes debugger information into the executable. Now restart the program and when it crashes, it will write a 'core-dump-file'. Start the debugger using a command similar to

```
gdb ~/RASPA/simulate/bin/simulate core
```

and type 'where' to obtain the line where the code crashed. It is also possible to just run the code inside the debugger.

## Acknowledgements

We would like to thank the following people for their help and input to improve the program, and for the very helpful discussions about the algorithms: Youn-Sang Bae, Tom Caremans, Juan Manuel Castillo Sanchez, Allison Dickey, Tina Düren, Gloria Emberger, Titus van Erp, Denise Ford, Houston Frost, Elena García-Pérez, Almudena Garcia Sanchez, Rachel Getman, Lev Sarkisov, Juan Jose Gutierrez Sevillano, John J. Low, Patrick Merkling, Patrick Ryan, Sudeep Punnathanam, Krista Walton, Chris Wilmer, Ozgur Yazaydin, and Decai Yu. Special thanks to Thijs Vlugt.