
MONASH University

ETC3550/ETC5550 Applied forecasting

Week 7: Seasonal exponential smoothing



ETS models

General notation E T S : **E**xponen**T**ial **S**moothing



Error **T**rend **S**eason

Error: Additive (`"A"`) or multiplicative (`"M"`)

ETS models

General notation E T S : **E**xponen**T**ial **S**moothing



Error **T**rend **S**eason

Error: Additive (`"A"`) or multiplicative (`"M"`)

Trend: None (`"N"`), additive (`"A"`), or damped (`"Ad"`).

ETS models

General notation E T S : **E**xponen**T**ial **S**moothing



Error **T**rend **S**eason

Error: Additive (`"A"`) or multiplicative (`"M"`)

Trend: None (`"N"`), additive (`"A"`), or damped (`"Ad"`).

Seasonality: None (`"N"`), additive (`"A"`) or multiplicative (`"M"`)

ETS(A,N,A): No trend, additive seasonal model

State equations	$\ell_t = \ell_{t-1} + \alpha\varepsilon_t$
	$s_t = s_{t-m} + \gamma\varepsilon_t$
Observation equation	$y_t = \ell_{t-1} + s_{t-m} + \varepsilon_t$
Forecast equation	$\hat{y}_{t+h\|t} = \ell_t + s_{t+h-m(k+1)}$

- Forecast errors: $\varepsilon_t = y_t - \hat{y}_{t|t-1} \sim \text{NID}(0, \sigma^2)$
- k is integer part of $(h-1)/m$.

ETS(A,A,A): Holt-Winters additive method

State equations	$\ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t$
	$b_t = b_{t-1} + \beta\varepsilon_t$
	$s_t = s_{t-m} + \gamma\varepsilon_t$
Observation equation	$y_t = \ell_{t-1} + b_{t-1} + s_{t-m} + \varepsilon_t$
Forecast equation	$\hat{y}_{t+h\|t} = \ell_t + hb_t + s_{t+h-m(k+1)}$

- Forecast errors: $\varepsilon_t = y_t - \hat{y}_{t|t-1} \sim \text{NID}(0, \sigma^2)$
- k is integer part of $(h-1)/m$.

ETS(M,A,M): Holt-Winters multiplicative method

State equations	$\ell_t = (\ell_{t-1} + b_{t-1})(1 + \alpha\varepsilon_t)$
	$b_t = b_{t-1} + \beta(\ell_{t-1} + b_{t-1})\varepsilon_t$
	$s_t = s_{t-m}(1 + \gamma\varepsilon_t)$
Observation equation	$y_t = (\ell_{t-1} + b_{t-1})s_{t-m}(1 + \varepsilon_t)$
Forecast equation	$\hat{y}_{t+h\|t} = (\ell_t + hb_t)s_{t+h-m(k+1)}$

- Relative forecast errors: $\varepsilon_t = \frac{y_t - \hat{y}_{t|t-1}}{\hat{y}_{t|t-1}} \sim \text{NID}(0, \sigma^2)$
- k is integer part of $(h-1)/m$.

All ETS models

Additive Error

Trend Component		Seasonal Component: N (None)	A (Additive)	M (Multiplicative)
N	(None)	A,N,N	A,N,A	A,N,M
A	(Additive)	A,A,N	A,A,A	A,A,M
A_d	(Additive damped)	A,A_d,N	A,A_d,A	A,A_d,M

All ETS models

Additive Error

Trend Component		Seasonal Component: N (None)	A (Additive)	M (Multiplicative)
N	(None)	A,N,N	A,N,A	A,N,M
A	(Additive)	A,A,N	A,A,A	A,A,M
A_d	(Additive damped)	A,A_d,N	A,A_d,A	A,A_d,M

Multiplicative Error

Trend Component		Seasonal Component: N (None)	A (Additive)	M (Multiplicative)
N	(None)	M,N,N	M,N,A	M,N,M
A	(Additive)	M,A,N	M,A,A	M,A,M
A_d	(Additive damped)	M,A_d,N	M,A_d,A	M,A_d,M

Additive error models

Trend	Seasonal		
	N	A	M
N	$y_t = \ell_{t-1} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \alpha\varepsilon_t$	$y_t = \ell_{t-1} + s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \alpha\varepsilon_t$ $s_t = s_{t-m} + \gamma\varepsilon_t$	$y_t = \ell_{t-1}s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \alpha\varepsilon_t/s_{t-m}$ $s_t = s_{t-m} + \gamma\varepsilon_t/\ell_{t-1}$
A	$y_t = \ell_{t-1} + b_{t-1} + \varepsilon_t$ $\ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t$ $b_t = b_{t-1} + \beta\varepsilon_t$	$y_t = \ell_{t-1} + b_{t-1} + s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t$ $b_t = b_{t-1} + \beta\varepsilon_t$ $s_t = s_{t-m} + \gamma\varepsilon_t$	$y_t = (\ell_{t-1} + b_{t-1})s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + b_{t-1} + \alpha\varepsilon_t/s_{t-m}$ $b_t = b_{t-1} + \beta\varepsilon_t/s_{t-m}$ $s_t = s_{t-m} + \gamma\varepsilon_t/(\ell_{t-1} + b_{t-1})$
A_d	$y_t = \ell_{t-1} + \phi b_{t-1} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha\varepsilon_t$ $b_t = \phi b_{t-1} + \beta\varepsilon_t$	$y_t = \ell_{t-1} + \phi b_{t-1} + s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha\varepsilon_t$ $b_t = \phi b_{t-1} + \beta\varepsilon_t$ $s_t = s_{t-m} + \gamma\varepsilon_t$	$y_t = (\ell_{t-1} + \phi b_{t-1})s_{t-m} + \varepsilon_t$ $\ell_t = \ell_{t-1} + \phi b_{t-1} + \alpha\varepsilon_t/s_{t-m}$ $b_t = \phi b_{t-1} + \beta\varepsilon_t/s_{t-m}$ $s_t = s_{t-m} + \gamma\varepsilon_t/(\ell_{t-1} + \phi b_{t-1})$

Multiplicative error models

Trend	Seasonal		
	N	**A**	**M**
N	$y_t = \ell_{t-1}(1+\varepsilon_t)$ $\ell_t = \ell_{t-1}(1+\alpha\varepsilon_t)$	$y_t = (\ell_{t-1}+s_{t-m})(1+\varepsilon_t)$ $\ell_t = \ell_{t-1}+\alpha(\ell_{t-1}+s_{t-m})\varepsilon_t$ $s_t = s_{t-m}+\gamma(\ell_{t-1}+s_{t-m})\varepsilon_t$	$y_t = \ell_{t-1}s_{t-m}(1+\varepsilon_t)$ $\ell_t = \ell_{t-1}(1+\alpha\varepsilon_t)$ $s_t = s_{t-m}(1+\gamma\varepsilon_t)$
A	$y_t = (\ell_{t-1}+b_{t-1})(1+\varepsilon_t)$ $\ell_t = (\ell_{t-1}+b_{t-1})(1+\alpha\varepsilon_t)$ $b_t = b_{t-1}+\beta(\ell_{t-1}+b_{t-1})\varepsilon_t$	$y_t = (\ell_{t-1}+b_{t-1}+s_{t-m})(1+\varepsilon_t)$ $\ell_t = \ell_{t-1}+b_{t-1}+\alpha(\ell_{t-1}+b_{t-1}+s_{t-m})\varepsilon_t$ $b_t = b_{t-1}+\beta(\ell_{t-1}+b_{t-1}+s_{t-m})\varepsilon_t$ $s_t = s_{t-m}+\gamma(\ell_{t-1}+b_{t-1}+s_{t-m})\varepsilon_t$	$y_t = (\ell_{t-1}+b_{t-1})s_{t-m}(1+\varepsilon_t)$ $\ell_t = (\ell_{t-1}+b_{t-1})(1+\alpha\varepsilon_t)$ $b_t = b_{t-1}+\beta(\ell_{t-1}+b_{t-1})\varepsilon_t$ $s_t = s_{t-m}(1+\gamma\varepsilon_t)$
$\mathbf{A_d}$	$y_t = (\ell_{t-1}+\phi b_{t-1})(1+\varepsilon_t)$ $\ell_t = (\ell_{t-1}+\phi b_{t-1})(1+\alpha\varepsilon_t)$ $b_t = \phi b_{t-1}+\beta(\ell_{t-1}+\phi b_{t-1})\varepsilon_t$	$y_t = (\ell_{t-1}+\phi b_{t-1}+s_{t-m})(1+\varepsilon_t)$ $\ell_t = \ell_{t-1}+\phi b_{t-1}+\alpha(\ell_{t-1}+\phi b_{t-1}+s_{t-m})\varepsilon_t$ $b_t = \phi b_{t-1}+\beta(\ell_{t-1}+\phi b_{t-1}+s_{t-m})\varepsilon_t$ $s_t = s_{t-m}+\gamma(\ell_{t-1}+\phi b_{t-1}+s_{t-m})\varepsilon_t$	$y_t = (\ell_{t-1}+\phi b_{t-1})s_{t-m}(1+\varepsilon_t)$ $\ell_t = (\ell_{t-1}+\phi b_{t-1})(1+\alpha\varepsilon_t)$ $b_t = \phi b_{t-1}+\beta(\ell_{t-1}+\phi b_{t-1})\varepsilon_t$ $s_t = s_{t-m}(1+\gamma\varepsilon_t)$

Estimating ETS models

- Smoothing parameters α, β, γ and ϕ, and the initial states ℓ_0, b_0, $s_0, s_{-1}, \ldots, s_{-m+1}$ are estimated by maximising the "likelihood" = the probability of the data arising from the specified model.
- For models with additive errors equivalent to minimising SSE.
- For models with multiplicative errors, **not** equivalent to minimising SSE.

Innovations state space models

Let $\boldsymbol{x}_t = (\ell_t, b_t, s_t, s_{t-1}, \dots, s_{t-m+1})$ and $\varepsilon_t \overset{\text{iid}}{\sim} \text{N}(0, \sigma^2)$.

$$
\begin{aligned}
y_t &= \underbrace{h(\boldsymbol{x}_{t-1})}_{\mu_t} + \underbrace{k(\boldsymbol{x}_{t-1})\varepsilon_t}_{e_t} \\
\boldsymbol{x}_t &= f(\boldsymbol{x}_{t-1}) + g(\boldsymbol{x}_{t-1})\varepsilon_t
\end{aligned}
$$

Additive errors

$k(\boldsymbol{x}_{t-1}) = 1.$ $\quad y_t = \mu_t + \varepsilon_t.$

Multiplicative errors

$k(\boldsymbol{x}_{t-1}) = \mu_t.$ $\quad y_t = \mu_t(1 + \varepsilon_t).$

$\varepsilon_t = (y_t - \mu_t)/\mu_t$ is relative error.

Innovations state space models

Estimation

$$L^*(\boldsymbol{\theta}, \boldsymbol{x}_0) = T\log\left(\sum_{t=1}^{T}\varepsilon_t^2\right) + 2\sum_{t=1}^{T}\log|k(\boldsymbol{x}_{t-1})|$$
$$= -2\log(\text{Likelihood}) + \text{constant}$$

- Estimate parameters $\boldsymbol{\theta} = (\alpha, \beta, \gamma, \phi)$ and initial states $\boldsymbol{x}_0 = (\ell_0, b_0, s_0, s_{-1}, \ldots, s_{-m+1})$ by minimizing L^*.

Parameter restrictions

Traditional region

- $0 < \alpha < 1$
- $0 < \beta < \alpha$
- $0 < \gamma < 1 - \alpha$
- $0.8 < \phi < 0.98$ — to prevent numerical difficulties.

Admissible region

- To prevent observations in the distant past having a continuing effect on current forecasts
- Usually (but not always) less restrictive than *traditional* region
- e.g., ETS(A,N,N): *traditional* $0 < \alpha < 1$
  admissible $0 < \alpha < 2$

Parameter restrictions

fable default:
intersection of both regions

Traditional region

- $0 < \alpha < 1$
- $0 < \beta < \alpha$
- $0 < \gamma < 1 - \alpha$
- $0.8 < \phi < 0.98$ — to prevent numerical difficulties.

Admissible region

- To prevent observations in the distant past having a continuing effect on current forecasts
- Usually (but not always) less restrictive than *traditional* region
- e.g., ETS(A,N,N): *traditional* $0 < \alpha < 1$
  admissible $0 < \alpha < 2$

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2\log(\text{L}) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2\log(\text{L}) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Corrected AIC

$$\text{AIC}_\text{c} = \text{AIC} + \frac{2k(k+1)}{T-k-1}$$

which is the AIC corrected (for small sample bias).

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2\log(\text{L}) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Corrected AIC

$$\text{AIC}_\text{c} = \text{AIC} + \frac{2k(k+1)}{T-k-1}$$

which is the AIC corrected (for small sample bias).

Bayesian Information Criterion

$$\text{BIC} = \text{AIC} + k[\log(T) - 2].$$

AIC and cross-validation

Minimizing the AIC assuming Gaussian residuals is asymptotically equivalent to minimizing one-step time series cross validation MSE.

Automatic forecasting

From Hyndman et al. (IJF, 2002):

- Apply each model that is appropriate to the data. Optimize parameters and initial values using MLE (or some other criterion).
- Select best method using AICc:
- Produce forecasts using best method.
- Obtain forecast intervals using underlying state space model.

Method performed very well in M3 competition.

Some unstable models

- Some of the combinations of (Error, Trend, Seasonal) can lead to numerical difficulties; see equations with **division by a state**.
- These are: ETS(A,N,M), ETS(A,A,M), ETS(A,A_d,M).

ETS models

Additive Error

Trend Component		Seasonal Component: N (None)	A (Additive)	M (Multiplicative)
N	(None)	A,N,N	A,N,A	~~A,N,M~~
A	(Additive)	A,A,N	A,A,A	~~A,A,M~~
A_d	(Additive damped)	A,A_d,N	A,A_d,A	~~A,A_d,M~~

Multiplicative Error

Trend Component		Seasonal Component: N (None)	A (Additive)	M (Multiplicative)
N	(None)	M,N,N	M,N,A	M,N,M
A	(Additive)	M,A,N	M,A,A	M,A,M
A_d	(Additive damped)	M,A_d,N	M,A_d,A	M,A_d,M

Residuals

Response residuals

$$\hat{e}_t = y_t - \hat{y}_{t|t-1}$$

Innovation residuals

Additive error model:

$$\hat{\varepsilon}_t = y_t - \hat{y}_{t|t-1}$$

Multiplicative error model:

$$\hat{\varepsilon}_t = \frac{y_t - \hat{y}_{t|t-1}}{\hat{y}_{t|t-1}}$$