


ETC3550/ETC5550

Applied forecasting

Week 5: Accuracy evaluation



Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions
- 4 Forecast evaluation
- 5 Time series cross-validation

Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions
- 4 Forecast evaluation
- 5 Time series cross-validation

Workshop tomorrow

The workshop is online tomorrow.

Zoom link on Ed Discussion Forum

Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions
- 4 Forecast evaluation
- 5 Time series cross-validation

Fitted values

- $\hat{y}_{t|t-1}$ is the forecast of y_t based on observations $y_1, \dots, y_{t-1}$.
- We call these “fitted values”.
- Sometimes drop the subscript: $\hat{y}_t \equiv \hat{y}_{t|t-1}$.
- Often not true forecasts since parameters are estimated on all data.

For example:

- $\hat{y}_t = \bar{y}$ for average method.
- $\hat{y}_t = y_{t-1} + (y_T - y_1)/(T - 1)$ for drift method.

Forecasting residuals

Residuals in forecasting: difference between observed value and its fitted value: $e_t = y_t - \hat{y}_{t|t-1}$.

Forecasting residuals

Residuals in forecasting: difference between observed value and its fitted value: $e_t = y_t - \hat{y}_{t|t-1}$.

Assumptions

- 1 $\{e_t\}$ uncorrelated. If they aren't, then information left in residuals that should be used in computing forecasts.
- 2 $\{e_t\}$ have mean zero. If they don't, then forecasts are biased.

Forecasting residuals

Residuals in forecasting: difference between observed value and its fitted value: $e_t = y_t - \hat{y}_{t|t-1}$.

Assumptions

- 1 $\{e_t\}$ uncorrelated. If they aren't, then information left in residuals that should be used in computing forecasts.
- 2 $\{e_t\}$ have mean zero. If they don't, then forecasts are biased.

Useful properties (for distributions & prediction intervals)

- 3 $\{e_t\}$ have constant variance.
- 4 $\{e_t\}$ are normally distributed.

ACF of residuals

- We assume that the residuals are white noise (uncorrelated, mean zero, constant variance). If they aren't, then there is information left in the residuals that should be used in computing forecasts.
- So a standard residual diagnostic is to check the ACF of the residuals of a forecasting method.
- We *expect* these to look like white noise.

Ljung-Box tests

r_k = autocorrelation of residual at lag k

Test *whole set* of r_k values simultaneously.

$$Q^* = T(T + 2) \sum_{k=1}^{\ell} (T - k)^{-1} r_k^2$$

where ℓ is max lag being considered and T is number of observations.

- My preferences: $\ell = 10$ for non-seasonal data, $\ell = 2m$ for seasonal data (where m is seasonal period).
- If data are WN, Q^* has χ^2 distribution with ℓ degrees of freedom.

Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions**
- 4 Forecast evaluation
- 5 Time series cross-validation

Forecast distributions

- A forecast $\hat{y}_{T+h|T}$ is (usually) the mean of the conditional distribution $y_{T+h} \mid y_1, \dots, y_T$.
- Most time series models produce normally distributed forecasts.
- The forecast distribution describes the probability of observing any future value.

Forecast distributions

Assuming residuals are normal, uncorrelated, $\text{sd} = \hat{\sigma}$:

Mean: $y_{T+h|T} \sim N(\bar{y}, (1 + 1/T)\hat{\sigma}^2)$

Naïve: $y_{T+h|T} \sim N(y_T, h\hat{\sigma}^2)$

Seasonal naïve: $y_{T+h|T} \sim N(y_{T+h-m(k+1)}, (k+1)\hat{\sigma}^2)$

Drift: $y_{T+h|T} \sim N(y_T + \frac{h}{T-1}(y_T - y_1), h\frac{T+h}{T}\hat{\sigma}^2)$

where k is the integer part of $(h-1)/m$.

Note that when $h = 1$ and T is large, these all give the same approximate forecast variance: $\hat{\sigma}^2$.

Prediction intervals

- A prediction interval gives a region within which we expect y_{T+h} to lie with a specified probability.
- Assuming forecast errors are normally distributed, then a 95% PI is

$$\hat{y}_{T+h|T} \pm 1.96\hat{\sigma}_h$$

where $\hat{\sigma}_h$ is the st dev of the h -step distribution.

- When $h = 1$, $\hat{\sigma}_h$ can be estimated from the residuals.

Prediction intervals

- Point forecasts are often useless without a measure of uncertainty (such as prediction intervals).
- Prediction intervals require a stochastic model (with random errors, etc).
- For most models, prediction intervals get wider as the forecast horizon increases.
- Use level argument to control coverage.
- Check residual assumptions before believing them.
- Prediction intervals are usually too narrow due to unaccounted uncertainty.

Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions
- 4 Forecast evaluation**
- 5 Time series cross-validation

Training and test sets



- A model which fits the training data well will not necessarily forecast well.
- A perfect fit can always be obtained by using a model with enough parameters.
- Over-fitting a model to data is just as bad as failing to identify a systematic pattern in the data.
- The test set must not be used for *any* aspect of model development or calculation of forecasts.
- Forecast accuracy is based only on the test set.

Forecast errors

Forecast “error”: the difference between an observed value and its forecast.

$$e_{T+h} = y_{T+h} - \hat{y}_{T+h|T},$$

where the training data is given by $\{y_1, \dots, y_T\}$

- Unlike residuals, forecast errors on the test set involve multi-step forecasts.
- These are *true* forecast errors as the test data is not used in computing $\hat{y}_{T+h|T}$.

Measures of forecast accuracy

y_{T+h} = $(T + h)$ th observation, $h = 1, \dots, H$

$\hat{y}_{T+h|T}$ = its forecast based on data up to time T .

$$e_{T+h} = y_{T+h} - \hat{y}_{T+h|T}$$

$$\text{MAE} = \text{mean}(|e_{T+h}|)$$

$$\text{MSE} = \text{mean}(e_{T+h}^2)$$

$$\text{MAPE} = 100\text{mean}(|e_{T+h}|/|y_{T+h}|)$$

$$\text{RMSE} = \sqrt{\text{mean}(e_{T+h}^2)}$$

Measures of forecast accuracy

y_{T+h} = $(T + h)$ th observation, $h = 1, \dots, H$

$\hat{y}_{T+h|T}$ = its forecast based on data up to time T .

$$e_{T+h} = y_{T+h} - \hat{y}_{T+h|T}$$

$$\text{MAE} = \text{mean}(|e_{T+h}|)$$

$$\text{MSE} = \text{mean}(e_{T+h}^2)$$

$$\text{RMSE} = \sqrt{\text{mean}(e_{T+h}^2)}$$

$$\text{MAPE} = 100\text{mean}(|e_{T+h}|/|y_{T+h}|)$$

- MAE, MSE, RMSE are all scale dependent.
- MAPE is scale independent but is only sensible if $y_t \gg 0$ for all t , and y has a natural zero.

Scaled Errors

Proposed by Hyndman and Koehler (IJF, 2006).

- For non-seasonal time series, scale errors using naïve forecasts:

$$q_{T+h} = \frac{e_{T+h}}{\frac{1}{T-1} \sum_{t=2}^T |y_t - y_{t-1}|}.$$

Scaled Errors

Proposed by Hyndman and Koehler (IJF, 2006).

- For non-seasonal time series, scale errors using naïve forecasts:

$$q_{T+h} = \frac{e_{T+h}}{\frac{1}{T-1} \sum_{t=2}^T |y_t - y_{t-1}|}.$$

- For seasonal time series, scale forecast errors using seasonal naïve forecasts:

$$q_{T+h} = \frac{e_{T+h}}{\frac{1}{T-m} \sum_{t=m+1}^T |y_t - y_{t-m}|}.$$

Scaled errors

Mean Absolute Scaled Error

$$\text{MASE} = \text{mean}(|q_{T+h}|)$$

Scaled errors

Mean Absolute Scaled Error

$$\text{MASE} = \text{mean}(|q_{T+h}|)$$

Root Mean Squared Scaled Error

$$\text{RMSSE} = \sqrt{\text{mean}(q_{T+h}^2)}$$

where

$$q_{T+h}^2 = \frac{e_{T+h}^2}{\frac{1}{T-m} \sum_{t=m+1}^T (y_t - y_{t-m})^2},$$


and we set $m = 1$ for non-seasonal data.

Outline

- 1 Workshop tomorrow
- 2 Residuals
- 3 Forecast distributions
- 4 Forecast evaluation
- 5 Time series cross-validation


Time series cross-validation

Traditional evaluation




Time series cross-validation

Traditional evaluation




Time series cross-validation




Time series cross-validation

Traditional evaluation




Time series cross-validation




Time series cross-validation

Traditional evaluation




Time series cross-validation




Time series cross-validation

Traditional evaluation




Time series cross-validation




Time series cross-validation

Traditional evaluation



Time series cross-validation



Workshop tomorrow

The workshop is online tomorrow.

Zoom link on Ed Discussion Forum