

ETF3231/5231

Business forecasting

Week 11: Dynamic
Regression <https://bf.numbat.space/>



Outline

- 1 Regression with ARIMA errors
- 2 Dynamic harmonic regression
- 3 Stochastic and deterministic trends

Outline

- 1 Regression with ARIMA errors
- 2 Dynamic harmonic regression
- 3 Stochastic and deterministic trends

Regression models

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \varepsilon_t,$$

- y_t modeled as function of k explanatory variables $x_{1,t}, \dots, x_{k,t}$.
- In regression, we assume that ε_t is WN.
- Now we want to allow ε_t to be autocorrelated.

Regression with ARIMA errors

Regression models

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \varepsilon_t,$$

- y_t modeled as function of k explanatory variables $x_{1,t}, \dots, x_{k,t}$.
- In regression, we assume that ε_t is WN.
- Now we want to allow ε_t to be autocorrelated.

Example: ARIMA(1,1,1) errors

$$\begin{aligned} y_t &= \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t, \\ (1 - \phi_1 B)(1 - B)\eta_t &= (1 + \theta_1 B)\varepsilon_t, \end{aligned}$$

where ε_t is white noise.

Example: $\eta_t = \text{ARIMA}(1,1,1)$

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Example: $\eta_t = \text{ARIMA}(1,1,1)$

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

- Be careful in distinguishing η_t from ε_t .
- Only the errors ε_t are assumed to be white noise.
- In ordinary regression, η_t is assumed to be white noise and so $\eta_t = \varepsilon_t$.

If we minimize $\sum \eta_t^2$ (by using ordinary regression):

- 1 Estimated coefficients $\hat{\beta}_0, \dots, \hat{\beta}_k$ are no longer optimal as some information ignored;
- 2 Statistical tests associated with the model (e.g., t-tests on the coefficients) are incorrect.
- 3 p -values for coefficients usually too small (“spurious regression”).
- 4 AIC of fitted models misleading.

If we minimize $\sum \eta_t^2$ (by using ordinary regression):

- 1 Estimated coefficients $\hat{\beta}_0, \dots, \hat{\beta}_k$ are no longer optimal as some information ignored;
- 2 Statistical tests associated with the model (e.g., t-tests on the coefficients) are incorrect.
- 3 p -values for coefficients usually too small (“spurious regression”).
- 4 AIC of fitted models misleading.
 - Minimizing $\sum \varepsilon_t^2$ avoids these problems.
 - Maximizing likelihood similar to minimizing $\sum \varepsilon_t^2$.

Regression with ARMA errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$

where η_t is an ARMA process.

- All variables in the model **must be stationary**.
- If we estimate the model while any of these are non-stationary, the estimated coefficients **can be incorrect**.
- **Difference** variables until all stationary.
- If necessary, apply same differencing to all variables.

Model with ARIMA(1,1,1) errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Model with ARIMA(1,1,1) errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Equivalent to model with ARIMA(1,0,1) errors

$$y'_t = \beta_1 x'_{1,t} + \cdots + \beta_k x'_{k,t} + \eta'_t,$$
$$(1 - \phi_1 B)\eta'_t = (1 + \theta_1 B)\varepsilon_t,$$

where $y'_t = y_t - y_{t-1}$, $x'_{t,i} = x_{t,i} - x_{t-1,i}$ and $\eta'_t = \eta_t - \eta_{t-1}$.

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Original data

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t$$

where $\phi(B)(1 - B)^d \eta_t = \theta(B)\varepsilon_t$

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Original data

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t$$

where $\phi(B)(1 - B)^d \eta_t = \theta(B)\varepsilon_t$

After differencing all variables

$$y'_t = \beta_1 x'_{1,t} + \cdots + \beta_k x'_{k,t} + \eta'_t.$$

where $\phi(B)\eta'_t = \theta(B)\varepsilon_t$,

$$y'_t = (1 - B)^d y_t, \quad x'_{i,t} = (1 - B)^d x_{i,t}, \quad \text{and} \quad \eta'_t = (1 - B)^d \eta_t$$

Regression with ARIMA errors

- In R, we can specify an $\text{ARIMA}(p, d, q)$ for the errors, and d levels of differencing will be applied to all variables $(y, x_{1,t}, \dots, x_{k,t})$ during estimation.
- Check that ε_t series looks like white noise.
- AICc can be calculated for final model.
- Repeat procedure for all subsets of predictors to be considered, and select model with lowest AICc value.

- To forecast a regression model with ARIMA errors, we need to forecast the regression part of the model and the ARIMA part of the model and combine the results.
- Some predictors are known into the future (e.g., time, dummies).
- Separate forecasting models may be needed for other predictors.
- Forecast intervals ignore the uncertainty in forecasting the predictors.

Outline

- 1 Regression with ARIMA errors
- 2 Dynamic harmonic regression
- 3 Stochastic and deterministic trends

Dynamic harmonic regression

Combine Fourier terms with ARIMA errors

Advantages


- it allows any length seasonality;
- for data with more than one seasonal period, you can include Fourier terms of different frequencies;
- the seasonal pattern is smooth for small values of K (but more wiggly seasonality can be handled by increasing K);
- the short-term dynamics are easily handled with a simple ARMA error.

Disadvantages

- seasonality is assumed to be fixed

Eating-out expenditure

```
aus_cafe <- aus_retail |> filter(  
  Industry == "Cafes, restaurants and takeaway food services",  
  year(Month) %in% 2004:2018  
) |> summarise(Turnover = sum(Turnover))  
aus_cafe |> autoplot(Turnover)
```




Eating-out expenditure

```
fit <- aus_cafe |> model(
  `K = 1` = ARIMA(log(Turnover) ~ fourier(K = 1) + PDQ(0,0,0)),
  `K = 2` = ARIMA(log(Turnover) ~ fourier(K = 2) + PDQ(0,0,0)),
  `K = 3` = ARIMA(log(Turnover) ~ fourier(K = 3) + PDQ(0,0,0)),
  `K = 4` = ARIMA(log(Turnover) ~ fourier(K = 4) + PDQ(0,0,0)),
  `K = 5` = ARIMA(log(Turnover) ~ fourier(K = 5) + PDQ(0,0,0)),
  `K = 6` = ARIMA(log(Turnover) ~ fourier(K = 6) + PDQ(0,0,0)))


glance(fit) |> select(.model, sigma2, log_lik, AIC, AICc, BIC)
```

.model	sigma2	log_lik	AIC	AICc	BIC
K = 1	0.002	317	-616	-615	-588
K = 2	0.001	362	-700	-698	-661
K = 3	0.001	394	-763	-761	-725
K = 4	0.001	427	-822	-818	-771
K = 5	0.000	474	-919	-917	-875
K = 6	0.000	474	-920	-918	-875


Eating-out expenditure




Eating-out expenditure




Eating-out expenditure




Eating-out expenditure



Eating-out expenditure



Eating-out expenditure



Outline

- 1 Regression with ARIMA errors
- 2 Dynamic harmonic regression
- 3 Stochastic and deterministic trends

Stochastic & deterministic trends

Deterministic trend

$$y_t = \beta_0 + \beta_1 t + \eta_t$$

where η_t is ARMA process.

Stochastic & deterministic trends

Deterministic trend

$$y_t = \beta_0 + \beta_1 t + \eta_t$$

where η_t is ARMA process.

Stochastic trend

$$y_t = \beta_0 + \beta_1 t + \eta_t$$

where η_t is ARIMA process with $d = 1$.

Stochastic & deterministic trends

Deterministic trend

$$y_t = \beta_0 + \beta_1 t + \eta_t$$

where η_t is ARMA process.

Stochastic trend

$$y_t = \beta_0 + \beta_1 t + \eta_t$$

where η_t is ARIMA process with $d = 1$.


Difference both sides until η_t is stationary:

$$y'_t = \beta_1 + \eta'_t$$

where η'_t is ARMA process.

Air transport passengers Australia

```
aus_airpassengers |>  
  autoplot(Passengers) +  
  labs(y = "Passengers (millions)",  
       title = "Total air passengers")
```



Air transport passengers Australia

Deterministic trend

```
fit_deterministic <- aus_airpassengers |>  
  model(ARIMA(Passengers ~ 1 + trend() + pdq(d = 0)))  
report(fit_deterministic)
```

Series: Passengers

Model: LM w/ ARIMA(1,0,0) errors

Coefficients:

	ar1	trend()	intercept
	0.9564	1.415	0.901
s.e.	0.0362	0.197	7.075

sigma^2 estimated as 4.343: log likelihood=-101

AIC=210 AICc=211 BIC=217

Air transport passengers Australia

Deterministic trend

```
fit_deterministic <- aus_airpassengers |>
  model(ARIMA(Passengers ~ 1 + trend() + pdq(d = 0)))
report(fit_deterministic)
```

Series: Passengers

Model: LM w/ ARIMA(1,0,0) errors

Coefficients:

	ar1	trend()	intercept
	0.9564	1.415	0.901
s.e.	0.0362	0.197	7.075

sigma^2 estimated as 4.343: log likelihood=-101

AIC=210 AICc=211 BIC=217

$$y_t = 0.901 + 1.415t + \eta_t$$

$$\eta_t = 0.956\eta_{t-1} + \varepsilon_t$$

$$\varepsilon_t \sim \text{NID}(0, 4.343).$$

Air transport passengers Australia

Stochastic trend

```
fit_stochastic <- aus_airpassengers |>  
  model(ARIMA(Passengers ~ 1 + pdq(d = 1)))  
report(fit_stochastic)
```

Series: Passengers

Model: ARIMA(0,1,0) w/ drift

Coefficients:

constant

1.419

s.e. 0.301

sigma^2 estimated as 4.271: log likelihood=-98.2

AIC=200 AICc=201 BIC=204

Air transport passengers Australia

Stochastic trend

```
fit_stochastic <- aus_airpassengers |>  
  model(ARIMA(Passengers ~ 1 + pdq(d = 1)))  
report(fit_stochastic)
```

Series: Passengers

Model: ARIMA(0,1,0) w/ drift

Coefficients:

constant

1.419

s.e. 0.301

$$y_t - y_{t-1} = 1.419 + \varepsilon_t,$$

$$y_t = y_0 + 1.419t + \eta_t$$


$$\eta_t = \eta_{t-1} + \varepsilon_t$$

$$\varepsilon_t \sim \text{NID}(0, 4.271).$$

sigma^2 estimated as 4.271: log likelihood=-98.2


AIC=200 AICc=201 BIC=204

Air transport passengers Australia



Air transport passengers Australia

```
aus_airpassengers |> autoplot(Passengers) +  
  autolayer(fit_stochastic |> forecast(h = 20), colour = "#0072B2", level = 95) +  
  autolayer(fit_deterministic |> forecast(h = 20), colour = "#D55E00", level = 95,  
    alpha = 0.65) +  
  labs(y = "Air passengers (millions)", title = "Forecasts from trend models")
```



Forecasting with trend

- Point forecasts are almost identical, but prediction intervals differ.
- Stochastic trends have much wider prediction intervals because the errors are non-stationary.
- Be careful of forecasting with deterministic trends too far ahead.