

# SPSA on the Noisy Function Testbed

Steffen Finck  
Vorarlberg University of Applied Sciences  
Hochschulstrasse 1  
Dornbirn, Austria  
steffen.finck@fhv.at

## ABSTRACT

This paper benchmarks the Simultaneous Perturbation Stochastic Algorithm (SPSA) [4] on the BBOB 2009 noisy testbed. SPSA is a widely used optimization algorithm with its main application in noisy optimization. The paper presents briefly the algorithm and used parameter setting for the testbed.

## Categories and Subject Descriptors

G.1.6 [Numerical Analysis]: Optimization Global Optimization, Unconstrained Optimization; F.2.1 [Analysis of Algorithms and Problem Complexity]: Numerical Algorithms and Problems

## General Terms

Algorithms

## Keywords

Benchmarking, Black-box optimization, evolutionary computation, stochastic optimization

## 1. INTRODUCTION

The SPSA algorithm is a very common and widely used optimization algorithm [5] and primarily designed for noisy optimization. In this paper the basic variant with a simple multistart procedure is presented. The main feature of SPSA is the use of just 2 function evaluations to determine the gradient, independent of the search space dimension DIM. As shown in [4, 6] this is advantageous (especially for large DIM) compared with common stochastic approximation algorithm which use  $2 \times \text{DIM}$  function evaluations to approximate the gradient. The here presented algorithm is coupled with a simple restart procedure to effectively use the given number of maximal function evaluations, similar to Fig. 3 in [2].

## 2. ALGORITHM PRESENTATION

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

GECCO'09, July 8–12, 2009, Montréal Québec, Canada.  
Copyright 2009 ACM 978-1-60558-505-5/09/07 ...\$5.00.

In Fig. 1 the main algorithm is presented.

The gain `ak` is used for the update of the current search point, while the gain `ck` is used for the test step of the gradient approximation. The determination of their initial values is shown in Fig. 2. To improve the performance of SPSA, `lambda` gradient approximations are averaged within one iteration before the update of the current search point. Thus, the here presented SPSA uses  $2 \times \text{lambda}$  function evaluations per iteration.

To effectively use the allowed number of function evaluations a simple restart procedure was implemented. It is shown in Fig. 3. To prevent infinite runs the procedure terminates if the maximal number of restarts is reached.

## 3. EXPERIMENTAL PROCEDURE

The gain rates were set to their recommended values `a0` = 0.602 and `c0` = 0.101, instead of the respective optimal values. All other parameters were set as recommended in [6]. The experiments were conducted on a Cluster with 2.44 GHz CPUs (machine\_type x86\_64) under Octave 3.0.2.

## 4. RESULTS

Results from experiments according to [2] on the benchmarks functions given in [1, 3] are presented in Figures 4 and 5 and in Tables 1 and 2.

## 5. CPU TIMING EXPERIMENT

For the timing experiment the same multistart algorithm was run on  $f_8$  and restarted until at least 30 seconds had passed (according to Figure 2 in [2]). The results were 1.2; 1.2; 1.2; 1.2; 1.3 and  $1.4 \times 10^{-4}$  seconds per function evaluation in dimension 2; 3; 5; 10; 20 and 40, respectively. The dependency of CPU time on the search space dimensionality is small.

## 6. CONCLUSION

This paper reports the result for the basic SPSA on the BBOB 2009 noisy testbed.

## Acknowledgments

The author would like to acknowledge the great work of the BBOB team with particular kudos the Anne Auger, Nikolaus Hansen and Raymond Ros. This work was supported by the Austrian Science Fund (FWF) under grant P19069-N18.

Figure 1: SPSA in Matlab

---

```
% simple spsa function
function [x,termvalue] = alg(FUN, x, parameter, maxGenerations, ftarget,...
                             DIM, maxfunevals)

% intialze counters
k = 1;

% initialize algorithm parameter
a0 = parameter(1);
alpha = parameter(2);
c0 = parameter(3);
gamma = parameter(4);
A = parameter(5);
lambda = parameter(6);

while 1

    % gain sequences ak and ck
    ak = a0 * (A + k)^(-alpha);
    ck = c0 * k^(-gamma);

    % gradient approximation with averaging of several approximations
    delta = 2*round(rand(DIM,lambda))-1;
    X = repmat(x,1,lambda);
    yplus = FUN(X + ck.*delta);
    yminus = FUN(X - ck.*delta);
    Gk = mean(repmat((yplus-yminus),DIM,1)./(2*ck.*delta),2);

    % update objectVector
    x = x - ak*Gk;

    % termination criterions
    fit = FUN(x);
    % stop if target or maxfunevals is reached
    if fit <= ftarget || feval(FUN, 'evaluations') >= maxfunevals
        termvalue = 1;
        break;
    end
    % stop if maxGenerations or fit is larger 1e30 (probably divergent
    % run)
    if k > maxGenerations || fit > 1e30
        termvalue = 0;
        break;
    end
    % stop if x has nan or inf entries
    if max(isnan(x)) == 1 || max(isinf(x)) == 1
        termvalue = 0;
        break;
    end

    % increase k
    k = k + 1;

end % of while loop

end % of function
```

---

$f_{101}$ in 5-D, N=15, mFE=513						$f_{101}$ in 20-D, N=15, mFE=2026						$f_{102}$ in 5-D, N=15, mFE=513						$f_{102}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	14	2.3e2	1.7e2	2.8e2	2.3e2	13	1.1e3	8.9e2	1.3e3	9.7e2	10	11	3.1e2	2.2e2	4.0e2	2.5e2	3	9.4e3	8.9e3	1.0e4	2.0e3		
1	9	6.0e2	5.0e2	6.9e2	3.9e2	12	1.7e3	1.5e3	2.0e3	1.5e3	1	8	7.1e2	6.1e2	8.1e2	3.6e2	0	48e+0	36e-1	10e+1	2.0e3		
1e-1	7	8.7e2	7.6e2	9.6e2	4.2e2	6	4.2e3	3.7e3	4.7e3	1.8e3	1e-1	4	1.7e3	1.6e3	1.9e3	4.9e2	.	.	.	.	.		
1e-3	4	1.8e3	1.7e3	1.9e3	5.1e2	4	7.0e3	6.6e3	7.3e3	2.0e3	1e-3	2	3.8e3	3.7e3	3.8e3	5.1e2	.	.	.	.	.		
1e-5	1	7.5e3	7.4e3	7.7e3	5.1e2	0	28e-2	35e-5	18e+0	2.0e3	1e-5	0	84e-2	52e-5	15e+0	4.5e2	.	.	.	.	.		
1e-8	0	73e-2	16e-6	89e-1	4.5e2	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{103}$ in 5-D, N=15, mFE=513						$f_{103}$ in 20-D, N=15, mFE=2013						$f_{104}$ in 5-D, N=15, mFE=514						$f_{104}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	6	9.7e2	8.1e2	1.1e3	4.0e2	5	4.7e3	4.1e3	5.4e3	1.7e3	10	0	14e+2	11e+1	38e+3	4.5e2	0	11e+4	12e+3	68e+4	1.8e3		
1	4	1.6e3	1.4e3	1.8e3	4.1e2	3	9.1e3	8.2e3	9.8e3	1.8e3	1	.	.	.	.	.	.	.	.	.	.		
1e-1	3	2.3e3	2.1e3	2.5e3	5.1e2	0	17e+1	16e-2	24e+1	8.9e1	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	1	7.4e3	7.1e3	7.7e3	5.1e2	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	0	19e+0	78e-4	78e+0	2.2e2	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{105}$ in 5-D, N=15, mFE=513						$f_{105}$ in 20-D, N=15, mFE=2004						$f_{106}$ in 5-D, N=15, mFE=513						$f_{106}$ in 20-D, N=15, mFE=2011					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	0	14e+2	42e+0	31e+3	4.5e2	0	26e+4	95e+3	60e+4	1.8e3	10	0	17e+2	21e+1	23e+3	5.0e2	0	35e+2	83e+1	19e+3	2.0e3		
1	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{107}$ in 5-D, N=15, mFE=513						$f_{107}$ in 20-D, N=15, mFE=2004						$f_{108}$ in 5-D, N=15, mFE=513						$f_{108}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	2	3.6e3	3.4e3	3.8e3	2.9e2	0	13e+1	54e+0	22e+1	8.9e2	10	1	7.4e3	7.2e3	7.7e3	5.1e2	0	21e+1	15e+1	33e+1	5.0e1		
1	0	30e+0	65e-1	48e+0	3.5e2	.	.	.	.	.	1	0	47e+0	24e+0	72e+0	2.2e2	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{109}$ in 5-D, N=15, mFE=513						$f_{109}$ in 20-D, N=15, mFE=2019						$f_{110}$ in 5-D, N=15, mFE=513						$f_{110}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	8	6.2e2	5.1e2	7.3e2	2.2e2	9	2.0e3	1.6e3	2.4e3	1.2e3	10	0	22e+3	45e+2	63e+3	1.2e2	0	31e+4	20e+4	52e+4	1.0e0		
1	5	1.3e3	1.2e3	1.4e3	4.5e2	5	5.2e3	4.6e3	5.7e3	2.0e3	1	.	.	.	.	.	.	.	.	.	.		
1e-1	2	3.8e3	3.8e3	3.8e3	5.1e2	1	2.9e4	2.8e4	3.0e4	2.0e3	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	0	69e-1	92e-3	40e+0	4.5e2	0	18e-1	18e-2	25e+1	1.3e3	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{111}$ in 5-D, N=15, mFE=513						$f_{111}$ in 20-D, N=15, mFE=2004						$f_{112}$ in 5-D, N=15, mFE=513						$f_{112}$ in 20-D, N=15, mFE=2016					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	0	43e+3	11e+3	15e+4	8.9e1	0	29e+4	18e+4	65e+4	8.9e1	10	0	85e+1	76e+0	69e+2	4.5e2	0	39e+2	27e+1	17e+3	1.8e3		
1	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{113}$ in 5-D, N=15, mFE=513						$f_{113}$ in 20-D, N=15, mFE=2004						$f_{114}$ in 5-D, N=15, mFE=513						$f_{114}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	1	7.6e3	7.5e3	7.7e3	5.1e2	0	22e+2	67e+1	40e+2	5.0e1	10	0	19e+1	32e+0	88e+1	7.0e1	0	15e+2	94e+1	34e+2	7.0e1		
1	0	29e+1	21e+0	12e+2	3.2e2	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{115}$ in 5-D, N=15, mFE=528						$f_{115}$ in 20-D, N=15, mFE=2040						$f_{116}$ in 5-D, N=15, mFE=513						$f_{116}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	3	2.4e3	2.2e3	2.6e3	5.1e2	2	1.4e4	1.3e4	1.5e4	1.3e3	10	0	33e+3	42e+2	24e+4	7.0e1	0	12e+4	72e+3	29e+4	5.0e1		
1	0	30e+0	63e-1	17e+1	4.0e2	0	63e+0	41e-1	20e+2	1.8e3	1	.	.	.	.	.	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{117}$ in 5-D, N=15, mFE=513						$f_{117}$ in 20-D, N=15, mFE=2004						$f_{118}$ in 5-D, N=15, mFE=513						$f_{118}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	0	27e+3	12e+2	12e+4	1.1e2	0	16e+4	73e+3	32e+4	1.0e0	10	0	96e+1	22e+1	10e+3	4.5e2	0	47e+2	12e+2	13e+4	1.8e3		
1	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		
$f_{119}$ in 5-D, N=15, mFE=519						$f_{119}$ in 20-D, N=15, mFE=2004						$f_{120}$ in 5-D, N=15, mFE=513						$f_{120}$ in 20-D, N=15, mFE=2004					
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>		
10	9	4.7e2	3.6e2	5.9e2	2.5e2	0	13e+1	53e+0	31e+1	5.0e1	10	4	1.6e3	1.4e3	1.8e3	3.8e2	0	83e+0	36e+0	22e+1	1.0e0		
1	0	74e-1	36e-1	26e+0	3.2e2	.	.	.	.	.	1	0	25e+0	36e-1	14e+1	1.4e2	.	.	.	.	.		
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.		
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.		
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.		
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.		

Figure 2: Determination of the initial values for the gains

---

```

function [a0,c0] = DetermineParameter(A,alpha,x,step,FUN,DIM)

    % generate matrix with trial vectors
    X = repmat(x,1,10);

    % c0
    dummy = FUN(X);
    c0 = max([std(dummy,1),1e-5]);

    % a0
    % generation of the simultaneous perturbation vector
    delta = 2*round(rand(DIM,10))-1;

    % function evaluation
    yplus = FUN(X + c0.*delta);
    yminus = FUN(X - c0.*delta);

    % gradient approximation
    gApprox = mean(repmat((yplus-yminus),DIM,1)./(2*c0.*delta),2);

    % mean of the magnitude of gradient element
    gMeanElement = abs(mean(gApprox));

    % determine parameter a
    a0 = step*(1+A)^alpha/gMeanElement;

end % of function

```

---

$f_{121}$ in 5-D, N=15, mFE=513						$f_{121}$ in 20-D, N=15, mFE=2044						$f_{122}$ in 5-D, N=15, mFE=513						$f_{122}$ in 20-D, N=15, mFE=2023								
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>
10	11	2.5e2	1.6e2	3.5e2	1.7e2	6	4.0e3	3.5e3	4.4e3	1.5e3	10	4	1.5e3	1.2e3	1.7e3	5.1e2	0	32e+0	14e+0	16e+1	1.0e0					
1	2	3.6e3	3.4e3	3.8e3	5.1e2	0	13e+0	52e-1	41e+0	1.8e3	1	0	13e+0	43e-1	25e+0	1.1e2	.	.	.	.	.	.	.	.	.	.
1e-1	1	7.6e3	7.6e3	7.7e3	5.1e2	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-3	0	42e-1	45e-2	35e+0	4.5e2	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
$f_{123}$ in 5-D, N=15, mFE=579						$f_{123}$ in 20-D, N=15, mFE=2181						$f_{124}$ in 5-D, N=15, mFE=531						$f_{124}$ in 20-D, N=15, mFE=2065								
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>
10	3	2.2e3	1.9e3	2.5e3	3.8e2	0	23e+0	14e+0	38e+0	1.0e3	10	9	4.2e2	3.0e2	5.4e2	2.1e2	8	2.3e3	1.8e3	2.8e3	1.3e3					
1	0	13e+0	72e-1	42e+0	1.0e2	.	.	.	.	.	1	0	78e-1	13e-1	31e+0	3.5e2	0	97e-1	62e-1	14e+0	1.3e3					
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
$f_{125}$ in 5-D, N=15, mFE=522						$f_{125}$ in 20-D, N=15, mFE=2013						$f_{126}$ in 5-D, N=15, mFE=513						$f_{126}$ in 20-D, N=15, mFE=2004								
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>
10	15	1.0e0	1.0e0	1.0e0	1.0e0	15	1.0e0	1.0e0	1.0e0	1.0e0	10	15	1.0e0	1.0e0	1.0e0	1.0e0	15	1.0e0	1.0e0	1.0e0	1.0e0					
1	15	1.1e2	7.9e1	1.4e2	1.1e2	11	1.1e3	7.8e2	1.5e3	8.8e2	1	4	1.5e3	1.3e3	1.7e3	4.0e2	0	25e-1	19e-1	37e-1	5.0e1					
1e-1	0	42e-2	17e-2	66e-2	2.8e2	0	77e-2	63e-2	13e-1	1.3e3	1e-1	0	19e-1	35e-2	44e-1	8.9e1	.	.	.	.	.	.	.	.	.	
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
$f_{127}$ in 5-D, N=15, mFE=527						$f_{127}$ in 20-D, N=15, mFE=2058						$f_{128}$ in 5-D, N=15, mFE=513						$f_{128}$ in 20-D, N=15, mFE=2013								
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>
10	15	1.0e0	1.0e0	1.0e0	1.0e0	15	1.0e0	1.0e0	1.0e0	1.0e0	10	1	7.5e3	7.4e3	7.7e3	5.1e2	0	76e+0	74e+0	80e+0	2.8e2					
1	13	1.5e2	8.4e1	2.1e2	1.4e2	8	2.3e3	1.8e3	2.9e3	9.4e2	1	0	39e+0	14e+0	60e+0	1.8e2	.	.	.	.	.	.	.	.	.	
1e-1	0	37e-2	23e-2	15e-1	1.2e2	0	97e-2	61e-2	18e-1	6.3e2	1e-1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
$f_{129}$ in 5-D, N=15, mFE=513						$f_{129}$ in 20-D, N=15, mFE=2004						$f_{130}$ in 5-D, N=15, mFE=513						$f_{130}$ in 20-D, N=15, mFE=2046								
$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	$\Delta f$	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>	#	ERT	10%	90%	RT <sub>succ</sub>
10	1	7.2e3	6.6e3	7.7e3	5.1e2	0	78e+0	74e+0	81e+0	3.2e2	10	7	7.2e2	5.8e2	8.6e2	3.1e2	0	77e+0	20e+0	84e+0	3.5e2					
1	0	35e+0	15e+0	55e+0	1.4e2	.	.	.	.	.	1	0	14e+0	20e-1	42e+0	2.2e2	.	.	.	.	.	.	.	.	.	.
1e-1	.	.	.	.	.	.	.	.	.	.	1e-1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-3	.	.	.	.	.	.	.	.	.	.	1e-3	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-5	.	.	.	.	.	.	.	.	.	.	1e-5	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
1e-8	.	.	.	.	.	.	.	.	.	.	1e-8	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.

Table 2: Shown are, for functions  $f_{121}$ - $f_{130}$  and for a given target difference to the optimal function value  $\Delta f$ : the number of successful trials (#); the expected running time to surpass  $f_{\text{opt}} + \Delta f$  (ERT, see Figure 4); the 10%-tile and 90%-tile of the bootstrap distribution of ERT; the average number of function evaluations in successful trials or, if none was successful, as last entry the median number of function evaluations to reach the best function value (RT<sub>succ</sub>). If  $f_{\text{opt}} + \Delta f$  was never reached, figures in *italics* denote the best achieved  $\Delta f$ -value of the median trial and the 10% and 90%-tile trial. Furthermore, N denotes the number of trials, and mFE denotes the maximum of number of function evaluations executed in one trial. See Figure 4 for the names of functions.

Figure 3: Restart procedure for the SPSA

---

```
function x = spsa(FUN, DIM, ftarget, maxfunevals)

% make sure to terminate
if isinf(maxfunevals) || maxfunevals > 1e5*DIM
    kmax = 1e5*DIM;
else
    kmax = maxfunevals;
end

% constant parameter
A = 0.1*kmax; % A approx 10% of max generations
gamma = 0.101; % reduction rate for ck (as recommended)
alpha = 0.602; % reduction rate for ak (as recommended)

% multistart such that ftarget is reached with reasonable prob.
for ilaunch = 1:100 % relaunch optimizer up to 100 times

    % restarts
    if ilaunch == 1 % initial scenario
        xstart = 8 * rand(DIM, 1) - 4; % random start solution
        step = 0.1; % parameter to determine a0
        lambda = 10; % number of gradient approximations
        [a0,c0] = DetermineParameter(A,alpha,xstart,step,FUN,DIM);

    else

        choice = round(3*rand) + 1;

        % if the xstart is changing, parameter a0 has to be newly
        % calculated

        switch choice

            case 1 % new point
                xstart = 8 * rand(DIM, 1) - 4;
                [a0,c0] = DetermineParameter(A,alpha,xstart,step,FUN,DIM);

            case 2 % improve old point
                xstart = x;
                [a0,c0] = DetermineParameter(A,alpha,xstart,step,FUN,DIM);

            case 3 % half the step size
                step = step/2;
                [a0,c0] = DetermineParameter(A,alpha,xstart,step,FUN,DIM);

            case 4 % increase lambda
                lambda = ceil(lambda * sqrt(2));

        end % switch case

    end


    % try spsa
    parameter = [a0,alpha,c0,gamma,A,lambda];
    [x,termvalue] = alg(FUN,xstart,parameter,kmax,ftarget,DIM,maxfunevals);

    if termvalue == 1
        break;
    end


end

end % of function
```

---



**Figure 4:** Expected Running Time (ERT, ●) to reach  $f_{\text{opt}} + \Delta f$  and median number of function evaluations of successful trials (+), shown for  $\Delta f = 10, 1, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-5}, 10^{-8}$  (the exponent is given in the legend of  $f_{101}$  and  $f_{130}$ ) versus dimension in log-log presentation. The  $\text{ERT}(\Delta f)$  equals to  $\#FES(\Delta f)$  divided by the number of successful trials, where a trial is successful if  $f_{\text{opt}} + \Delta f$  was surpassed during the trial. The  $\#FES(\Delta f)$  are the total number of function evaluations while  $f_{\text{opt}} + \Delta f$  was not surpassed during the trial from all respective trials (successful and unsuccessful), and  $f_{\text{opt}}$  denotes the optimal function value. Crosses (×) indicate the total number of function evaluations  $\#FES(-\infty)$ . Numbers above ERT-symbols indicate the number of successful trials. Annotated numbers on the ordinate are decimal logarithms. Additional grid lines show linear and quadratic scaling.



**Figure 5: Empirical cumulative distribution functions (ECDFs), plotting the fraction of trials versus running time (left) or  $\Delta f$ .** Left subplots: ECDF of the running time (number of function evaluations), divided by search space dimension  $D$ , to fall below  $f_{\text{opt}} + \Delta f$  with  $\Delta f = 10^k$ , where  $k$  is the first value in the legend. Right subplots: ECDF of the best achieved  $\Delta f$  divided by  $10^k$  (upper left lines in continuation of the left subplot), and best achieved  $\Delta f$  divided by  $10^{-8}$  for running times of  $D, 10D, 100D \dots$  function evaluations (from right to left cycling black-cyan-magenta). Top row: all results from all functions; second row: moderate noise functions; third row: severe noise functions; fourth row: severe noise and highly-multimodal functions. The legends indicate the number of functions that were solved in at least one trial. FEvals denotes number of function evaluations,  $D$  and DIM denote search space dimension, and  $\Delta f$  and Df denote the difference to the optimal function value.

## 7. REFERENCES

- [1] S. Finck, N. Hansen, R. Ros, and A. Auger. Real-parameter black-box optimization benchmarking 2009: Presentation of the noisy functions. Technical Report 2009/21, Research Center PPE, 2009.
- [2] N. Hansen, A. Auger, S. Finck, and R. Ros. Real-parameter black-box optimization benchmarking 2009: Experimental setup. Technical Report RR-6828, INRIA, 2009.
- [3] N. Hansen, S. Finck, R. Ros, and A. Auger. Real-parameter black-box optimization benchmarking 2009: Noisy functions definitions. Technical Report RR-6869, INRIA, 2009.
- [4] J. C. Spall. Multivariate Stochastic Approximation Using a Simultaneous Perturbation Gradient Approximation. *IEEE Transactions on Automatic Control*, 37(3):332–341, March 1992.
- [5] J. C. Spall. An Overview of the Simultaneous Perturbation Method for Efficient Optimization. *Johns Hopkins APL Technical Digest*, 19:482–492, 1998.
- [6] J. C. Spall. *Introduction to Stochastic Search and Optimization*. John Wiley & Sons, Hoboken, NJ, 2003.