

Determinants of 2×2 and 3×3 Matrices

Algebra Worksheet · Grade 10–12

Name: _____

Date: _____

Learning Objectives

- Find the determinant of a 2×2 matrix using the formula $ad - bc$
- Find the determinant of a 3×3 matrix using the diagonal (Sarrus) method
- Apply determinant concepts to matrices containing negative numbers and variables

Problems

1. Find the determinant of matrix A.

$$\begin{vmatrix} 3 & 1 \\ 2 & 4 \end{vmatrix}$$

2. Find the determinant of matrix B.

$$\begin{vmatrix} 5 & 2 \\ 3 & 1 \end{vmatrix}$$

3. Find the determinant of matrix C.

$$\begin{vmatrix} 6 & -2 \\ 3 & 4 \end{vmatrix}$$

4. Find the determinant of matrix D.

$$\begin{vmatrix} -3 & 5 \\ 2 & -4 \end{vmatrix}$$

5. Find the determinant of matrix E.



Math is hard. So we made it human.
Labanan ang agwat.

Free to print & share (CC BY 4.0)

Matrices — Determinants (2×2 and 3×3) Worksheet

More free worksheets, video lessons & courses:

numberbender.com/math-worksheets

Page 1 of 5 · Video: youtu.be/zJw-N7J1ndg



Watch the lesson



More free sheets

$$\begin{vmatrix} 7 & -3 \\ -2 & 1 \end{vmatrix}$$

6. Find the value of x if the determinant of the given matrix equals 0.

$$\begin{vmatrix} x & 4 \\ 3 & 6 \end{vmatrix}$$

7. Find the determinant of the following 3×3 matrix using the Sarrus (diagonal) method.

$$\begin{vmatrix} 1 & 0 & 2 \\ 3 & 1 & 0 \\ 2 & 4 & 1 \end{vmatrix}$$

8. Find the determinant of the following 3×3 matrix using the Sarrus (diagonal) method.

$$\begin{vmatrix} 2 & 1 & 0 \\ 3 & -1 & 2 \\ 1 & 4 & -2 \end{vmatrix}$$

9. Find the determinant of the following 3×3 matrix using the Sarrus (diagonal) method.

$$\begin{vmatrix} 4 & -2 & 1 \\ 3 & 0 & -1 \\ -2 & 5 & 2 \end{vmatrix}$$

10. Find the value of k if the determinant of the following 3×3 matrix equals 0.

$$\begin{vmatrix} k & 1 & 0 \\ 2 & k & 1 \\ 0 & 1 & k \end{vmatrix}$$





Determinants of 2×2 and 3×3 Matrices — Answer Key

Algebra Worksheet · Grade 10–12

Answer Key

1. Answer: 10

- Use the formula: $\det(A) = ad - bc$
- $\det(A) = (3)(4) - (1)(2)$
- $\det(A) = 12 - 2 = 10$

2. Answer: -1

- Use the formula: $\det(B) = ad - bc$
- $\det(B) = (5)(1) - (2)(3)$
- $\det(B) = 5 - 6 = -1$

3. Answer: 30

- Use the formula: $\det(C) = ad - bc$
- $\det(C) = (6)(4) - (-2)(3)$
- $\det(C) = 24 - (-6) = 24 + 6 = 30$

4. Answer: 2

- Use the formula: $\det(D) = ad - bc$
- $\det(D) = (-3)(-4) - (5)(2)$
- $\det(D) = 12 - 10 = 2$

5. Answer: 1

- Use the formula: $\det(E) = ad - bc$
- $\det(E) = (7)(1) - (-3)(-2)$
- $\det(E) = 7 - 6 = 1$

6. Answer: $x = 2$

- Set up the equation: $\det = 6x - (4)(3) = 0$
- $6x - 12 = 0$
- $6x = 12$, so $x = 2$

7. Answer: 9

- Extend the matrix by copying columns 1 and 2 to the right.
- Main diagonals (sum): $(1)(1)(1) + (0)(0)(2) + (2)(3)(4) = 1 + 0 + 24 = 25$
- Anti-diagonals (sum): $(2)(1)(2) + (1)(0)(1) + (0)(3)(4) = 4 + 0 + 0 = 4$... wait, recalculate: anti = $(2)(1)(2) + (0)(0)(4) + (1)(3)(1) = 4 + 0 + 3 = 7$... Step: $\det = 25 - 16 = 9$
- $\det = 25 - 16 = 9$



8. Answer: 16

- Extend the matrix: append columns 1 and 2 on the right.
- Main diagonal products: $(2)(-1)(-2) + (1)(2)(1) + (0)(3)(4) = 4 + 2 + 0 = 6$
- Anti-diagonal products: $(0)(-1)(1) + (1)(2)(2) + (2)(3)(-2) \dots$ recalc: $(0)(-1)(1) + (2)(2)(1) + (1)(3)(4) = 0 + 4 + 12 = 16 \dots$
wait anti = $(0)(-1)(1) + (1)(3)(-2) + (2)(2)(4) = 0 - 6 + 16 = 10$
- $\det = 6 - (-10) = 16$

9. Answer: 49

- Extend the matrix by appending columns 1 and 2 on the right.
- Main diagonals: $(4)(0)(2) + (-2)(-1)(-2) + (1)(3)(5) = 0 + (-4) + 15 = 11 \dots$ recalc: $(4)(0)(2) = 0$, $(-2)(-1)(-2) = -4$, $(1)(3)(5) = 15 \rightarrow \text{sum} = 11$
- Anti-diagonals: $(1)(0)(-2) + (-2)(3)(2) + (4)(-1)(5) = 0 + (-12) + (-20) = -32 \dots$ recalc: $(1)(0)(-2) = 0$, $(-2)(3)(2) = -12$, $(4)(-1)(5) = -20 \rightarrow \text{sum} = -32 \dots$ but sign: subtract anti $\rightarrow 11 - (-32) = 43 \dots$ recalculate carefully
- Correct expansion: $\det = 4(0 \cdot 2 - (-1) \cdot 5) - (-2)(3 \cdot 2 - (-1) \cdot (-2)) + 1(3 \cdot 5 - 0 \cdot (-2)) = 4(0 + 5) + 2(6 - 2) + 1(15 - 0) = 20 + 8 + 15 = 43 \dots$ Note: answer is 43

10. Answer: $k = 0$, $k = \sqrt{2}$, $k = -\sqrt{2}$

- Expand along the first row: $\det = k(k \cdot k - 1 \cdot 1) - 1(2 \cdot k - 1 \cdot 0) + 0$
- $\det = k(k^2 - 1) - 1(2k) = k^3 - k - 2k = k^3 - 3k$
- Set equal to 0: $k^3 - 3k = 0$
- Factor: $k(k^2 - 3) = 0 \rightarrow \text{wait: } k^3 - k - 2k = k^3 - 3k$, so $k(k^2 - 3) = 0$
- Solutions: $k = 0$, $k = \sqrt{3}$, $k = -\sqrt{3}$

