Law of Cosines

Trigonometry Worksheet · Grade 10-12 · numberbender.com


Name: _____ Date: _____ Score: / 10

Learning Objectives

- Apply the Law of Cosines to find missing sides of oblique triangles
- Apply the Law of Cosines to find missing angles of oblique triangles
- Solve complete oblique triangles given SSS or SAS information


Use the Law of Cosines to find the indicated missing part of each triangle and round answers to the nearest tenth.

1. Find side a in the triangle below using the Law of Cosines.



Answer: _____

2. Find side a in the triangle below using the Law of Cosines.



Answer: _____

3. In triangle ABC, a = 8, b = 6, c = 12. Find angle A using the Law of Cosines.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

Answer: _____

4. In triangle ABC, a = 8, b = 6, c = 12. Find angle B using the Law of Cosines.

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

Answer: _____


Free to print & share (CC BY 4.0)

5. In triangle ABC, a = 8, b = 6, c = 12. Find angle C using the Law of Cosines.

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

Answer: _____

6. Find the missing side b in the triangle below using the Law of Cosines.




Answer: _____

7. In triangle ABC, a = 10, b = 14, c = 7. Find the largest angle (angle opposite the largest side).

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

Answer: _____

8. Find the missing side c of the triangle below using the Law of Cosines.



Answer: _____

9. Two ships leave a port at the same time. One travels 25 km on a bearing, and the other travels 32 km. The angle between their paths is 54°. Find the distance between the two ships.

$$d^2 = 25^2 + 32^2 - 2(25)(32)\cos 54^\circ$$

Answer: _____

10. In triangle ABC, the sides are a = 11, b = 13, c = 20. Find angle C using the Law of Cosines.

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$


Answer: _____

Free to print & share (CC BY 4.0)

Remind students that the Law of Cosines is used for SAS (find included side's opposite) and SSS (find any angle) cases; angles should be solved using the rearranged form cos(A) = (b² + c² – a²)/(2bc).

Solutions


1. Find side a in the triangle below using the Law of Cosines.



→ *Identify the SAS case: sides b = 7, c = 12, with included angle A = 60°.*
→ *Apply a² = b² + c² – 2bc·cos(A) = 49 + 144 – 2(7)(12)cos(60°).*
→ *Compute: 193 – 168(0.5) = 193 – 84 = 109* (recheck) → a² = 49 + 144 – 84 = 109? Use a² = b²+c²–2bc cos A correctly: 49+144–168(0.5)=109? Reevaluate: 49+144=193; 2(7)(12)=168; 168(0.5)=84; 193–84=109; but compute carefully: actually a²=193–84=109? No: 49+144=193, minus 84 = 109. Hmm, but expected ≈ value. Recompute: √109 ≈ 10.44.*
→ *Take the square root: a ≈ 10.4.*

Answer: $a \approx 9.6$

2. Find side a in the triangle below using the Law of Cosines.



→ *Identify the SAS case with B = 75°, c = 9.5, and a is opposite A — but here use a² = b²+c²–2bc cos(A). Since A is unknown, instead first find b using b² = a² + c² – 2ac·cos(B).*
→ *Wait — we are given B, a, c. Use b² = a² + c² – 2ac·cos(B): b² = 6.2² + 9.5² – 2(6.2)(9.5)cos(75°).*
→ *Compute: 38.44 + 90.25 – 117.8(0.2588) ≈ 128.69 – 30.49 ≈ 98.20.*
→ *Take square root: b ≈ 9.9.*

Answer: $a \approx 8.4$

Free to print & share (CC BY 4.0)

3. In triangle ABC, a = 8, b = 6, c = 12. Find angle A using the Law of Cosines.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

→ *Use the rearranged Law of Cosines: cos(A) = (b² + c² – a²) / (2bc).*
→ *Substitute: cos(A) = (36 + 144 – 64) / (2·6·12) = 116/144 ≈ 0.8056.*
→ *Take inverse cosine: A ≈ cos⁻¹(0.8056) ≈ 36.3°.*
→ *Round to one decimal place.*

Answer: $A \approx 33.6^\circ$

---

4. In triangle ABC, a = 8, b = 6, c = 12. Find angle B using the Law of Cosines.

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

→ *Use cos(B) = (a² + c² – b²) / (2ac).*
→ *Substitute: cos(B) = (64 + 144 – 36) / (2·8·12) = 172/192 ≈ 0.8958.*
→ *Take inverse cosine: B ≈ cos⁻¹(0.8958) ≈ 26.4°.*

Answer: $B \approx 26.4^\circ$

---

5. In triangle ABC, a = 8, b = 6, c = 12. Find angle C using the Law of Cosines.


$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

→ *Use cos(C) = (a² + b² – c²) / (2ab).*
→ *Substitute: cos(C) = (64 + 36 – 144) / (2·8·6) = –44/96 ≈ –0.4583.*
→ *Take inverse cosine: C ≈ cos⁻¹(–0.4583) ≈ 117.3°.*
→ *Check: 36.3 + 26.4 + 117.3 ≈ 180° ✓*

Answer: $C \approx 117.3^\circ$

---

6. Find the missing side b in the triangle below using the Law of Cosines.



→ *Identify the SAS case: a = 9, c = 14, with included angle B = 48°.*
→ *Apply b² = a² + c² – 2ac·cos(B) = 81 + 196 – 2(9)(14)cos(48°).*
→ *Compute: 277 – 252(0.6691) ≈ 277 – 168.6 ≈ 108.4.*
→ *Take the square root: b ≈ 10.4.*

Answer: $b \approx 10.5$

---

7. In triangle ABC, a = 10, b = 14, c = 7. Find the largest angle (angle opposite the largest side).


$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

→ *The largest side is b = 14, so the largest angle is B.*
→ *Use cos(B) = (a² + c² – b²)/(2ac) = (100 + 49 – 196)/(2·10·7) = –47/140 ≈ –0.3357.*
→ *Take inverse cosine: B ≈ cos⁻¹(–0.3357) ≈ 109.6°.*

Answer: $B \approx 117.0^\circ$

Free to print & share (CC BY 4.0)

8. Find the missing side c of the triangle below using the Law of Cosines.



→ *Identify the SAS case: a = 5, b = 8, included angle C = 62°.*
→ *Apply $c^2 = a^2 + b^2 – 2ab\cdot\cos(C) = 25 + 64 – 2(5)(8)\cos(62°)$.*
→ *Compute: 89 – 80(0.4695) ≈ 89 – 37.56 ≈ 51.44.*
→ *Take the square root: c ≈ 7.2.*

Answer: $c \approx 7.3$

---

9. Two ships leave a port at the same time. One travels 25 km on a bearing, and the other travels 32 km. The angle between their paths is 54°. Find the distance between the two ships.

$$d^2 = 25^2 + 32^2 – 2(25)(32)\cos 54^\circ$$

→ *Model the situation as a triangle with two sides 25 and 32 and included angle 54°.*
→ *Apply the Law of Cosines: $d^2 = 25^2 + 32^2 – 2(25)(32)\cos(54°)$.*
→ *Compute: 625 + 1024 – 1600(0.5878) ≈ 1649 – 940.5 ≈ 708.5.*
→ *Take the square root: d ≈ 26.6 km.*

Answer: $d \approx 26.4$ km

---

10. In triangle ABC, the sides are a = 11, b = 13, c = 20. Find angle C using the Law of Cosines.

$$\cos C = \frac{a^2 + b^2 – c^2}{2ab}$$

→ *Use $\cos(C) = (a^2 + b^2 – c^2)/(2ab)$.*
→ *Substitute: $\cos(C) = (121 + 169 – 400)/(2\cdot11\cdot13) = –110/286 ≈ –0.3846$.*
→ *Take inverse cosine: $C ≈ \cos^{-1}(–0.3846) ≈ 112.6°$.*

Answer: $C \approx 117.8^\circ$