

Self perception in bacteria: quorum sensing with acylated homoserine lactones

Clay Fuqua* and E Peter Greenberg†

A variety of Gram-negative bacteria produce membrane permeant, acylated homoserine lactone (HL) pheromones that act as cell density cues. Synthesis and response to these factors requires proteins homologous to the LuxI acylhomoserine lactone synthase and the LuxR transcription factor from *Vibrio fischeri*. Recent genetic and biochemical studies have begun to provide a mechanistic understanding of acyl HL dependent gene regulation. Examination of the role of acyl HLs in diverse bacteria positions LuxR–LuxI type systems within an increasingly broad regulatory context and suggests that, in some bacteria, they comprise a global regulatory circuit.

Addresses

*Department of Biology, Trinity University, San Antonio, TX 78212, USA; e-mail: cfuqua@trinity.edu

†Department of Microbiology, University of Iowa, Iowa City, IA 52242, USA; e-mail: epgreen@blue.weeg.uiowa.edu

Current Opinion in Microbiology 1998, 1:183–189

<http://biomednet.com/elecreff/1369527400100183>

© Current Biology Ltd ISSN 1369-5274

Abbreviations

ACP acyl carrier protein
HL homoserine lactone
MTA 5-methylthioadenosine
SAM S-adenosylmethionine

Introduction

In animals and plants, membrane-permeant signal molecules, including steroid and brassinosteroid hormones, nitric oxide and methyl jasmonate, are well established as important regulators of development, cell differentiation and metabolism. Considering the widespread occurrence of such diffusible signals in eukaryotic organisms, it is not entirely surprising that prokaryotes also employ membrane-permeant signal molecules. Over the past few years, it has become clear that a diverse collection of Gram-negative bacteria produce acylated homoserine lactone (HL) derivatives that function in cell–cell communication. Bacteria utilize acyl HLs to monitor the density of cells within their own population in a process called quorum sensing [1–4]. Each cell in the population produces a low basal level of the diffusible acyl HL via the activity of an acyl HL synthase, usually a member of the LuxI family of proteins. As the bacterial population density rises, the concentration of the acyl HL also increases. At a sufficiently high population density, in essence a bacterial quorum, the accumulated acyl HL interacts with a receptor protein, usually a member of the LuxR family of transcription regulators, that acts to control a specific

constellation of acyl HL dependent genes. In different bacteria, the basic mechanism of acyl HL regulation is conserved but the target genes are extremely varied. The first acyl HL quorum sensor to be fully described was that of the marine bacterium *Vibrio fischeri* that regulates bioluminescence (*lux*) gene expression (for a review see [5]); however, acyl HL quorum sensors in other bacteria regulate a range of functions including virulence genes in *Pseudomonas aeruginosa*, conjugal transfer in *Agrobacterium tumefaciens*, swarming motility in *Serratia liquefaciens*, and antibiotic production in *Erwinia carotovora* [1,6]. New acyl HL quorum sensors with different regulatory targets are reported monthly and the past five years have seen the list of bacteria known to employ these regulatory systems grow from certain bioluminescent marine vibrios to a large group of phylogenetically diverse bacteria.

The purpose of this review is to highlight several topics within the area of acyl HL quorum sensing that have seen particularly rapid progress during the past two years (see [2–4] for more comprehensive reviews). We will discuss the biochemical basis of acyl HL synthesis and perception as well as the global nature of at least a subset of these regulatory systems.


Enzymology of acyl HL synthesis

Acyl HLs have the general structure of an acyl chain of variable length and chemistry conjugated via an amide linkage to a lactonized homoserine moiety (Figure 1). Genetic studies of *luxI* and related genes clearly indicated a role in synthesis of the acyl HL signal. Only recently, however, has it been established that LuxI-type proteins are in fact acyl HL synthases that recruit their substrates from normal metabolic precursors.

Studies of LuxI and its homologs TraI from *A. tumefaciens* and RhII from *P. aeruginosa* show that these proteins catalyze the ligation of S-adenosylmethionine (SAM) and a fatty acyl chain derived from acyl–acyl carrier protein (ACP) conjugates ([7,8,9]; MR Parsek, EP Greenberg, unpublished data). A catalytic model has been proposed that entails the nucleophilic attack of the C1 position of the acyl–ACP conjugate by the amino nitrogen of SAM, generating an amide linkage (Figure 1). The lactonization of SAM, subsequent to, or concomitant with, amide bond formation, results in ring formation and release of the acyl HL, as well as the formation of the side product 5-methylthioadenosine (MTA).

Although this basic mechanism is likely to be correct, there remain a number of catalytic details that are unclear; for example, acyl transferases that interact with ACP-linked

Figure 1



Structure and function of LuxR-type (R) and LuxI-type (I) proteins. Proteins are depicted in linear form. Amino acid residues conserved in all members of the protein family, or implicated from mutational analysis, are marked by vertical lines. Conserved amino acid residues are in bold and positions where mutations have been found to abolish activity (*) are indicated. For LuxI-type proteins, positions where inactivating mutations have been isolated in LuxI or RhII, but not both, (^) are indicated. The upper portion of the figure depicts a proposed model for acyl HL synthesis by LuxI-type proteins. The lower portion of the figure illustrates transcriptional stimulation by LuxR-type transcriptional activators. H-T-H, helix-turn-helix. SH, sulfhydryl group on 5'-phosphopantetheine prosthetic group of ACP.

substrates often form transient covalent linkages with the acyl chains via ester or thioester bonds at cysteine or serine residues. The existence of a conserved cysteine residue and a conserved serine residue (Cys68 and Ser104 in LuxI) prompted the proposal that acyl HL synthesis might involve a transient acylation mechanism [2,8[•],10]. Directed mutational analyses of the LuxI, TraI and RhII acyl HL synthases, however, have revealed that the conserved cysteine (as well as several nonconserved cysteines in each protein) is dispensable for activity ([11,12]; C Fuqua, unpublished data). A mutation in the conserved serine in TraI abolishes activity, but this residue is not essential in LuxI and RhII, suggesting an indirect role in TraI function. Although these results demonstrate that transient acylation does not occur at these residues, it remains possible that a covalent linkage occurs with another residue in the protein.

Mutational analyses of LuxI and RhII have indicated a role for conserved charged residues clustered in the amino-terminal halves (residues 25–104 in LuxI; see Figure 1) of these proteins in acyl HL synthesis [11,12]. Seven out of the ten residues that are absolutely conserved among all LuxI-type proteins are charged (Arg25, Glu44, Asp46, Asp49, Arg70, Glu100, and Arg104 in LuxI), and mutations in any of these residues can lead to a loss of activity. Many other enzymes that utilize SAM as a substrate, namely, the methyl transferases, also employ charged residues within defined SAM-binding sites [13,14]; however, there are no discernable SAM-binding sites in the amino acid sequences of acyl HL synthases. Mutations in the three nonpolar conserved residues (Pro29, Trp35 and Pro84 in LuxI) significantly reduce, but do not abolish, activity [11,12]. Although a region in the carboxy-terminal half of LuxI was implicated in catalysis (residues 133–164; see Figure 1), mutations within the corresponding region of RhII do not disrupt activity [12].

Putative acyl HL synthases, AinS and LuxM, that share no homology with LuxI have been identified in *V. fischeri* and *V. harveyi*, respectively. These two proteins share a region of strong sequence conservation with each other and are required for acyl HL synthesis [15,16[•]], and thus appear to represent a second family of acyl HL synthases. Preliminary *in vitro* studies suggest that AinS will synthesize its cognate acyl HL (*N*-octanoyl homoserine lactone) using SAM and octanoyl-ACP (BII Hanzelka, PV Dunlap, EP Greenberg, unpublished data). Further comparisons between these proteins and LuxI-type acyl HL synthases should help to establish general enzymatic requirements for acyl HL synthesis.

Acyl HL perception and transcriptional activation by LuxR-type proteins

Genetic analyses of LuxR suggest that it is composed of two functional modules or domains: an amino-terminal domain (residues 20–156) with an acyl HL binding region (residues 79–120) and a carboxy-terminal transcription

regulation domain (residues 160–250), which includes a helix–turn–helix DNA-binding motif (residues 190–210; see Figure 1). Mutations in key residues within the amino-terminal acyl HL binding domain abolish or severely reduce responsiveness to *N*-3-oxohexanoyl HL (Figure 1; see [17,18]). Furthermore, the amino-terminal half of the protein (residues 1–193) is sufficient to sequester radiolabeled *N*-3-oxohexanoyl HL when expressed in *Escherichia coli* and deletions or missense mutations resulting in acyl HL nonresponsive LuxR alleles abolish this binding [19].

The DNA-binding domains of LuxR-type proteins share sequence similarity with a much larger group of transcription factors (the so-called LuxR or FixJ superfamily; see [20,21]). LuxR molecules consisting of amino-terminal deletion polypeptides (residues 156–250 and 162–250) show acyl HL independent activation of *lux* genes [22]. These results suggested that in the absence of acyl HL, the amino-terminal half of the protein blocks the function of the DNA-binding domain. Interaction with the acyl HL abolishes this inhibition and allows transcriptional activation. Similar amino-terminal inhibitory domains have been described for FixJ and NarL, two component-type response regulators of the LuxR superfamily that do not interact with acyl HLs but are instead phosphorylated at a conserved aspartate residue [21,23]. The recent three-dimensional structure of NarL reveals that the amino-terminal domain of this protein in the inactive, nonphosphorylated state effectively blocks residues within the helix–turn–helix motif, presumably preventing DNA binding [24[•]].

In addition, the observation of dominant-negative effects when inactive alleles of both LuxR and the TraR protein from *A. tumefaciens* are coexpressed with the wild-type proteins suggest that transcriptional activation of target promoters requires formation of multimers [25,26]. It remains unclear how the inhibitory effect of the amino-terminal acyl HL binding domain is integrated with multimerization. Perhaps binding of the pheromone elaborates a sequestered dimerization site and, in turn, dimerization facilitates DNA binding. The mechanistic integration of these two processes is possibly reflected by the observation that amino acid substitutions that result in acyl HL independent *luxR* proteins have been isolated across the length of the protein (residues 9–236; see [27,28]).

Transcriptional activation of the *lux* operon by LuxR requires a *cis*-acting DNA element, a 20 base pair inverted repeat centered at about –40, now called the *lux* box [1,29]. Similar *lux* box-like sequences are found upstream of at least some promoters regulated by LuxR-type proteins in other bacteria [1,30–33]. Specific DNA binding by LuxR-type proteins has been reported. Using a truncated, acyl HL independent allele of LuxR, Stevens *et al.* [34,35] demonstrated specific binding to the *lux* regulatory

DNA; however, binding to the *lux* box and *lux* promoter, as assessed by DNase I protection analysis, was only observed in the presence of RNA polymerase. It was therefore impossible to determine which of the two proteins was responsible for the observed protection. It is unclear whether this apparent synergism reflects the mechanism of activation by LuxR or is an artifact of using the acyl HL independent protein. More recently, the full length LasR protein from *P. aeruginosa*, purified as a glutathione-S-transferase (GST) fusion, was reported to bind to *lux* box-like sequences upstream of the *lasB* promoter sequences in the absence of RNA polymerase [36]. Further studies are required to confirm this report.

There are several acyl HL regulatory systems, thus far exclusively from the plant pathogenic *Erwinia* species, in which the LuxR-type protein acts as a transcriptional repressor rather than an activator ([37]; S Beck von Bodman, personal communication). As yet, little is known about LuxR-type repressors, although their mechanism of action is likely to deviate substantially from those employed by LuxR-type transcriptional activators.

Global gene regulation by acyl HL quorum sensors

The discovery of acyl HL based quorum sensing in diverse bacteria has provided several examples of acyl HLs that regulate multiple genes of vastly different function within the same bacterium. The most extensively studied of these is found in the opportunistic plant and animal pathogen *P. aeruginosa* (Figure 2), where recent genetic analyses have revealed a broad range of target genes. The LasR transcriptional regulator and *N*-3-oxododecanoyl HL (synthesized by the LasI protein) were identified by the role they play in regulating the expression of the *lasB* gene, which encodes an elastase involved in virulence [38,39]. LasR also affects the expression of the *lasA* gene (a second elastase), the *apr* alkaline protease, and may indirectly influence exotoxin A (ETA) activity (Figure 2; [40,41]). In addition, LasR controls the expression of the gene for a second LuxR-type protein called RhlR [42•,43•]. The RhlR protein was identified as a regulator of genes required for rhamnolipid biosynthesis (*rhlAB*) and requires the activity of the RhlI acyl HL synthase to generate its inducing ligand *N*-butyryl HL [44–47]. *P. aeruginosa* mutants deficient in *N*-butyryl HL synthesis do not produce the secondary metabolites pyocyanin and hydrogen cyanide, and also manifest defects in elastase gene expression [44,48]. Furthermore, expression of *xcp* genes encoding proteins of the outer membrane component of the general secretion pathway is activated by LasR and RhlR [49•]. Finally, there is a report that exogenous *N*-3-oxohexanoyl HL promotes phenazine antibiotic synthesis, although the mechanism for this is unclear [50].

In a possible explanation for the alarmingly broad regulatory net of acyl HLs in *P. aeruginosa*, recent reports

suggest that RhlR and *N*-butyryl HL are required for expression of the *rpoS* stationary-phase sigma factor gene (Figure 2; [42•,51]). In *E. coli* there are a wide variety of genes that are known to be part of the RpoS regulon [52], and it is assumed that the *P. aeruginosa* *rpoS* gene product has a similarly wide regulatory influence. Alterations in specific functions in a *las* or *rhl* mutant background may reflect direct control via the quorum-sensing cascade, or indirect control by RpoS. Clearly, in addition to the range of functions that have been shown to be definitively within the Las–Rhl signal cascade, the effect of placing *rpoS* under acyl HL control should greatly broaden the Las–Rhl regulon.


Although population density (acyl HL) and nutritional status of the cell (RpoS) are not obligately linked, they do overlap. It is not entirely surprising that bacteria have evolved mechanisms for integrating responses to both. An interesting example where both conditions are likely is in natural bacterial surface films. Indeed, recent work directly demonstrates the production of acyl HL signals by naturally occurring biofilms [53]. The importance of acyl HL quorum sensing in biofilm metabolism is an area of active investigation. Studies with *P. aeruginosa* mutants suggest that *lasI* is essential for proper biofilm development [54].

LasR and *N*-3-oxododecanoyl HL stand at the head of an impressive and rapidly expanding list of quorum-dependent functions in *P. aeruginosa*; however, there are several reasons not to consider LasR as a master switch. Recent studies suggest that *lasR* gene expression is itself regulated by the Vfr protein (virulence factor regulator, a homolog of the *E. coli* catabolite repressor protein) in response to an as yet unidentified signal [55,56]. Furthermore, it has also been proposed that *rhlR* and perhaps *lasR* are under the control of the GacA protein, a two-component response regulator and, in fact, a member of the LuxR/FixJ superfamily [57]. GacA and its sensor kinase partner LemA are known to regulate the expression of a variety of virulence genes in pathogenic pseudomonads in response to an unidentified environmental signal [58]. GacA and LemA also control the expression of the *phzI* acyl HL synthase that directly regulates phenazine antibiotic synthesis in the pseudomonad *P. aureofaciens* ([59]; S Chancey, LS Pierson, personal communication); therefore, in at least two pseudomonads, GacA–LemA regulation influences the function of a LuxR–LuxI type quorum sensor. Future research should address the ways in which LuxR–LuxI systems are integrated into the general regulatory circuits of a bacterial cell.

Conclusion

Over recent years, acyl HL dependent gene regulation has received increasing recognition as an important form of cell–cell communication in Gram-negative bacteria. Often, the genes controlled by LuxR–LuxI type quorum sensors are involved in important microbial processes,

Figure 2



Quorum sensing and global regulation in *P. aeruginosa*. A proposed regulatory cascade incorporating the *las* and *rhl* quorum sensors. Dashed arrows indicate activation of target promoters. Solid open arrows indicate enzymatic activity: kinase activity for LemA and acyl HL synthesis by LasI and RhlI. Squiggles indicate translation products. Although the response regulator GacA is shown in association with its cognate sensor kinase LemA, there is as yet no experimental evidence to support this speculation. Unidentified environmental signals to which GacA (LemA) and Vfr respond are indicated as question marks, as are those genes transcribed by *P. aeruginosa* σ^S RNA polymerase.

including microbial–host interactions such as pathogenesis and symbiosis. As our understanding of the biochemical mechanisms by which bacteria synthesize and respond to acyl HLs increases, human intervention designed to manipulate this regulation and to harness or mollify quorum-dependent gene products will become possible. Furthermore, the observation that acyl HL regulation can act in a global fashion suggests that a wide range of functions may be targeted with a single strategy.

Acknowledgements

We wish to thank Susanne Beck von Bodman and Sandy Pierson for sharing unpublished results. Research in the authors' laboratories is supported by the National Science Foundation (MCB-9723837) to WCF and the National Science Foundation, Cystic Fibrosis Foundation and Office of Naval Research to EP Greenberg.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Fuqua WC, Winans SC, Greenberg EP: **Quorum sensing in bacteria: the LuxR/LuxI family of cell density-responsive transcriptional regulators.** *J Bacteriol* 1994, **176**:269-275.
2. Fuqua C, Winans SC, Greenberg EP: **Census and consensus in bacterial ecosystems: the LuxR–LuxI family of quorum-sensing transcriptional regulators.** *Annu Rev Microbiol* 1996, **50**:591-624.
3. Gray K: **Intercellular communication and group behavior in bacteria.** *Trends Microbiol* 1997, **5**:184-188.
4. Swift S, Throup JP, Williams P, Salmon GPC, Stewart GSAB: **Quorum sensing: a population density component in the determination of bacterial phenotype.** *Trends Biochem Sci* 1996, **21**:214-219.
5. Dunlap PV, Greenberg EP: **Role of intercellular chemical communication in the *Vibrio fischeri*–monocentric fish symbiosis.** In *Microbial Cell–Cell Interactions*. Edited by Dworkin M. Washington DC: American Society for Microbiology; 1991:219-253.
6. Eberl L, Winson MK, Sternberg C, Stewart GSAB, Christiansen G, Chhabra SR, Bycroft BW, Williams P, Molin S, Givskov M: **Involvement of *N*-acyl-L-homoserine lactone autoinducers in control of multicellular behavior of *Serratia liquefaciens*.** *Mol Microbiol* 1996, **20**:127-136.
7. Hanzelka BL, Greenberg EP: **Quorum sensing in *Vibrio fischeri*: evidence that *S*-adenosylmethionine is the amino acid substrate for autoinducer synthesis.** *J Bacteriol* 1996, **178**:5291-5294.
8. Moré MJ, Finger LD, Stryker JL, Fuqua C, Eberhard A, Winans SC: **Enzymatic synthesis of a quorum-sensing autoinducer through use of defined substrates.** *Science* 1996, **272**:1655-1658.
9. Schaefer AL, Val DL, Hanzelka BL, Cronan JE, Greenberg EP: **Generation of cell-to-cell signals in quorum sensing: acyl homoserine activity of purified *Vibrio fischeri* LuxI protein.** *Proc Natl Acad Sci USA* 1996, **93**:9505-9509.
10. Sitnikov D, Schineller JB, Baldwin TO: **Transcriptional regulation of bioluminescence genes from *Vibrio fischeri*.** *Mol Microbiol* 1995, **17**:801-812.
11. Hanzelka BL, Stevens AM, Parsek MR, Crone TJ, Greenberg EP: **Mutational analysis of the *Vibrio fischeri* LuxI polypeptide: critical regions of an autoinducer synthase.** *J Bacteriol* 1997, **179**:4882-4887.
12. Parsek MR, Schaefer AL, Greenberg EP: **Analysis of random and site-directed mutations in *rhlI*, a *Pseudomonas aeruginosa* gene encoding an acylhomoserine lactone synthase.** *Mol Microbiol* 1997, **26**:301-310.
13. Kumar S, Cheng X, Klimasauskas S, Mi S, Posfai J, Roberts RJ, Wilson GG: **The DNA (cytosine-5) methyltransferases.** *Nucleic Acids Res* 1994, **22**:1-10.
14. Labahn J, Granzin J, Schluckebier G, Robinson DP, Jack WE, Schildkraut I, Saenger W: **Three-dimensional structure of the adenine-specific DNA methyltransferase MtaI in complex with the cofactor *S*-adenosylmethionine.** *Proc Natl Acad Sci USA* 1994, **91**:10957-10961.
15. Bassler BL, Wright M, Showalter RE, Silverman MR: **Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence.** *Mol Microbiol* 1993, **9**:773-786.
16. Gilson L, Kuo A, Dunlap PV: **AinS and a new family of autoinducer synthesis proteins.** *J Bacteriol* 1995, **177**:6946-6951.
17. Shadel GS, Young R, Baldwin TO: **Use of regulated cell lysis in a lethal genetic selection in *Escherichia coli*: identification of the autoinducer-binding region of the LuxR protein from *Vibrio fischeri* ATCC 7744.** *J Bacteriol* 1990, **172**:3980-3987.
18. Slock J, Kolibachuk D, Greenberg EP: **Critical regions of the *Vibrio fischeri* LuxR protein defined by mutational analysis.** *J Bacteriol* 1990, **172**:3974-3979.
19. Hanzelka BL, Greenberg EP: **Evidence that the N-terminal region of the *Vibrio fischeri* LuxR protein constitutes an autoinducer-binding domain.** *J Bacteriol* 1995, **177**:815-817.
20. Henikoff S, Wallace JC, Brown JP: **Finding protein similarities with nucleotide sequence databases.** *Methods Enzymol* 1990, **183**:111-132.
21. Kahn D, Ditta G: **Modular structure of FixJ: homology of the transcriptional activation domain with the –35 binding domain of sigma factors.** *Mol Microbiol* 1991, **5**:987-997.
22. Choi SH, Greenberg EP: **The C-terminal region of the *Vibrio fischeri* LuxR protein contains an inducer-independent lux gene activating domain.** *Proc Natl Acad Sci USA* 1991, **88**:11115-11119.
23. Da Re S, Bertagnoli S, Fourment J, Reyat J-M, Kahn D: **Intramolecular signal transduction within the FixJ transcriptional activator: *in vitro* evidence for the inhibitory effect of the phosphorylatable regulatory domain.** *Nucl Acids Res* 1994, **9**:1555-1561.
24. Baikalov I, Schroder I, Kaczor-Grzekowiak M, Grzekowiak K, Gunsalus RP, Dickerson RE: **Structure of the *Escherichia coli* response regulator NarL.** *Biochemistry* 1996, **35**:11053-11061.
25. Choi SH, Greenberg EP: **Genetic evidence for multimerization of LuxR, the transcriptional activator of *Vibrio fischeri* luminescence.** *Mol Mar Biol Biotechnol* 1992, **1**:408-413.
26. Zhu J, Winans SC: **Activity of the quorum-sensing regulator TraR of *Agrobacterium tumefaciens* is inhibited by a truncated, dominant defective TraR-like protein.** *Mol Microbiol* 1998, **27**:289-298.
27. Poellinger KA, Lee JP, Parales JV, Greenberg EP: **Intragenic suppression of a luxR mutation: characterization of an autoinducer-independent LuxR.** *FEMS Microbiol Lett* 1995, **129**:97-102.
28. Sitnikov DM, Shadel GS, Baldwin TO: **Autoinducer-independent mutants of the LuxR transcriptional activator exhibit differential effects on the two lux promoters of *Vibrio fischeri*.** *Mol Gen Genet* 1996, **252**:622-625.

The identification of a gene other than *luxI* that directs the synthesis of an acyl homoserine lactone (HL) in *Vibrio fischeri* is described. The product of this gene, AinS, shows no similarity to any member of the LuxI family. Rather, it is related to a protein implicated in acyl HL synthesis in *V. harveyi*. This suggests that these gene products represent a second, newly recognized family of autoinducer synthases.

This paper provides structural information supporting the view of inhibition of DNA binding by an amino-terminal regulatory domain in a two-component response regulator related to LuxR.

29. Devine JH, Shadel GS, Baldwin TO: Identification of the operator of the *lux* regulon from the *Vibrio fischeri* strain ATCC7744. *Proc Natl Acad Sci USA* 1989, **86**:5688-5692.
30. Fuqua C, Winans SC: Localization of the OccR-activated and TraR-activated promoters that express two ABC-type permeases and the *traR* gene of the Ti plasmid pTiR10. *Mol Microbiol* 1996, **20**:1199-1210.
31. Fuqua C, Winans SC: Conserved cis-acting promoter elements are required for density-dependent transcription of *Agrobacterium tumefaciens* conjugal transfer genes. *J Bacteriol* 1996, **178**:435-440.
32. Gray KM, Passador L, Iglewski BH, Greenberg EP: Interchangeability and specificity of components from the quorum-sensing regulatory systems of *Vibrio fischeri* and *Pseudomonas aeruginosa*. *J Bacteriol* 1994, **176**:3076-3080.
33. Rust L, Pesci EC, Iglewski BH: Analysis of the *Pseudomonas aeruginosa* elastase (*lasB*) regulatory region. *J Bacteriol* 1996, **178**:1134-1140.
34. Stevens AM, Dolan KM, Greenberg EP: Synergistic binding of the *Vibrio fischeri* LuxR transcriptional activator domain and RNA polymerase to the *lux* promoter region. *Proc Natl Acad Sci USA* 1994, **91**:12619-12623.
35. Stevens AM, Greenberg EP: Quorum sensing in *Vibrio fischeri*: essential elements for activation of the luciferase genes. *J Bacteriol* 1997, **179**:557-562.
36. You Z, Fukushima J, Ishiwata T, Chang B, Kurata M, Kawamoto S, Williams P, Okuda K: Purification and characterization of LasR as a DNA-binding protein. *FEMS Microbiol Lett* 1996, **142**:301-307.
37. Beck Von Bodman S, Farrand SK: Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia stewartii* require induction by an *N*-acylhomoserine lactone autoinducer. *J Bacteriol* 1995, **177**:5000-5008.
38. Gambello MJ, Iglewski BH: Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *J Bacteriol* 1991, **173**:3000-3009.
39. Pearson JP, Gray KM, Passador L, Tucker KD, Eberhard A, Iglewski BH, Greenberg EP: Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc Natl Acad Sci USA* 1994, **91**:197-201.
40. Gambello MJ, Kaye S, Iglewski BH: LasR of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (*apr*) and an enhancer of exotoxin A expression. *Infect Immun* 1993, **61**:1180-1184.
41. Toder DS, Gambello MJ, Iglewski BH: *Pseudomonas aeruginosa* LasA: a second elastase under the transcriptional control of LasR. *Mol Microbiol* 1991, **5**:2003-2010.
42. Latifi A, Foglino M, Tanaka K, Williams P, Lazdunski A: A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhlR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol Microbiol* 1996, **21**:1137-1146.
- This paper provides strong evidence that LasR and RhlR are involved in a complex regulatory network. LasR appears to regulate *rhlR* and other genes. RhlR activates the stationary phase sigma factor gene *rpoS* and other genes. This paper and reference [43**] are complementary.
43. Pesci EC, Pearson JP, Seed PC, Iglewski BH: Regulation of *las* and *rhl* quorum sensing in *Pseudomonas aeruginosa*. *J Bacteriol* 1997, **179**:3127-3132.
- The authors develop the view of hierarchical control of gene expression by LasR, and present a model for this control. LasR activates the expression of several genes including *rhlR*. RhlR regulates a second set of genes including the stationary phase sigma factor gene *rpoS*. See also [42**].
44. Latifi A, Winson MK, Foglino M, Bycroft BW, Stewart GSAB, Lazdunski A, Williams P: Multiple homologues of LuxR and LuxI control expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Mol Microbiol* 1995, **17**:333-343.
45. Pearson JP, Passador L, Iglewski BH, Greenberg EP: A second *N*-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 1995, **92**:1490-1494.
46. Ochsner UA, Koch AK, Reiser J: Isolation and characterization of a regulatory gene affecting rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. *J Bacteriol* 1994, **176**:2044-2054.
47. Ochsner UA, Reiser J: Autoinducer-mediated regulation of rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 1995, **92**:6424-6428.
48. Brint JM, Ohman DE: Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhlI, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *J Bacteriol* 1995, **177**:7155-7163.
49. Chapon-Herve V, Akrim M, Latifi A, Lazdunski A, Bally M: Regulation of the *xcp* secretion pathway by multiple quorum-sensing modulons in *Pseudomonas aeruginosa*. *Mol Microbiol* 1997, **24**:1169-1178.
- The findings presented in this paper considerably expand the influence of the *las* and *rhl* quorum sensors to components of the general secretion pathway and potentially those secretion substrates dependent on its function.
50. Stead P, Rudd BAM, Bradshaw H, Noble D, Dawson MJ: Induction of phenazine antibiotic biosynthesis in cultures of *Pseudomonas aeruginosa* by *L*-*N*-(3-oxohexanoyl)-homoserine lactone. *FEMS Microbiol Lett* 1996, **140**:15-22.
51. Hengge-Aronis R: Survival of hunger and stress: the role of *rpoS* in early stationary phase regulation in *Escherichia coli*. *Cell* 1993, **72**:165-168.
52. Hengge-Aronis R: Regulation of gene expression during entry into stationary phase. In *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Edited by Neidhardt FC. Washington DC: American Society for Microbiology Press; 1995:1497-1512.
53. McLean RJC, Whiteley M, Stickler DJ, Fuqua WC: Evidence of autoinducer activity in naturally occurring biofilms. *FEMS Microbiol Lett* 1997, **154**:259-263.
54. Davis DG, Paresk MR, Pearson JP, Iglewski BH, Costerton JW, Greenberg EP: The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science* 1998, in press.
55. Albus AM, Pesci EC, Runyen-Janecky LJ, West SEH, Iglewski BH: Vfr controls quorum sensing in *Pseudomonas aeruginosa*. *J Bacteriol* 1997, **179**:3928-3935.
56. West SEH, Sample AK, Runyen-Janecky LJ: The *vfr* gene product, required for *Pseudomonas aeruginosa* exotoxin A and protease production, belongs to the cyclic AMP receptor protein family. *J Bacteriol* 1994, **176**:7532-7542.
57. Reimann C, Beyeler M, Latifi A, Winteler H, Foglino M, Lazdunski A, Haas D: The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer *N*-butyryl-homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol Microbiol* 1997, **24**:309-319.
58. Rahme LG, Stevens EJ, Wolford SF, Shao J, Tompkins RG, Ausubel FM: Common virulence factors for bacterial pathogenicity in plants and animals. *Science* 1995, **268**:1899-1902.
59. Wood DW, Pierson LS: The *phzI* gene of *Pseudomonas aureofaciens* 30-84 is responsible for the production of a diffusible signal required for phenazine antibiotic production. *Gene* 1996, **168**:49-53.