

DRAFT: NCT

RA

2021-04-02

1 NCT models

Abbreviations: ODE = ordinary differential equations.

1.1 GSR’03 model of NCT

Ran gradient. First we implement the “minimal Ran gradient system” from [GSR03]. The equations are shown in Table 1 and the constants are collected in Table 2. The “dynamic capacity” Ex is an optional maximal steady-state (positive) flux of nuclear $\text{Ran} \cdot \text{GTP}$ to cytoplasmic $\text{Ran} \cdot \text{GDP}$, which we determine using the additional equation

$$\frac{d}{dt} \text{Ex} = k_{\text{Ex}} [\text{Ran} \cdot \text{GTP}]_{\text{nuc}}, \quad k_{\text{Ex}} := 10 \text{ s}^{-2}, \quad \text{initial } \text{Ex} := 0 \text{ } \mu\text{M s}^{-1}. \quad (1)$$

The fluxes are in units of concentration/time ($\mu\text{M s}^{-1}$). The ones across the nuclear boundary have positive sign when exiting the nucleus and are normalized to the nuclear volume. The *amount* exiting the nucleus per unit of time is $\text{flux} \times V_{\text{nuc}}$.


Simulating the ODE across the scenarios of [GSR03] we obtain results that are sufficiently close to the original, see Table 3. Importantly, a 1000-fold nuclear enrichment of $\text{Ran} \cdot \text{GTP}$ is sustained in steady-state.

Code: [d56d16f/code/20210225-GSR/v1](#)

Coupling to transport. A coupling of the Ran gradient to importin–cargo transport was proposed in [GSR03, Fig. 6A]. We formulate a version of it in Table 4.

With the constants from Table 5, the steady-state of the model (reached after some $10 \times 10^4 \text{ s}$) is reported in Fig. 1. Nuclear accumulation of free cargo is over 20-fold. Sensitivity analysis shows that, in relative terms, the final nuclear concentration of free cargo depends most strongly on k_{knockoff} (and the volume of the nucleus). In particular, doubling k_{knockoff} almost doubles the concentration.

Code: [2a2199d/code/20210225-GSR/v2](#)



2

The following account for the cytoplasmic species. Here, [...] abbreviates the (cytoplasmic) concentration of the complex $\text{RanBP1} \cdot \text{Ran} \cdot \text{GTP}$.

$$\frac{d}{dt}[\text{Ran} \cdot \text{GDP}]_{\text{cyt}} = F_{\text{Ran} \cdot \text{GDP}} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} + \text{GAP} + \text{GAP}_{\text{RanBP1}} + \text{Ex} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} \quad (2a)$$

$$\frac{d}{dt}[\text{Ran} \cdot \text{GTP}]_{\text{cyt}} = F_{\text{Ran} \cdot \text{GTP}} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} - \text{GAP} - k_{\text{on}}^{\text{rbp}}[\text{RanBP1}][\text{Ran} \cdot \text{GTP}]_{\text{cyt}} + k_{\text{off}}^{\text{rbp}}[\dots] \quad (2b)$$

$$\frac{d}{dt}[\text{RanBP1} \cdot \text{Ran} \cdot \text{GTP}] = -\text{GAP}_{\text{RanBP1}} + k_{\text{on}}^{\text{rbp}}[\text{RanBP1}][\text{Ran} \cdot \text{GTP}]_{\text{cyt}} - k_{\text{off}}^{\text{rbp}}[\dots] \quad (2c)$$

The following account for the nuclear species. Following [GSR03], E denotes free RCC1.

$$\frac{d}{dt}[\text{Ran} \cdot \text{GDP}]_{\text{nuc}} = -F_{\text{Ran} \cdot \text{GDP}} + r_8[\text{IntC}] - r_1[\text{E}][\text{Ran} \cdot \text{GDP}]_{\text{nuc}} \quad (3a)$$

$$\frac{d}{dt}[\text{Ran} \cdot \text{GTP}]_{\text{nuc}} = -F_{\text{Ran} \cdot \text{GTP}} + r_4[\text{IntA}] - r_5[\text{E}][\text{Ran} \cdot \text{GTP}]_{\text{nuc}} - \text{Ex} \quad (3b)$$

The nucleotide-exchange reaction $\text{Ran} \cdot \text{GDP} + \text{GTP} \rightleftharpoons \text{Ran} \cdot \text{GTP} + \text{GDP}$ is catalyzed by RCC1. It is modeled as in [Kle+95, Fig. 6] / [GSR03, Fig. 1] with three intermediates. Note that it depends on the availability of GDP and GTP.

$$\frac{d}{dt}[\text{IntA}] = -(r_4 + r_6)[\text{IntA}] + r_5[\text{E}][\text{Ran} \cdot \text{GTP}]_{\text{nuc}} + r_3[\text{GTP}][\text{IntB}] \quad (4a)$$

$$\frac{d}{dt}[\text{IntB}] = r_6[\text{IntA}] + r_2[\text{IntC}] - (r_3[\text{GTP}] + r_7[\text{GDP}])[\text{IntB}] \quad (4b)$$

$$\frac{d}{dt}[\text{IntC}] = -(r_2 + r_8)[\text{IntC}] + r_1[\text{E}][\text{Ran} \cdot \text{GDP}]_{\text{nuc}} + r_7[\text{GDP}][\text{IntB}] \quad (4c)$$

Constraints on the total concentration:

$$\text{Free RCC1 :} \quad [\text{E}] = \text{RCC1}_{\text{total}} - ([\text{IntA}] + [\text{IntB}] + [\text{IntC}]) \quad (5a)$$

$$\text{Free RanBP1 :} \quad [\text{RanBP1}] = \text{RanBP1}_{\text{total}} - [\text{RanBP1} \cdot \text{Ran} \cdot \text{GTP}] \quad (5b)$$

Gradient-driven fluxes from the nucleus to the cytoplasm:

$$F_{\text{Ran} \cdot \text{GTP}} = D_{\text{Ran} \cdot \text{GTP}} ([\text{Ran} \cdot \text{GTP}]_{\text{nuc}} - [\text{Ran} \cdot \text{GTP}]_{\text{cyt}}) \quad (6a)$$

$$F_{\text{Ran} \cdot \text{GDP}} = D_{\text{Ran} \cdot \text{GDP}} ([\text{Ran} \cdot \text{GDP}]_{\text{nuc}} - [\text{Ran} \cdot \text{GDP}]_{\text{cyt}}) \quad (6b)$$

RanGAP hydrolyzes the γ -phosphate of $\text{Ran} \cdot \text{GTP}$. This is more efficient when $\text{Ran} \cdot \text{GTP}$ is bound to RanBP1 [Bis+95], reducing the IC50 seven-fold [GSR03, Table I, p. 1091].

$$\text{GAP} = k_{\text{GAP}}[\text{RanGAP}]/(1 + K_{\text{GAP}}/[\text{Ran} \cdot \text{GTP}]_{\text{cyt}}) \quad (7a)$$

$$\text{GAP}_{\text{RanBP1}} = k'_{\text{GAP}}[\text{RanGAP}]/(1 + K'_{\text{GAP}}/[\text{RanBP1} \cdot \text{Ran} \cdot \text{GTP}]) \quad (7b)$$

Table 1: The minimal Ran gradient system from [GSR03, Fig. 2]. Ex is an additional potentially useful flux of nuclear $\text{Ran} \cdot \text{GTP}$ to cytoplasmic $\text{Ran} \cdot \text{GDP}$, set by default to zero.

(2a)	$V_{\text{nuc}} = 1.2 \text{ pl}, \quad V_{\text{cyt}} = 1.8 \text{ pl}$	[GSR03, Table II]
(2a)	initial condition $[\text{Ran} \cdot \text{GDP}]_{\text{cyt}} = 5 \mu\text{M}$	[GSR03, Table II]
(2b)–(2c)	$k_{\text{on}}^{\text{rbp}} = 0.3 \mu\text{M}^{-1} \text{ s}^{-1}, \quad k_{\text{off}}^{\text{rbp}} = 4 \times 10^{-4} \text{ s}^{-1}$	[GSR03, Supp. Table A]
(3a)–(4c)	$r_1 = 74 \mu\text{M}^{-1} \text{ s}^{-1}, \quad r_8 = 55 \text{ s}^{-1}$ $r_7 = 11 \mu\text{M}^{-1} \text{ s}^{-1}, \quad r_2 = 21 \text{ s}^{-1}$ $r_3 = 0.6 \mu\text{M}^{-1} \text{ s}^{-1}, \quad r_6 = 19 \text{ s}^{-1}$ $r_5 = 100 \mu\text{M}^{-1} \text{ s}^{-1}, \quad r_4 = 55 \text{ s}^{-1}$	[GSR03, Supp. Table A] [Kle+95, Fig. 6]
(4a)–(4c)	$[\text{GTP}] = 500 \mu\text{M}, \quad [\text{GDP}] = 1.6 \mu\text{M}$	[GSR03, Table II]
(5a)	$\text{RCC1}_{\text{total}} = 0.7 \mu\text{M}$	[GSR03, Supp. Table B]
(5b)	$\text{RanBP1}_{\text{total}} = 2 \mu\text{M}$	[GSR03, Fig. 4]
(6a)	$D_{\text{Ran} \cdot \text{GTP}} = 0.03 \text{ s}^{-1}$	[GSR03, Table II]
(6b)	$D_{\text{Ran} \cdot \text{GDP}} = 0.12 \text{ s}^{-1}$	
(7a)	$k_{\text{GAP}} = 10.6 \text{ s}^{-1}, \quad K_{\text{GAP}} = 0.7 \mu\text{M}$	[GSR03, Supp. Table A]
(7b)	$k'_{\text{GAP}} = 10.8 \text{ s}^{-1}, \quad K'_{\text{GAP}} = 0.1 \mu\text{M}$	[GSR03, Table I]
(7a)–(7b)	cytoplasmic $[\text{RanGAP}] = 0.7 \mu\text{M}$	[GSR03, Table II / ST B]

Table 2: Constants for the “standard simulation condition” of §1.1 at 25 °C. Except for (2a), all species are initialized to zero at $t = 0$.

Condition	Affected parameters	Nuclear RanGTP, μM	Cytoplasmic RanGTP, nM	Dynamic capacity, $\mu\text{M/s}$
“Standard”	See Table 2	4.26 (4.3)	7.75 (7.7)	0.59 (0.60)
Omission of RanBP1	$\text{RanBP1}_{\text{total}} := 0$	4.27 (4.3)	8.13 (8.1)	0.59 (0.60)
200% RCC1	$\text{RCC1}_{\text{total}}$	3.95 (4.0)	7.17 (7.1)	0.59 (0.60)
50% RCC1	$\text{RCC1}_{\text{total}}$	4.31 (4.3)	7.82 (7.7)	0.58 (0.60)
10% RCC1	$\text{RCC1}_{\text{total}}$	3.59 (3.6)	6.50 (6.4)	0.46 (0.48)
1% RCC1	$\text{RCC1}_{\text{total}}$	1.40 (1.4)	2.52 (2.5)	0.075 (0.08)
GTP:GDP = 500:0	$[\text{GDP}] := 0 \mu\text{M}$	4.80 (4.8)	8.72 (8.6)	0.59 (0.60)
GTP:GDP = 500:50	$[\text{GDP}] := \frac{1}{10}[\text{GTP}]$	0.98 (0.8)	1.76 (1.5)	0.57 (0.58)
GTP:GDP = 500:500	$[\text{GDP}] := [\text{GTP}]$	0.12 (0.12)	0.22 (0.21)	0.34 (0.34)
Saturating NTF2	$D_{\text{Ran} \cdot \text{GDP}} := 0.48 \text{ s}^{-1}$	5.12 (5.1)	9.32 (9.2)	2.18 (2.2)
No NTF2	$D_{\text{Ran} \cdot \text{GDP}} := D_{\text{Ran} \cdot \text{GTP}}$	2.55 (2.5)	4.60 (4.5)	0.15 (0.16)
200% RanGAP	$[\text{RanGAP}]$	4.27 (4.3)	3.95 (3.9)	0.59 (0.60)
50% RanGAP	$[\text{RanGAP}]$	4.26 (4.3)	14.9 (14)	0.59 (0.60)
50% permeability	$D_{\text{Ran} \cdot \text{GTP}}$	4.91 (4.9)	4.44 (4.4)	0.59 (–)
200% permeability	$D_{\text{Ran} \cdot \text{GTP}}$	3.41 (3.4)	12.4 (12.3)	0.59 (–)
400% permeability	$D_{\text{Ran} \cdot \text{GTP}}$	2.46 (2.5)	18.0 (17.8)	0.59 (–)

Table 3: Steady-state concentrations for the simulation scenarios from [GSR03, Table II/III], with their results shown in brackets. Value for $D_{\text{Ran} \cdot \text{GDP}}$ is from [GSR03, Fig. 3].

The following equations comprise the handling of cargo by $\text{Imp}\beta$ in the cytoplasm.

$$R_{\text{cyt}} := -k_{\text{on}}^{\text{R}}[\text{Imp}\beta][\text{Ran} \cdot \text{GTP}]_{\text{cyt}} + k_{\text{off}}^{\text{R}}[\text{Imp}\beta \cdot \text{Ran} \cdot \text{GTP}]_{\text{cyt}} \quad (8a)$$

$$C_{\text{cyt}} := -k_{\text{on}}^{\text{C}}[\text{Imp}\beta][\text{Cargo}]_{\text{cyt}} + k_{\text{off}}^{\text{C}}[\text{Imp}\beta \cdot \text{Cargo}]_{\text{cyt}} \quad (8b)$$

$$\frac{d}{dt}[\text{Imp}\beta \cdot \text{Ran} \cdot \text{GTP}]_{\text{cyt}} = -R_{\text{cyt}} + F_{\text{Imp}\beta \cdot \text{Ran} \cdot \text{GTP}} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} - \text{GAP}_{\text{Imp}\beta} + \text{Knockoff}_{\text{cyt}} \quad (8c)$$

$$\frac{d}{dt}[\text{Imp}\beta]_{\text{cyt}} = +R_{\text{cyt}} + C_{\text{cyt}} + F_{\text{Imp}\beta} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} + \text{GAP}_{\text{Imp}\beta} \quad (8d)$$

$$\frac{d}{dt}[\text{Imp}\beta \cdot \text{Cargo}]_{\text{cyt}} = -C_{\text{cyt}} + F_{\text{Imp}\beta \cdot \text{Cargo}} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} - \text{Knockoff}_{\text{cyt}} \quad (8e)$$

$$\frac{d}{dt}[\text{Cargo}]_{\text{cyt}} = +C_{\text{cyt}} + F_{\text{Cargo}} \frac{V_{\text{nuc}}}{V_{\text{cyt}}} + \text{Knockoff}_{\text{cyt}} \quad (8f)$$

The flux of the reaction



is called **Knockoff**. It is modeled as a one-way reaction with forward rate k_{knockoff} . The previous equations are modified accordingly:

$$\frac{d}{dt}[\text{Ran} \cdot \text{GDP}]_{\text{cyt}} = (2a) + \text{GAP}_{\text{Imp}\beta} \quad (2a')$$

$$\frac{d}{dt}[\text{Ran} \cdot \text{GTP}]_{\text{cyt}} = (2b) + R_{\text{cyt}} - \text{Knockoff}_{\text{cyt}} \quad (2b')$$

Analogous nuclear equations (without **GAP**) are implemented but are omitted here. Analogously to (6a)/(6b) we have the additional nuclear-to-cytoplasmic diffusion fluxes

$$F_{\text{Imp}\beta \cdot \text{Ran} \cdot \text{GTP}}, \quad F_{\text{Imp}\beta}, \quad F_{\text{Imp}\beta \cdot \text{Cargo}}, \quad F_{\text{Cargo}}. \quad (10)$$

Table 4: Equations for the coupling of the minimal Ran gradient system from §1.1 to importin-mediated cargo transport.

(8a)	$k_{\text{on}}^{\text{R}} = 0.096 \mu\text{M}^{-1} \text{s}^{-1}, \quad k_{\text{off}}^{\text{R}} = 4.8 \times 10^{-6} \text{s}^{-1}$	[GSR03, Supp. Table A], [RM05, Table II]
(8b)	$k_{\text{on}}^{\text{C}} = 0.49 \mu\text{M}^{-1} \text{s}^{-1}, \quad k_{\text{off}}^{\text{C}} = 0.017 \text{s}^{-1}$	[Cat+01, below Fig. 3], [RM05, Table II]
(9)	$k_{\text{knockoff}} = 2 \times 10^{-2} \mu\text{M}^{-1} \text{s}^{-1}$	[RM05, Table II]
(10)	$D_{\text{Imp}\beta \cdot \text{Ran} \cdot \text{GTP}} = 0.07 \text{s}^{-1}, \quad D_{\text{Imp}\beta} = 0.4 \text{s}^{-1}$ $D_{\text{Imp}\beta \cdot \text{Cargo}} = 0.25 \text{s}^{-1}, \quad D_{\text{Cargo}} = 5 \times 10^{-4} \text{s}^{-1}$	[RM05, Table III]

Table 5: Constants for the $\text{Imp}\beta$ -mediated transport from §1.1 / Table 4.

References

- [Bis+95] F. R. Bischoff, H. Krebber, E. Smirnova, W. Dong, and H. Ponstingl. “Co-activation of RanGTPase and inhibition of GTP dissociation by Ran–GTP binding protein RanBP1”. In: *The EMBO Journal* 14.4 (Feb. 1995), pp. 705–715. DOI: [10.1002/j.1460-2075.1995.tb07049.x](https://doi.org/10.1002/j.1460-2075.1995.tb07049.x) (cit. on p. 3).
- [Kle+95] C. Klebe, H. Prinz, A. Wittinghofer, and R. S. Goody. “The Kinetic Mechanism of Ran-Nucleotide Exchange Catalyzed by RCC1”. In: *Biochemistry* 34.39 (Oct. 1995), pp. 12543–12552. DOI: [10.1021/bi00039a008](https://doi.org/10.1021/bi00039a008) (cit. on pp. 3, 4).
- [Cat+01] B. Catimel, T. Teh, M. R. Fontes, I. G. Jennings, D. A. Jans, G. J. Howlett, E. C. Nice, and B. Kobe. “Biophysical Characterization of Interactions Involving Importin- α during Nuclear Import”. In: *Journal of Biological Chemistry* 276.36 (Sept. 2001), pp. 34189–34198. DOI: [10.1074/jbc.m103531200](https://doi.org/10.1074/jbc.m103531200) (cit. on pp. 2, 5, 6).
- [GSR03] D. Görlich, M. J. Seewald, and K. Ribbeck. “Characterization of Ran-driven cargo transport and the RanGTPase system by kinetic measurements and computer simulation”. In: *The EMBO Journal* 22.5 (Mar. 2003), pp. 1088–1100. DOI: [10.1093/emboj/cdg113](https://doi.org/10.1093/emboj/cdg113) (cit. on pp. 1, 3–5).
- [RM05] G. Riddick and I. G. Macara. “A systems analysis of importin- α – β mediated nuclear protein import”. In: *Journal of Cell Biology* 168.7 (Mar. 2005), pp. 1027–1038. DOI: [10.1083/jcb.200409024](https://doi.org/10.1083/jcb.200409024) (cit. on pp. 2, 5, 6).

TODOs:

1. p.1. ? [Cat+01] and [RM05] discuss the reaction $\text{Imp}\beta \cdot \text{Cargo} \rightleftharpoons \text{Imp}\beta^* \cdot \text{Cargo}$