

Numerical Linear Algebra

1 Matrices, Vectors, and Norms

1.1 Matrix-Vector Multiplication

Given a matrix $A \in \mathbb{C}^{m \times n}$ and a vector $x \in \mathbb{C}^n$, the matrix-vector product $Ax = b \in \mathbb{C}^m$ is defined as:

$$b_i = \sum_{j=1}^n a_{ij}x_j, \quad \text{for } i = 1, \dots, m$$

where a_{ij} are the entries of the matrix A .

Observation

The transformation $x \mapsto Ax$ is a linear transformation from \mathbb{C}^n to \mathbb{C}^m , i.e., it satisfies:

$$A(x + y) = Ax + Ay, \quad A(\alpha x) = \alpha Ax, \quad \text{for all } x, y \in \mathbb{C}^n, \alpha \in \mathbb{C}$$

1.1.1 A Matrix times a Vector

$$b = Ax = \sum_{j=1}^n x_j a_j$$

where a_j is the j -th column of A .

$$\begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

Example: Vandermonde Matrix

A Vandermonde matrix is defined as:

$$A = \begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_m & x_m^2 & \cdots & x_m^{n-1} \end{bmatrix} \in \mathbb{C}^{m \times n}$$

Observation

Let fix the sequence $\{x_1, x_2, \dots, x_m\}$. If p and q are polynomials of degree at most $n - 1$ and α is a scalar, then:

1. $(p + q)$ is a polynomial of degree at most $n - 1$, and so are $\alpha p, \alpha q$.
2. $(p + q)(x_i) = p(x_i) + q(x_i)$ for $i = 1, \dots, m$.
3. $(\alpha p)(x_i) = \alpha p(x_i)$ for $i = 1, \dots, m$.

Observation

Suppose we have a vector $c \in \mathbb{C}^n$ representing the coefficients of a polynomial $p(x) = c_0 + c_1x + c_2x^2 + \cdots + c_{n-1}x^{n-1}$. Then

$$p(x_i) = (Ac)_i = c_0 + c_1x_i + c_2x_i^2 + \cdots + c_{n-1}x_i^{n-1}$$

Any polynomial of degree at most $n - 1$ can be represented as

$$p(x) = \begin{bmatrix} 1 & x & x^2 & \cdots & x^{n-1} \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ c_2 \\ \vdots \\ c_{n-1} \end{bmatrix}$$

1.2 Matrix-Matrix Multiplication

Given matrices $A \in \mathbb{C}^{l \times m}$ and $C \in \mathbb{C}^{m \times n}$, the matrix-matrix product $B = AC \in \mathbb{C}^{l \times n}$ is defined as:

$$b_{ij} = \sum_{k=1}^m a_{ik}c_{kj}, \quad \text{for } i = 1, \dots, l, j = 1, \dots, n$$

So,

$$B = \begin{bmatrix} b_1 & b_2 & \cdots & b_n \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & \cdots & a_m \end{bmatrix} \begin{bmatrix} c_1 & c_2 & \cdots & c_n \end{bmatrix}$$

$$b_j = Ac_j = \sum_{k=1}^m c_{kj}a_k, \quad j = 1, \dots, n$$

Example: Outer Product

Given two vectors $u \in \mathbb{C}^{m \times 1}$ and $v \in \mathbb{C}^{1 \times n}$, the outer product $uv^T \in \mathbb{C}^{m \times n}$ is defined as:

$$\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_m \end{bmatrix} \begin{bmatrix} v_1 & v_2 & \cdots & v_n \end{bmatrix} = \begin{bmatrix} u_1v_1 & u_1v_2 & \cdots & u_1v_n \\ u_2v_1 & u_2v_2 & \cdots & u_2v_n \\ \vdots & \vdots & \ddots & \vdots \\ u_mv_1 & u_mv_2 & \cdots & u_mv_n \end{bmatrix}$$

Example

Let $B \in \mathbb{C}^{m \times n}$, let a_1, a_2, \dots, a_n be the columns of $A \in \mathbb{C}^{m \times n}$ and let $R \in \mathbb{C}^{n \times n}$ be an upper triangular matrix with all its superdiagonal entries equal to 1 such that

$$B = \begin{bmatrix} b_1 & b_2 & \cdots & b_n \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix} \begin{bmatrix} 1 & 1 & \cdots & 1 \\ 0 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Then,

$$b_j = Ar_j = \sum_{k=1}^j a_k, \quad j = 1, \dots, n$$

which is known as the indefinite integral operation.

1.3 Range and Null Space

Theorem

The range of a matrix $A \in \mathbb{C}^{m \times n}$ is the space spanned by its columns, i.e. it's column space.

"The set of vectors that can be written as Ax for some $x \in \mathbb{C}^n$."

1.3.1 Null Space and Rank

The null space of a matrix $A \in \mathbb{C}^{m \times n}$ is defined as:

$$\text{span}\{x \in \mathbb{C}^n : Ax = 0\}$$

The rank of a matrix $A \in \mathbb{C}^{m \times n}$ is defined as the dimension of its range:

$$\text{rank}(A) = \dim(\text{range}(A))$$

Theorem

For any matrix $A \in \mathbb{C}^{m \times n}$, $m \geq n$, A has full rank if and only if it maps no two distinct vectors to the same vector:

$$Ax = Ay \implies x = y, \forall x, y \in \mathbb{C}^n \quad \text{and} \quad Ax \neq Ay \implies x \neq y, \forall x, y \in \mathbb{C}^n$$

1.3.2 Non-singular Matrices

A non-singular matrix, or invertible matrix is a square matrix $A \in \mathbb{C}^{n \times n}$ that has full rank.

1.3.3 Inverse of a Matrix

The matrix $A \in \mathbb{C}^{n \times n}$ is the inverse of Z if and only if:

$$AZ = ZA = I = [e_1 \ e_2 \ \cdots \ e_n]$$

where e_i is the i -th standard basis vector. Furthermore, the inverse of a matrix is unique and

$$\text{rank}(A) = n = \text{rank}(Z)$$

Theorem

For $A \in \mathbb{C}^{n \times n}$, the following statements are equivalent:

1. A has an inverse A^{-1} .
2. A has full rank, i.e. $\text{rank}(A) = n$.
3. The columns of A span \mathbb{C}^n , i.e. $\text{range}(A) = \mathbb{C}^n$.
4. The columns of A are linearly independent, i.e. $\text{null}(A) = \{0\}$.
5. 0 is not an eigenvalue of A .
6. 0 is not a singular value of A .
7. $|A| \neq 0$.

1.4 Orthogonal Vectors and Matrices

1.4.1 Complex Conjugate and Conjugate Transpose

Given $z \in \mathbb{C}$, the complex conjugate of z is denoted by \bar{z} or z^* , where if $z = x + iy$, then $\bar{z} = x - iy$.

The conjugate transpose of a matrix $A \in \mathbb{C}^{m \times n}$ is denoted by A^* , where $A^* = \bar{A}^T$.

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \implies A^* = \begin{bmatrix} \bar{a}_{11} & \bar{a}_{21} & \bar{a}_{31} \\ \bar{a}_{12} & \bar{a}_{22} & \bar{a}_{32} \end{bmatrix}$$

1.4.2 Hermitian Matrices and Skew-Hermitian Matrices

A matrix $A \in \mathbb{C}^{n \times n}$ is called Hermitian if $A^* = A$. (Or $A = A^T$ if $A \in \mathbb{R}^{n \times n}$.)

A matrix $A \in \mathbb{C}^{n \times n}$ is called Skew-Hermitian if $A^* = -A$. (Or $A = -A^T$ if $A \in \mathbb{R}^{n \times n}$.)

1.4.3 Inner Product and Norm

Given two vectors $x, y \in \mathbb{C}^n$, the inner product is defined as:

$$\langle x, y \rangle = y^* x = \sum_{i=1}^n \bar{y}_i x_i$$

The norm of a vector $x \in \mathbb{C}^n$ is defined as:

$$\|x\| = \sqrt{\langle x, x \rangle} = \sqrt{\sum_{i=1}^n |x_i|^2}$$

The modulus of a complex number $z \in \mathbb{C}$ is defined as:

$$|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2} \quad \text{where } z = x + iy$$

The angle θ between two vectors $x, y \in \mathbb{C}^n$ is defined as:

$$\cos(\theta) = \frac{\langle x, y \rangle}{\|x\| \|y\|}$$

Observation

The inner product is bilinear, which means:

1. $\langle x_1 + x_2, y \rangle = \langle x_1, y \rangle + \langle x_2, y \rangle$
2. $\langle x, y_1 + y_2 \rangle = \langle x, y_1 \rangle + \langle x, y_2 \rangle$
3. $\langle \alpha x, \beta y \rangle = \alpha \bar{\beta} \langle x, y \rangle$ for all $\alpha, \beta \in \mathbb{C}$

For all vectors or matrices A, B of compatible dimensions, we have:

$$(AB)^* = B^* A^*$$

1.4.4 Orthogonal Vectors

Two vectors $x, y \in \mathbb{C}^n$ are orthogonal if $\langle x, y \rangle = 0$.

Observation

If x and y are orthogonal, then they are perpendicular.

With $x = \{x_1, x_2, \dots, x_n\}$ and $y = \{y_1, y_2, \dots, y_n\}$, x and y are orthogonal if and only if:

$$\langle x_i, y_j \rangle = 0, \quad \forall i, j = 1, 2, \dots, n$$

1.4.5 Orthonormal Vectors

A set of vectors $\{x_1, x_2, \dots, x_n\} \subset \mathbb{C}^n$ is orthonormal if:

$$\langle x_i, x_j \rangle = 0 \text{ for } i \neq j \quad \text{and} \quad \langle x_i, x_i \rangle = \|x_i\|^2 = 1$$

Observation

The vectors in an orthogonal set are linearly independent.

1.4.6 Orthogonal and Unitary Matrices

A matrix $Q \in \mathbb{R}^{n \times n}$ is orthogonal if its columns form an orthonormal set, i.e. $Q^T Q = Q Q^T = I$.

A matrix $U \in \mathbb{C}^{n \times n}$ is unitary if its columns form an orthonormal set, i.e. $U^* U = U U^* = I$.

Observation

If $U \in \mathbb{C}^{n \times n}$ is unitary, then $U^{-1} = U^*$ and

$$\begin{bmatrix} u_1^* \\ u_2^* \\ \vdots \\ u_n^* \end{bmatrix} \begin{bmatrix} u_1 & u_2 & \cdots & u_n \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

which implies that $\langle u_i, u_j \rangle = \delta_{ij}$ where δ_{ij} is the Kronecker delta, i.e. $\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$.

2 Vector Spaces

A vector space \mathcal{V} over a field \mathbb{F} (usually \mathbb{R} or \mathbb{C}) is a set of objects called vectors, along with two operations: vector addition and scalar multiplication, satisfying the following axioms for all $u, v, w \in \mathcal{V}$ and $\alpha, \beta \in \mathbb{F}$:

1. (Closure under addition) $u + v \in \mathcal{V}$
2. (Commutativity) $u + v = v + u$
3. (Associativity) $(u + v) + w = u + (v + w)$
4. (Existence of additive identity) There exists an element $0 \in \mathcal{V}$ such that $u + 0 = u$
5. (Existence of additive inverses) For each $u \in \mathcal{V}$, there exists an element $-u \in \mathcal{V}$ such that $u + (-u) = 0$
6. (Closure under scalar multiplication) $\alpha u \in \mathcal{V}$
7. (Distributivity of scalar multiplication with respect to vector addition) $\alpha(u + v) = \alpha u + \alpha v$
8. (Distributivity of scalar multiplication with respect to field addition) $(\alpha + \beta)u = \alpha u + \beta u$
9. (Associativity of scalar multiplication) $\alpha(\beta u) = (\alpha\beta)u$
10. (Existence of multiplicative identity) $1u = u$ where 1 is the multiplicative identity in \mathbb{F}

2.1 Linear Subspaces

A subset \mathcal{W} of a vector space \mathcal{V} is a subspace if \mathcal{W} is itself a vector space under the operations of addition and scalar multiplication defined on \mathcal{V} :

1. (Closure under addition) $u + v \in \mathcal{W}$
2. (Closure under scalar multiplication) $\alpha u \in \mathcal{W}$

2.2 Linear Independence

A set of vectors $\{v_1, v_2, \dots, v_k\} \subset \mathcal{V}$ is linearly independent if the only solution to

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_k v_k = 0$$

is $\alpha_1 = \alpha_2 = \dots = \alpha_k = 0$. If there exists a non-trivial solution, the set is linearly dependent.

2.3 Basis and Dimension

A basis of a vector space \mathcal{V} is a set of linearly independent vectors that spans \mathcal{V} . The dimension of \mathcal{V} , denoted $\dim(\mathcal{V})$, is the number of vectors in any basis of \mathcal{V} .

2.4 Inner Product and Norms

An inner product on a vector space \mathcal{V} over \mathbb{F} is a function $\langle \cdot, \cdot \rangle : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{F}$. The standard inner product on \mathbb{C}^n is defined as:

$$\langle x, y \rangle = y^* x = \sum_{i=1}^n \bar{y}_i x_i$$

A norm on a vector space \mathcal{V} is a function $\| \cdot \| : \mathcal{V} \rightarrow \mathbb{R}$ satisfying:



1. (Non-negativity) $\|v\| \geq 0$
2. (Definiteness) $\|v\| = 0$ if and only if $v = 0$
3. (Homogeneity) $\|\alpha v\| = |\alpha| \|v\|$ for all $\alpha \in \mathbb{F}$ and $v \in \mathcal{V}$
4. (Triangle inequality) $\|u + v\| \leq \|u\| + \|v\|$ for all $u, v \in \mathcal{V}$

The p -norm on \mathbb{C}^n is defined as:

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$$

for $1 \leq p < \infty$, and the ∞ -norm is defined as:

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$$



2.4.1 Lemma

Let $\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ be two norms on a finite-dimensional vector space \mathcal{V} . Then there exist positive constants c_1 and c_2 such that for all $v \in \mathcal{V}$:

$$c_1\|v\|_\alpha \leq \|v\|_\beta \leq c_2\|v\|_\alpha$$

$\|\cdot\|_\alpha$ and $\|\cdot\|_\beta$ are said to be equivalent norms. In \mathbb{R}^n :

$$\|x\|_2 \leq \|x\|_1 \leq \sqrt{n}\|x\|_2$$

$$\|x\|_\infty \leq \|x\|_2 \leq \sqrt{n}\|x\|_\infty$$

$$\|x\|_\infty \leq \|x\|_1 \leq n\|x\|_\infty$$

3 Matrix Norms

A matrix norm is a function $\|\cdot\| : \mathbb{C}^{m \times n} \rightarrow \mathbb{R}$ satisfying:

1. (Non-negativity) $\|A\| \geq 0$
2. (Definiteness) $\|A\| = 0$ if and only if $A = 0$
3. (Homogeneity) $\|\alpha A\| = |\alpha|\|A\|$ for all $\alpha \in \mathbb{C}$ and $A \in \mathbb{C}^{m \times n}$
4. (Triangle inequality) $\|A + B\| \leq \|A\| + \|B\|$ for all $A, B \in \mathbb{C}^{m \times n}$
5. (Sub-multiplicativity) $\|AB\| \leq \|A\|\|B\|$ for all $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times p}$

3.1 Frobenius Norm

The Frobenius norm of a matrix $A \in \mathbb{C}^{m \times n}$ is defined as:

$$\|A\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2} = \sqrt{\text{trace}(A^*A)}$$

3.2 Induced Norms

The induced norm (or operator norm) of a matrix $A \in \mathbb{C}^{m \times n}$ is defined as:

$$\|A\| = \sup_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \sup_{\|x\|=1} \|Ax\|$$

3.3 Consistent Norms

Let $\|\cdot\|_{m \times n}$, $\|\cdot\|_{n \times p}$ and $\|\cdot\|_{m \times p}$ be three matrix norms on $\mathbb{C}^{m \times n}$, $\mathbb{C}^{n \times p}$ and $\mathbb{C}^{m \times p}$ respectively. They are said to be consistent if:

$$\|AB\|_{m \times p} \leq \|A\|_{m \times n} \|B\|_{n \times p}, \quad \forall A \in \mathbb{C}^{m \times n}, B \in \mathbb{C}^{n \times p}$$

3.4 Properties of Induced Norms

1. $\|Ax\| \leq \|A\|\|x\|$ for all $A \in \mathbb{C}^{m \times n}$ and $x \in \mathbb{C}^n$
2. $\|AB\| \leq \|A\|\|B\|$ for all $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times p}$
3. $\|A\|_\infty = \max_{1 \leq i \leq m} \sum_{j=1}^n |a_{ij}|$ (maximum absolute row sum)
4. $\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^m |a_{ij}|$ (maximum absolute column sum)
5. $\|A\|_2 = \sqrt{\lambda_{\max}(A^*A)}$ (with λ_{\max} the largest eigenvalue)
6. $\|A\|_2 = \|A^T\|_2$
7. $\|A\|_2 = \max |\lambda_i(A)|$ if A is normal (i.e. $A^*A = AA^*$)
8. $\|A\|_F = \sqrt{\text{trace}(A^*A)}$

So, in general,

$$\|A\|_\alpha = \max_{x \neq 0} \frac{\|Ax\|_\alpha}{\|x\|_\alpha} = \max_{\|x\|_\alpha=1} \|Ax\|_\alpha$$

The Frobenius norm is consistent with the euclidean vector norm, but it is not an induced norm.