

Relative Covariance

Nuno Carvalho

October 6, 2026

***Note:** Parts of this document may be AI-written. My general workflow is to write sections of the article myself, laying out the general points I want to hit on and going into detail where relevant. I then converse with an LLM to identify errors in the logic or math and opportunities to use clearer or more generalizable notation. I then have the LLM make the edits to reflect our proposed changes, including adding more detailed derivations.*

This article describes how the phenotypic covariance between two individuals i and j can be modeled by limited information we have about the pair. I derive formulas from the starting trait generative model and only introduce assumptions at the point the derivation requires it. I start with a simple trait architecture and iteratively add more complexity.

1 Additive genetics under random mating with independent environment

We begin with the simplest architecture: a trait determined by an additive genetic component and an independent environmental component, in a randomly mating population.

1.1 The phenotype model

For an individual i we write the (standardized) phenotype as

$$y_i = g_i + e_i, \tag{1}$$

where g_i is the genetic value and e_i the environmental deviation. We take y to be standardized across the population, $E[y] = 0$ and $\text{Var}[y] = 1$, and both components to be mean-centered, $E[g] = E[e] = 0$, with variances $\text{Var}[g] = V_A$ and $\text{Var}[e] = V_E$.

We then introduce the following assumptions.

[Additivity] Additive genetic value

The genetic value is a fixed additive function of the individual's own genotype,

$$g_i = \sum_{k=1}^M \beta_k x_{ik},$$

where β_k is the per-standardized-allele effect of variant k , x_{ik} is the standardized genotype ($E[x_{ik}] = 0$, $\text{Var}[x_{ik}] = 1$), and M is the number of variants in the genome. Non-causal variants are included with $\beta_k = 0$. Because we assume an additive model, the genetic value can also be called a *breeding value*, although we stick to the former term.

Commonly bundled into: the additive model; no dominance, no gene–gene interactions (GxG, epistasis), and no gene–environment interactions (GxE) — each would make the genetic value depend on more than the individual’s own genotype in an additive way.

[GE-indep] Gene–environment independence

An individual’s environmental component is independent of their own genetic component:

$$g_i \perp e_i$$

Commonly bundled into: no gene–environment correlation or interaction, within an individual.

[No-IGE] No indirect genetic effects

One individual’s genetic component is independent of another individual’s environmental component:

$$g_i \perp e_j \quad (i \neq j).$$

Commonly bundled into: no indirect (e.g. parental) genetic effects — one individual’s genotype does not shape another’s environment.

[Env-indep] Independent environments

The environmental component is independent across individuals, including relatives:

$$e_i \perp e_j \quad (i \neq j).$$

Commonly bundled into: no shared/common environment (C); no environmental confounder shared by the pair (e.g. population stratification).

Because $g_i \perp e_i$ ([GE-indep]), $\text{Cov}[g_i, e_i] = 0$ and the phenotypic variance decomposes as $\text{Var}[y] = V_A + V_E$. Narrow-sense heritability is typically defined as:

$$h^2 = \frac{V_A}{\text{Var}[y]}$$

Since $\text{Var}[y] = 1$, $h^2 = V_A$ under this model.

1.2 Phenotypic covariance is genotypic covariance

For a single fixed pair, y_i and y_j are two fixed values, so “their covariance” only has meaning as an average over an ensemble of pairs. The ensemble we use is the set of all pairs that share a given piece of *relatedness information* $\mathcal{I}_{ij}$ about the pair, and the object of interest is the conditional cross-product

$$C_{ij}(\mathcal{I}) := \mathbb{E}[y_i y_j \mid \mathcal{I}_{ij}]. \quad (2)$$

If $\mathbb{E}[y_i \mid \mathcal{I}] = 0$, then $C_{ij}(\mathcal{I}) = \text{Cov}[y_i, y_j \mid \mathcal{I}]$. For most information $\mathcal{I}$, like the realized relatedness or degree between two relatives, this equivalence is applicable. An exception is when the information we are modelling contains individuals’ actual genotype information, for which $\mathbb{E}[y_i \mid \text{genotypes}] = g_i \neq 0$, therefore the expected cross-product of two individuals is not equivalent to their covariance given the fixed information.

Using (1) together with [No-IGE] and [Env-indep], every cross term involving e vanishes in expectation (each factors through a mean-zero environmental term), so the phenotypic cross-product reduces to the genetic one:

$$C_{ij}(\mathcal{I}) = \mathbb{E}[(g_i + e_i)(g_j + e_j) \mid \mathcal{I}] = \mathbb{E}[g_i g_j \mid \mathcal{I}]. \quad (3)$$

The rest of the article evaluates (3) at four levels of resolution for $\mathcal{I}$.

1.3 Level 0 — the exact genetic cross-product

At the finest resolution we condition on the realized genotypes of i and j at the causal variants, for which we know the true causal effect sizes. Then g_i and g_j are fixed, and by [Additivity],

$$C_{ij} = \mathbb{E}[y_i y_j \mid \mathbf{x}_i, \mathbf{x}_j, \boldsymbol{\beta}] = g_i g_j = \sum_k \sum_{k'} \beta_k \beta_{k'} x_{ik} x_{jk'} \quad (4)$$

Under our model and assumptions so far, this returns the true expected cross-product of two individuals’ phenotypes. The genetic cross-product between two individuals is, fundamentally, an effect-size-weighted function of the genotypes they actually carry at causal sites. Notably, this formula does not require any knowledge of the relatedness of the two individuals.

1.4 Level 1 — genome-wide allelic sharing

If we do not know the causal effect sizes, then we cannot apply (4). We therefore adopt the framework whereby variant effects sizes are random effects. Because we do not have any variant-specific information for distinguishing variants apart, their priors are identical up to the first two moments:

[Zero-mean-effects] Effects are mean-zero

The expected effect of every variant is zero:

$$E[\beta_k] = 0.$$

Commonly bundled into: the infinitesimal / random-effects model; effects are as likely to raise the trait as to lower it.

[Exchangeable-effects] Every variant carries the same effect variance

Every variant has the same expected squared effect:

$$E[\beta_k^2] = E[\beta_{k'}^2].$$

Commonly bundled into: the infinitesimal / random-effects model; equal per-standardized-allele effect variance across variants.

[Uncorrelated-effects] Effects are uncorrelated across variants

The effects of two different variants have zero expected cross-product:

$$E[\beta_k \beta_{k'}] = 0 \quad (k \neq k').$$

Commonly bundled into: the infinitesimal / random-effects model, whose effects are drawn independently across variants.

In the following assumption, we preclude genotype information, a form of variant-specific information, from influencing our priors for variants' effect sizes up to the second moment. Assortative mating and population stratification violate this assumption by “leaking” information regarding the first two moments of variant effect sizes. We are effectively imposing a constraint on any non-random mating that is correlated with effect size distributions. Later, in **[Non-IBD-indep]**, we impose a maximal version of this constraint.

[Genotype-independent-effects] Observing genotypes does not change the effects' first two moments

Conditioning on the genotypes leaves the first and second moments of the effects unchanged:

$$E[\beta_k | \mathbf{X}] = E[\beta_k], \quad E[\beta_k \beta_{k'} | \mathbf{X}] = E[\beta_k \beta_{k'}] \quad \text{for all } k, k'.$$

Commonly bundled into: effects drawn independently of the genotypes; in particular, no selection and no assortative mating, both of which make the genotypes informative about the effects.

Taking the expectation of (4) over the effects:

$$\mathbb{E}[g_i g_j \mid \mathbf{x}_i, \mathbf{x}_j] = \sum_k \sum_{k'} \mathbb{E}[\beta_k \beta_{k'} \mid \mathbf{X}] x_{ik} x_{jk'} = \sum_k \sum_{k'} \mathbb{E}[\beta_k \beta_{k'}] x_{ik} x_{jk'},$$

the second step by [\[Genotype-independent-effects\]](#). By [\[Uncorrelated-effects\]](#), $\mathbb{E}[\beta_k \beta_{k'}] = 0$ whenever $k \neq k'$; the entire off-diagonal ($k \neq k'$) collapses regardless of any correlation between the genotypes. What remains is the diagonal, weighted by each variant’s effect variance, which by [\[Exchangeable-effects\]](#) is the same for every variant:

$$\mathbb{E}[g_i g_j \mid \mathbf{x}_i, \mathbf{x}_j] = \sum_k \mathbb{E}[\beta_k^2] x_{ik} x_{jk}.$$

The additive variance is the total of these: since $\text{Var}[x_{ik}] = 1$ and the off-diagonal effect terms vanish, $V_A = \text{Var}[g_i] = \sum_k \mathbb{E}[\beta_k^2]$, and thus $\mathbb{E}[\beta_k^2] = V_A/M$. Effectively, the per-variant effect weighting of (4) has been replaced by a uniform weighting across all variants. Therefore

$$C_{ij}(\mathbf{x}_i, \mathbf{x}_j) = V_A \cdot \frac{1}{M} \sum_k x_{ik} x_{jk}. \quad (5)$$

Unlike at Level 0, this cross-product is also a covariance: $\mathbb{E}[g_i \mid \mathbf{x}_i, \mathbf{x}_j] = \sum_k \mathbb{E}[\beta_k \mid \mathbf{X}] x_{ik} = 0$ by [\[Genotype-independent-effects\]](#) and [\[Zero-mean-effects\]](#). The quantity multiplying V_A is the genome-wide genotype cross-product (i.e. allelic sharing), which we’ll define by:

$$A_{ij} := \frac{1}{M} \sum_k x_{ik} x_{jk} \quad (6)$$

This can be thought of as an identity-by-state (IBS)-based measure of “relatedness”, and is exactly the (i, j) entry of what is typically computed as a Genetic Relatedness Matrix (GRM), $\mathbf{A}$. Therefore, we can further simplify:

$$\boxed{C_{ij}(A_{ij}) = V_A A_{ij}} \quad (7)$$

1.5 Level 2 — realized relatedness

The problem with calculating A_{ij} is that we may not have genotyped every causal variant that contributes to a trait. As long as some of the causal variants are not in perfect correlation with a genotyped variant, then the covariance of two individuals’ phenotypes will not fully take into account the contributions of the ungenotyped causal variants. However, when individuals are related, their IBS-based relatedness from genotyped variants is informative of the IBS-based relatedness of ungenotyped variants. Therefore, the former can be used to capture the contributions of the latter.

When individuals inherit a genomic segment from a common ancestor, then all of the variants within that genomic segment will be identical-by-state, as long as we assume that de-novo mutations have not caused the inherited alleles to be different. These genomic segments and their variants are said to be identical-by-descent (IBD).

[No-mutation] IBD implies IBS

Alleles that are identical-by-descent are also identical-by-state: descent is transmitted without mutation.

Commonly bundled into: the infinite-sites / no-recurrent-mutation assumption.

When two individuals' alleles are IBD, then they are always IBS; when they are not IBD, they can still be IBS. Therefore, we can decompose the IBS-based relatedness A_{ij} into a sum over IBD allele pairs and a sum over non-IBD allele pairs. Write each standardized genotype as the sum of its two per-allele (maternal and paternal) standardized values, $x_{ik} = z_{ik}^{(m)} + z_{ik}^{(p)}$. Since $\text{Var}[x_{ik}] = 1$ and the two alleles are independent ([No-inbreeding]), each per-allele value has $E[z] = 0$ and $\text{Var}[z] = \frac{1}{2}$. The genotype cross-product then expands over the four ordered allele pairs, and A_{ij} splits by IBD status:

$$\begin{aligned} A_{ij} &= \frac{1}{M} \sum_k x_{ik} x_{jk} = \frac{1}{M} \sum_k \sum_{u,v \in \{m,p\}} z_{ik}^{(u)} z_{jk}^{(v)} \\ &= \underbrace{\frac{1}{M} \sum_{\text{IBD pairs}} z_{ik}^{(u)} z_{jk}^{(v)}}_{\text{IBD}} + \underbrace{\frac{1}{M} \sum_{\text{non-IBD pairs}} z_{ik}^{(u)} z_{jk}^{(v)}}_{\text{non-IBD}}. \end{aligned} \quad (8)$$

For an IBD allele pair, the two per-allele values are copies of the same ancestral allele and, by [No-mutation], of the same state; they are therefore equal, so their expected cross-product is $E[(z_{ik}^{(u)})^2] = \text{Var}[z_{ik}^{(u)}] = \frac{1}{2}$. For non-IBD allele pairs, we have to make an assumption on how they are distributed.

[Non-IBD-indep] Independence of non-IBD alleles

Any two alleles that are *not* IBD are independent draws from the population allele distribution.

Commonly bundled into: between individuals: random mating, no population structure, and no assortative mating.

[No-inbreeding] No inbreeding

The maternally and paternally inherited alleles of an individual are not IBD with each other.

Commonly bundled into: a randomly mating, outbred population (Hardy–Weinberg within individuals).

Two consequences follow, and both are used throughout. First, an individual's own two alleles, not being IBD, are independent draws from the population by [Non-IBD-indep],

and so are uncorrelated with each other. A variant's genotype variance is therefore the sum of its two allele variances, $2p_k(1-p_k)$, which is what makes the standardized genotype x_{ik} have variance 1 and each of its two per-allele values variance $\frac{1}{2}$. Second, IBD is an equivalence relation on alleles — two are IBD exactly when they are copies of the same ancestral allele — and the assumption says no class may contain both alleles of one individual. At most two of the four ordered allele pairs between a pair of individuals can therefore be IBD, which is what keeps the number of alleles they share IBD at a variant in $\{0, 1, 2\}$.

By the definition of independent draws, the expected cross-product of a non-IBD allele pair is therefore $E[z_{ik}^{(u)}]E[z_{jk}^{(v)}] = 0$. Letting $\text{IBD}_{ij}(k) \in \{0, 1, 2\}$ denote the number of alleles the pair shares IBD at variant k , exactly $\text{IBD}_{ij}(k)$ of the four ordered allele pairs are IBD — each contributing $\frac{1}{2}$ — and the rest contribute 0, so

$$E[x_{ik}x_{jk} \mid \text{IBD}_{ij}(k)] = \frac{1}{2}\text{IBD}_{ij}(k).$$

A variant thus contributes expected cross-product 0, $\frac{1}{2}$, or 1 according to whether the pair shares 0, 1, or 2 alleles IBD there. We define the pair's *realized relatedness* π_{ij} as the average IBD state across variants, halved — equivalently, the fraction of the pair's genome that is IBD:

$$\pi_{ij} := \frac{1}{2M} \sum_k \text{IBD}_{ij}(k). \quad (9)$$

Conditioning on the pair's realized IBD states and using the per-variant identity $E[x_{ik}x_{jk} \mid \text{IBD}_{ij}(k)] = \frac{1}{2}\text{IBD}_{ij}(k)$, linearity of expectation gives the expected allelic sharing:

$$E[A_{ij} \mid \{\text{IBD}_{ij}(k)\}] = \frac{1}{M} \sum_k \frac{1}{2} \text{IBD}_{ij}(k) = \frac{1}{2M} \sum_k \text{IBD}_{ij}(k) = \pi_{ij}.$$

This depends on the IBD states only through π_{ij} , so we may condition on π_{ij} alone. Crucially, it required no assumption that the per-variant IBD states are independent — and indeed they are not: an IBD state changes only at a cross-over within a co-inherited segment, so physically linked variants have strongly correlated IBD states. Linearity of expectation is insensitive to this correlation.

Combining with (7), the pair's expected phenotypic cross-product is their realized relatedness scaled by the additive variance:

$$\boxed{E[y_i y_j \mid \pi_{ij}] = V_A E[A_{ij} \mid \pi_{ij}] = V_A \pi_{ij}}. \quad (10)$$

1.6 Level 3 — genetic degree

In cases where we cannot measure π_{ij} based on per-variant IBD states, we can still estimate it if we know the genetic degree d_{ij} between the individuals.

[Genetic-degree] Degree determines expected relatedness

The genetic degree d_{ij} fixes the distribution of IBD sharing through Mendelian segregation, so that the expected relatedness given the degree is a single number $\bar{\pi}_{d_{ij}}$:

$$E[\pi_{ij} \mid d_{ij}] = \bar{\pi}_{d_{ij}}.$$

Furthermore, under the outbreeding assumption ([No-inbreeding]) — applied here not only to i and j but to the entire pedigree connecting them, so that it contains no inbreeding loops — $\bar{\pi}_{d_{ij}} = 2^{-d_{ij}}$, which is not fully derived here. We can then obtain the expected phenotypic covariance given the degree d_{ij} :

$$E[y_i y_j \mid d_{ij}] = E[E[y_i y_j \mid \pi_{ij}] \mid d_{ij}] = V_A E[\pi_{ij} \mid d_{ij}] = V_A \bar{\pi}_{d_{ij}}. \quad (11)$$

1.7 Summary

Each level conditions on coarser information than the one above and is its conditional expectation; the rightmost column lists the assumption(s) newly required to descend one step.

Explicitly, each level is the conditional expectation of the finer level above it, averaging out one additional layer of information:

$$\begin{aligned} E[y_i y_j \mid \mathbf{x}_i, \mathbf{x}_j] &= E[g_i g_j \mid \mathbf{x}_i, \mathbf{x}_j] = V_A A_{ij}, \\ E[y_i y_j \mid \pi_{ij}] &= E[V_A A_{ij} \mid \pi_{ij}] = V_A \pi_{ij}, \\ E[y_i y_j \mid d_{ij}] &= E[V_A \pi_{ij} \mid d_{ij}] = V_A \bar{\pi}_{d_{ij}}, \end{aligned}$$

starting from the Level-0 value $E[y_i y_j \mid \mathbf{x}_i, \mathbf{x}_j, \boldsymbol{\beta}] = g_i g_j$. Equivalently, the ladder telescopes into a single nested expectation,

$$E[y_i y_j \mid d_{ij}] = E\left[E\left[E[g_i g_j \mid \mathbf{x}_i, \mathbf{x}_j] \mid \pi_{ij}\right] \mid d_{ij}\right] = V_A \bar{\pi}_{d_{ij}}.$$

This is valid down the generative chain $d_{ij} \rightarrow \pi_{ij} \rightarrow (\mathbf{x}_i, \mathbf{x}_j) \rightarrow y$: conditioned on the finer variable, the phenotype does not depend further on the coarser one, so the tower property applies at each step.

Level	Conditioned on	$E[y_i y_j \mid \cdot]$	New assumptions
0	causal genotypes	$g_i g_j$	[Additivity] (given [No-IGE] , [Env-indep])
1	realized sharing A_{ij}	$V_A A_{ij}$	[Zero-mean-effects] , [Exchangeable-effects] , [Uncorrelated-effects] , [Genotype-independent-effects]
2	realized relatedness π_{ij}	$V_A \pi_{ij}$	[No-mutation] , [Non-IBD-indep] , [No-inbreeding]
3	genetic degree d_{ij}	$V_A \bar{\pi}_{d_{ij}}$	[Genetic-degree]

The property that makes this a ladder is worth naming, because the next architecture breaks it. At each step, the finer variable *screens off* the coarser one: once we condition on π_{ij} , knowing d_{ij} as well would tell us nothing more about $y_i y_j$, and likewise for the other steps. Screening is what licenses the tower property, and it is the reason the four information sets can be arranged in a single descending chain rather than a more complicated structure.

2 Additive genetics under assortative mating

We will now do away with the assumption of random mating (particularly implied in [\[Non-IBD-indep\]](#)) and instead provide a model for assortative mating that will let us quantify its impact on phenotypic covariance.

The phenotype model remains the same: we assume an additive genetic architecture ([\[Additivity\]](#)) with an independent environmental component ([\[GE-indep\]](#)). As a consequence of the assortative mating model we adopt below, covariance between the environmental and genetic components between some individuals will arise, breaking the weakest forms of the assumptions [\[No-IGE\]](#) and [\[Env-indep\]](#). We also still assume no new mutations ([\[No-mutation\]](#)) and no inbreeding ([\[No-inbreeding\]](#)).

2.1 The mating model

We assume a model of assortative mating based on mates' similarity in a sorting factor, which is also our focal trait.

[\[Primary-AM\]](#) Primary Assortative Mating

Mating is non-random such that two mates' phenotypes are correlated for some trait at the time of mating:

$$\text{Corr}[y_m, y_p] = \rho_y,$$

where the subscripts m and p index the maternal and paternal member of a mated pair. Matching is on the phenotype *only*: any covariance between the mates' trait components is a downstream consequence of this phenotypic matching, and is not specified separately. Following the convention that a subscript names the component being correlated, we write ρ_g for the induced correlation between mates' genetic values: $\text{Corr}[g_m, g_p] = \rho_g^{(t)}$, where the superscript records that this induced correlation changes from one generation to the next, unlike ρ_y , which is held fixed across generations.

Commonly bundled into: primary phenotypic assortment, as opposed to social homogamy (matching on a shared environment) or genetic homogamy (matching on genotype directly); it also excludes convergence, where mates grow more similar after mating. ρ_y is often written μ or r in the literature.

2.2 Dynamics and equilibrium

To make modelling the dynamics of populations easier, we will make the following assumption:

[Discrete-generations] Discrete, non-overlapping generations

The population is partitioned into generations indexed by t , and every individual mates within their own generation.

Commonly bundled into: no overlapping generations and no age structure; in particular no mating between individuals of different generations, and no individual contributing offspring to more than one generation.

When starting from a randomly mating population, one generation of assortative mating causes the immediate offspring generation to exhibit different phenotypic and genetic properties on expectation. Luckily, if the parameters of assortative mating remain fixed, an equilibrium is reached such that stable properties of the population can be studied. We will first explore how assortative mating affects statistical properties of the population during the first few generations, then we will solve for the parameter values that lead to an equilibrium state.

2.2.1 Path Analysis

First, we'll introduce path analysis, a very helpful tool for modelling the covariance between variables in a causal system, such as genetic inheritance and assortative mating. Path analysis was introduced by Sewall Wright [1], as a way of decomposing an observed correlation between variables into the contributions of a hypothesized causal structure, and set out systematically thirteen years later as the method of path coefficients [2]. Wright applied it to mating systems in the same period [3, 4]; together with Fisher's

treatment of the same problem [5], that work is the origin of the classical results on how assortment changes the correlation between relatives — the results this section rederives. The explanations below are adapted from the Supplementary Note of Sunde et al. [6], which introduced the co-path edge type (described below). Unlike Sunde et al. [6], we do not assume that the variables are standardized, so we will keep track of variances explicitly.

A *path diagram* is a graphical representation of a generative model. Each variable is a node — drawn as a rectangle if it is observed and an ellipse if it is latent — and the nodes are joined by three kinds of edges:

- A **directed** edge $a \rightarrow b$ labelled with a coefficient c says that a is a cause of b , contributing ca to b 's structural equation. Collecting the arrows that point *into* a node recovers that node's equation, so the diagram and the model equations carry the same content.
- A **bidirected** edge $a \leftrightarrow b$ carries covariance between the *disturbances* of a and b . A variable's disturbance is what remains of it once its causes in the diagram are accounted for: if the arrows into b give $b = c_1a_1 + \dots + c_ka_k + \epsilon_b$, then ϵ_b is b 's disturbance, and for a variable with no incoming arrows the disturbance is the variable itself. The self-loop $a \leftrightarrow a$ is the case $b = a$: it is $\text{Var}[\epsilon_a]$, the part of $\text{Var}[a]$ not supplied by a 's causes, so a variable with no incoming arrows carries its entire variance on such a loop.
- A **co-path** $a - b$, drawn as a plain line with no arrowheads, is covariance attributable to *matching*: a and b came to be associated because they were paired on their values, not because they share a cause.

The covariance between *any* two variables in the diagram, latent or observed, can be calculated quickly by following Wright's tracing rules. Enumerate every *valid chain* joining the two variables; their covariance is the sum over chains of the product of the coefficients each chain passes through. A valid chain between a and b has the shape

$$a \leftarrow \dots \leftarrow u \leftrightarrow v \rightarrow \dots \rightarrow b :$$

trace *backward* along directed edges from a to some ancestor u , cross *exactly one* bidirected edge $u \leftrightarrow v$ (possibly a self-loop, $u = v$), then trace *forward* along directed edges from v down to b . A chain turns around exactly once, and it turns around on a bidirected edge, because that is the only place variance enters the system.

Co-paths extend this. A chain may pass *through* a co-path and keep going: a co-path joins two valid chains of the kind above into a longer one,

$$\underbrace{a \leftarrow \dots \leftrightarrow \dots \rightarrow y_m}_{\text{valid chain}} - \underbrace{y_p \leftarrow \dots \leftrightarrow \dots \rightarrow b}_{\text{valid chain}} .$$

Two further rules apply [6]: a chain can neither start nor end with a co-path, and a chain cannot use two co-paths belonging to the same mating process. It *may* cross co-paths of different mating processes.

A bidirected edge is a statement about disturbances, so a chain terminates on it: a disturbance has no causes in the diagram, so there is nothing to keep tracing back through, and the chain cannot reach past y_m to the things that caused y_m . A co-path is a statement about the matched variables themselves, so a chain continues backward through it,

which encodes the fact that matching two people on their phenotypes induces correlation between *all the causes* of those phenotypes — their genetic values, and every individual variant behind them. This is the mathematical basis behind the [\[Primary-AM\]](#) assumption.

The quantity multiplied along a chain when it crosses a co-path is not the correlation ρ_y itself, but

$$\mu^{(t)} = \frac{\rho_y}{\sqrt{\text{Var}[y_m^{(t)}] \text{Var}[y_p^{(t)}]}} = \frac{\rho_y}{V_Y^{(t)}}, \quad (12)$$

because a co-path with coefficient μ contributes $\text{Cov}[a, b] = \mu \text{Var}[a] \text{Var}[b]$ [6]. The two coincide only when the matched variables have unit variance. Note the use of a superscript to denote the generation a quantity belongs to, since $V_Y^{(t)}$ changes from one generation to the next under assortative mating (described below). A value of μ carried unchanged from one generation to the next is therefore wrong, which is the practical reason we label a co-path with its correlation in brackets, as $[\rho_y]$, and leave $\mu^{(t)}$ to be derived from the diagram's own variances.

Tracing chains by hand is instructive for a single pair and hopeless for a pedigree, since the number of chains grows multiplicatively with depth. The diagrams and covariances in this section were produced with **pathMgr**, a small package written alongside this writeup: it takes a model specification, returns the covariance between any two of its variables as an algebraic expression, and emits the path diagram to go with it. It computes each covariance in two independent ways — by enumerating chains exactly as above, and in closed form from the matrix identity $\Sigma = \mathbf{F}(\mathbf{I} - \mathbf{A})^{-1} \mathbf{S}(\mathbf{I} - \mathbf{A})^{-\top} \mathbf{F}^\top$ — and their agreement is the correctness check.

That identity is the diagram written as matrices, one row and column per variable. Collecting all the variables into a vector $\mathbf{v}$, the matrix $\mathbf{A}$ holds the **directed** edges — A_{ij} is the coefficient on the arrow from j to i — so the whole system of structural equations is $\mathbf{v} = \mathbf{A}\mathbf{v} + \boldsymbol{\epsilon}$, where $\boldsymbol{\epsilon}$ is the vector of disturbances. The symmetric matrix $\mathbf{S}$ holds the **bidirected** edges: S_{ij} is the value of $i \leftrightarrow j$, and its diagonal S_{ii} is the self-loop, the variance of i 's disturbance. $\mathbf{F}$ merely selects rows, keeping the variables we want reported and discarding the rest.

The inverse is what does the tracing. In an acyclic model $\mathbf{A}$ is nilpotent, so $(\mathbf{I} - \mathbf{A})^{-1} = \mathbf{I} + \mathbf{A} + \mathbf{A}^2 + \dots$, and the (i, u) entry of that sum is the total over all directed paths from u to i of the product of their coefficients — one term per path, $\mathbf{A}^n$ contributing the paths of length n . Sandwiching $\mathbf{S}$ between it and its transpose therefore gives, for latent and observed variables alike,

$$\Sigma_{ij} = \sum_u \sum_v \underbrace{[(\mathbf{I} - \mathbf{A})^{-1}]_{iu}}_{i \leftarrow \dots \leftarrow u} \underbrace{S_{uv}}_{u \leftrightarrow v} \underbrace{[(\mathbf{I} - \mathbf{A})^{-1}]_{jv}}_{v \rightarrow \dots \rightarrow j},$$

which is Wright's rule term for term: a backward leg, exactly one bidirected edge, a forward leg, summed over every chain, indexed by the $u \leftrightarrow v$ it turns around on. The two methods are the same statement — one enumerated, one summed in closed form — which is why their agreement tests the implementation rather than the mathematics. Co-paths are the one thing that sits outside this identity and contributes further terms; there, the chain rules given above are the definition.

2.2.2 Consequences of assortative mating on first generation of mates

We assume a population has been randomly mating as described by assumption [\[Non-IBD-indep\]](#) up to some generation $t = 0$, which can also be referred to as the base or randomly-mating population. The parameters defined in the previous section, such as the additive variance V_A and heritability h^2 , are measured in this base population and denoted by a (0), e.g. $V_A^{(0)}$ and $h_{(0)}^2$, since these terms do not remain fixed across generations under assortative mating. Note that V_E remains fixed.

Building the model from haploid alleles

To properly model the effects of assortative mating, it pays off to think in terms of individual haploid alleles, rather than diploid genotypes. We previously defined $z_{ik}^{(m)}$ and $z_{ik}^{(p)}$ as the standardized values of the maternally- and paternally-inherited alleles at variant k for individual i , and $x_{ik} = z_{ik}^{(m)} + z_{ik}^{(p)}$ as the standardized diploid genotype. Each haploid allele $a^{(m)}$ and $a^{(p)}$ can take on two values, 0 or 1, and therefore each has a variance of $p_k(1 - p_k)$, where p_k is the allele frequency at variant k . Typically, diploid genotypes are standardized by their expected variance when the two alleles are independent, which is $2p_k(1 - p_k)$, resulting in a standardized variance of 1, as used in the previous section. Under this standardization, each haploid allele has a standardized variance of $\frac{1}{2}$, and if we assume that the allele frequency does not change over time, then neither does this quantity:

[\[Constant-freq\]](#) Allele frequencies do not change

The allele frequency p_k of each variant is the same across every generation. As a consequence, the standardization divides by a fixed quantity, and the standardized variance of each haploid allele is the same in every generation:

$$z_{ik} = \frac{a_{ik} - p_k}{\sqrt{2p_k(1 - p_k)}} \implies \text{Var}[z_{ik}] = \frac{\text{Var}[a_{ik}]}{2p_k(1 - p_k)} = \frac{p_k(1 - p_k)}{2p_k(1 - p_k)} = \frac{1}{2},$$

where $a_{ik} \in \{0, 1\}$ is the unstandardized haploid allele count and p_k is the allele frequency at variant k .

Commonly bundled into: no selection on the trait, no drift, no migration; in a finite population it is the large- N_e limit [\[7\]](#).

In contrast, the standardized diploid genotype can have non-unit variance and change over time if the two alleles are correlated, as will be demonstrated in the case of assortative mating. Our path diagram will therefore start with haploid alleles which are summed together to form diploid genotypes, which are then summed to form the additive genetic value, which is then added to the environmental value to form the phenotype.


The diagram

Mates in generation $t = 0$ are formed according to [Primary-AM]. Below is the path diagram for a single pair that assortatively mated on the focal trait with strength ρ_y . The trait's genetic component is the additive sum of two causal variants, which is the smallest example that shows a *cross-variant* covariance; the formulas we state work for any number of variants.

One chain is traced on the diagram to show the rules in use. It runs from a single allele in one mate to a single allele at a *different* variant in the other — a covariance that is zero under random mating, and for which there is no common ancestor to appeal to. Multiplying the coefficients along it gives

$$\text{Cov}[z_{m,1}^{(m)}, z_{p,2}^{(p)}] = \frac{1}{2} \cdot \beta_1 \cdot \mu^{(0)} \cdot \beta_2 \cdot \frac{1}{2} = \frac{\beta_1 \beta_2 \rho_y}{4 V_Y^{(0)}},$$

Because there are four such allele pairs behind any one pair of genotypes, they sum back to $\text{Cov}[x_{m,1}, x_{p,2}] = \beta_1 \beta_2 \rho_y / V_Y^{(0)}$.



Doing the same for every pair of variables in the diagram gives the two tables below. Within an individual (Table 1) nothing has changed from the randomly mating case: these are base-population individuals, so their genotypes are still uncorrelated with each other and with their own environment.

	$x_{i,1}$	$x_{i,2}$	g_i	e_i	y_i
$x_{i,1}$	1	0	β_1	0	β_1
$x_{i,2}$	0	1	β_2	0	β_2
g_i	β_1	β_2	$V_A^{(0)}$	0	$V_A^{(0)}$
e_i	0	0	0	V_E	V_E
y_i	β_1	β_2	$V_A^{(0)}$	V_E	$V_Y^{(0)}$

Table 1: Covariances within one individual $i \in \{m, p\}$, identical for both members of the pair. Here $V_A^{(0)} = \beta_1^2 + \beta_2^2$ and $V_Y^{(0)} = V_A^{(0)} + V_E$. The last column — each variable’s covariance with its own phenotype — is the only part of this table needed for computing the covariance between variables across mates.

The co-path induces covariance between the factors pertaining to mates such that:

$$\text{Cov}[a_m, b_p] = \text{Cov}[a_m, y_m] \cdot \mu^{(0)} \cdot \text{Cov}[y_p, b_p] \quad (13)$$

for any variables a and b belonging to the maternal and paternal member respectively. The entire cross-mate block is therefore the outer product of a single vector — the last column of Table 1 — scaled by $\mu^{(0)} = \rho_y/V_Y^{(0)}$.

$\times \mu^{(0)}$	$x_{p,1}$	$x_{p,2}$	g_p	e_p	y_p
$x_{m,1}$	β_1^2	$\beta_1\beta_2$	$\beta_1V_A^{(0)}$	β_1V_E	$\beta_1V_Y^{(0)}$
$x_{m,2}$	$\beta_1\beta_2$	β_2^2	$\beta_2V_A^{(0)}$	β_2V_E	$\beta_2V_Y^{(0)}$
g_m	$\beta_1V_A^{(0)}$	$\beta_2V_A^{(0)}$	$(V_A^{(0)})^2$	$V_A^{(0)}V_E$	$V_A^{(0)}V_Y^{(0)}$
e_m	β_1V_E	β_2V_E	$V_A^{(0)}V_E$	V_E^2	$V_EV_Y^{(0)}$
y_m	$\beta_1V_Y^{(0)}$	$\beta_2V_Y^{(0)}$	$V_A^{(0)}V_Y^{(0)}$	$V_EV_Y^{(0)}$	$(V_Y^{(0)})^2$

Table 2: Covariances between the two mates. **Every entry carries a factor $\mu^{(0)} = \rho_y/V_Y^{(0)}$** , shown once in the corner rather than repeated in twenty-five cells. The block is symmetric, and is the outer product of $(\beta_1, \beta_2, V_A^{(0)}, V_E, V_Y^{(0)})$ with itself — a direct consequence of (13). Setting $\rho_y = 0$ returns every entry to zero, recovering the randomly mating case of Section 1.

Three key results from Table 2 are highlighted here. We also generalize from two variants to M variants, where $g_i = \sum_{k=1}^M \beta_k x_{ik}$.

- $\text{Cov}[x_{mk}, x_{pl}] = \beta_k \beta_l \rho_y / V_Y^{(0)} \neq 0$, for *every* pair of variants k and l . Alleles at different loci in different individuals are correlated, signed by the product of their effects and breaking the [\[Non-IBD-indep\]](#) assumption. Note that in the base population, the within-individual covariance $\text{Cov}[x_{mk}, x_{ml}]$ is still exactly zero for $k \neq l$; the disequilibrium exists only *between* the mates.

- $\text{Cov}[e_m, g_p] = V_A^{(0)} V_E \rho_y / V_Y^{(0)} \neq 0$. One mate's environment is correlated with the other's genetic value, which is the break of [\[Env-indep\]](#) and [\[No-IGE\]](#) flagged at the start of this section. Despite one mate's genotype not causally affecting the other's environment, the two still become correlated due to the phenotypic matching.
- $\rho_g^{(0)} = \text{Corr}[g_m, g_p] = \frac{(V_A^{(0)})^2 \rho_y / V_Y^{(0)}}{V_A^{(0)}} = \rho_y \frac{V_A^{(0)}}{V_Y^{(0)}} = \rho_y h_{(0)}^2$. Mates' genetic values are correlated according to the strength of assortative mating and the heritability of the sorting trait. Equivalently, this is the sum of the first result over all M^2 ordered pairs of variants, weighted by their effects: $\sum_k \sum_l \beta_k \beta_l \cdot \beta_k \beta_l \rho_y / V_Y^{(0)} = (\sum_k \beta_k^2)^2 \rho_y / V_Y^{(0)}$.

2.2.3 Consequences of assortative mating on first generation of offspring

Next, we will extend our path diagram to contain two offspring of an assortatively mated pair. For each variant, all offspring inherit one allele from each parent, but which allele they inherit from that parent (the maternal vs paternal allele) is uniformly random. For the purposes of tracking covariances, we model this by drawing an edge from each parental allele to their respective inherited allele in the offspring and adding an exogenous source of variance. The directed edges have a coefficient of $\frac{1}{2}$ to reflect that the expected inherited allele in the offspring is the average of the two alleles of the parent being inherited from.

Write o for an offspring of the mated pair, and let $B_{ok}^{(m)} \in \{0, 1\}$ indicate which of the mother's two alleles was transmitted at variant k . We arbitrarily label the maternal allele $z_{mk}^{(m)}$ and the paternal allele $z_{mk}^{(p)}$ such that if $B_{ok}^{(m)} = 1$, then the offspring inherits the maternal allele, and if $B_{ok}^{(m)} = 0$ then the offspring inherits the paternal allele.

$$z_{ok}^{(m)} = B_{ok}^{(m)} z_{mk}^{(m)} + (1 - B_{ok}^{(m)}) z_{mk}^{(p)}, \quad B_{ok}^{(m)} \sim \text{Bernoulli}(\tfrac{1}{2}), \quad B_{ok}^{(m)} \perp z_{mk}. \quad (14)$$

A path diagram represents the *linear projection* of the random variable: the best prediction of the offspring's allele from the mother's two, which is their average,

$$\mathbb{E}[z_{ok}^{(m)} \mid z_{mk}^{(m)}, z_{mk}^{(p)}] = \tfrac{1}{2} z_{mk}^{(m)} + \tfrac{1}{2} z_{mk}^{(p)} = \tfrac{1}{2} x_{mk}. \quad (15)$$

The exogenous variance is what the projection leaves behind. Subtracting (15) from (14) gives the *segregation deviation*

$$z_{ok}^{\perp(m)} := z_{ok}^{(m)} - \tfrac{1}{2} x_{mk} = (B_{ok}^{(m)} - \tfrac{1}{2})(z_{mk}^{(m)} - z_{mk}^{(p)}), \quad (16)$$

which is mean-zero and uncorrelated with both of the mother's alleles — a disturbance in the sense of Section 2.2.1, and drawn as one. The offspring's paternally inherited allele $z_{ok}^{(p)}$ is built the same way from the father's two alleles, with its own independent $B_{ok}^{(p)}$ and residual $z_{ok}^{\perp(p)}$.

Covariance between alleles

We now define the covariances that will be repeatedly used in deriving the consequences of assortative mating. For an individual i of generation t and variants k and l where $u, v \in \{m, p\}$,

$$\boxed{a_{kl}^{(t)} := \text{Cov}[z_{ik}^{(u)}, z_{il}^{(v)}] \quad (u \neq v), \quad s_{kl}^{(t)} := \text{Cov}[z_{ik}^{(u)}, z_{il}^{(u)}]} \quad (17)$$

In words, a denotes the covariance between alleles *across* gametes, while s denotes the covariance between alleles on the *same* gamete. When $k = l$, $s_{kk}^{(t)}$ is the variance of an individual allele, which is fixed for all generations due to how we standardized variables according to [\[Constant-freq\]](#):

$$s_{kk}^{(t)} \equiv \frac{1}{2} \quad \text{for all } t, \quad (18)$$

We can also write the covariance between two genotypes in the same individual as the sum of four allelic covariances:

$$\text{Cov}[x_{ik}, x_{il}] = \sum_{u,v \in \{m,p\}} \text{Cov}[z_{ik}^{(u)}, z_{il}^{(v)}] = 2(s_{kl}^{(t)} + a_{kl}^{(t)}), \quad (19)$$

In the case of $k = l$, it becomes the variance of genotypes, since $s_{kk}^{(t)} \equiv \frac{1}{2}$:

$$\begin{aligned} \text{Var}[x_{ik}] &= 2(s_{kk}^{(t)} + a_{kk}^{(t)}) \\ \boxed{\text{Var}[x_{ik}] &= 1 + 2a_{kk}^{(t)}} \end{aligned} \quad (20)$$

The third quantity we need is the covariance between a single allele and the phenotype of the individual carrying it. Expanding the phenotype into its causes and then each genotype into its two alleles,

$$\boxed{c_k^{(t)} := \text{Cov}[z_{ik}^{(u)}, y_i] = \sum_{l=1}^M \beta_l \text{Cov}[z_{ik}^{(u)}, x_{il}] = \sum_{l=1}^M \beta_l (s_{kl}^{(t)} + a_{kl}^{(t)})} \quad (21)$$

The first step drops e_i , which is uncorrelated with every allele of the same individual by [\[GE-indep\]](#). The second splits each genotype x_{il} into the allele sitting on the same gamete as $z_{ik}^{(u)}$, contributing $s_{kl}^{(t)}$, and the one on the opposite gamete, contributing $a_{kl}^{(t)}$. A single allele therefore reaches the phenotype through *every* variant the individual carries, not only through its own.

Summing over alleles in the same way gives the additive variance of the generation:

$$\begin{aligned} V_A^{(t)} &= \text{Var}\left[\sum_k \beta_k x_{ik}\right] = \sum_k \sum_l \beta_k \beta_l \text{Cov}[x_{ik}, x_{il}] \\ &= 2 \sum_k \sum_l \beta_k \beta_l (s_{kl}^{(t)} + a_{kl}^{(t)}) \end{aligned} \quad (22)$$

$$\boxed{V_A^{(t)} = 2 \sum_k \beta_k c_k^{(t)}}$$

with $V_Y^{(t)} = V_A^{(t)} + V_E$, since assortment on the phenotype does not change the environmental variance. Similarly, we define generation-specific heritability as follows:

$$\boxed{h_{(t)}^2 := \frac{V_A^{(t)}}{V_Y^{(t)}} = \frac{V_A^{(t)}}{V_A^{(t)} + V_E} = \frac{2 \sum_k \beta_k c_k^{(t)}}{2 \sum_k \beta_k c_k^{(t)} + V_E}} \quad (23)$$

We'll also confirm here that $\rho_g^{(t)} = \frac{\rho_y V_A^{(t)}}{V_Y^{(t)}}$. A single co-path joins the two mates, so (13) applies with $\mu^{(t)}$ in place of $\mu^{(0)}$, and a genetic value's covariance with its own carrier's phenotype is $\text{Cov}[g_i, g_i + e_i] = V_A^{(t)}$ by **[GE-indep]**. Both mates have genetic variance $V_A^{(t)}$, so the denominator of the correlation is $V_A^{(t)}$ as well:

$$\rho_g^{(t)} = \frac{\text{Cov}[g_m, g_p]}{\sqrt{\text{Var}[g_m] \text{Var}[g_p]}} = \frac{\overbrace{\text{Cov}[g_m, y_m]}^{V_A^{(t)}} \cdot \mu^{(t)} \cdot \overbrace{\text{Cov}[y_p, g_p]}^{V_A^{(t)}}}{V_A^{(t)}} = \mu^{(t)} V_A^{(t)} \quad (24)$$

$$\boxed{\rho_g^{(t)} = \frac{\rho_y V_A^{(t)}}{V_Y^{(t)}} = \rho_y h_{(t)}^2}$$

The genetic correlation between mates is the phenotypic one scaled by that generation's heritability, exactly as it was in the base population.

We can also now write the variance of the segregation deviation. The indicator $B_{ok}^{(m)}$ takes values in $\{0, 1\}$, so $(B_{ok}^{(m)} - \frac{1}{2})^2 = \frac{1}{4}$ with probability one, and it is independent of the alleles it selects between. Squaring (16) and taking expectations therefore factors cleanly:

$$\text{Var}[z_{ok}^{\perp(m)}] = \mathbb{E}\left[\left(B_{ok}^{(m)} - \frac{1}{2}\right)^2\right] \text{Var}[z_{mk}^{(m)} - z_{mk}^{(p)}] = \frac{1}{4} \left(\underbrace{\frac{1}{2}}_{s_{kk}^{(t)}} + \underbrace{\frac{1}{2}}_{s_{kk}^{(t)}} - 2 \underbrace{a_{kk}^{(t)}}_{\text{Cov}[z_{mk}^{(m)}, z_{mk}^{(p)}]} \right), \quad (25)$$

where the parent is of generation t . Collecting terms,

$$\boxed{\text{Var}[z_{ok}^{\perp(m)}] = \frac{1}{4} - \frac{1}{2} a_{kk}^{(t)}} \quad (26)$$

So a parent whose two alleles are more alike contributes *less* segregation variance: it matters less which of the two was transmitted.

First generation of offspring

In a generation-0 individual, their own alleles are uncorrelated with each other. At the same variant, the two alleles are non-IBD by **[No-inbreeding]** and hence independent by **[Non-IBD-indep]**; at different variants, the base population is in linkage equilibrium, which is the within-individual zero already noted below Table 2:

$$\boxed{a_{kl}^{(0)} = 0 \text{ for all } k, l, \quad s_{kl}^{(0)} = 0 \text{ } (k \neq l), \quad c_k^{(0)} = \frac{1}{2} \beta_k} \quad (27)$$

The third follows from the first two: with every off-diagonal term of (21) equal to zero, only $l = k$ survives, leaving $\beta_k s_{kk}^{(0)} = \frac{1}{2} \beta_k$. Setting $a_{kk}^{(0)} = 0$ in (20) recovers $\text{Var}[x_{ik}] = 1$, and in (26) recovers the familiar segregation variance of $\frac{1}{4}$.

In a generation-1 individual, this is no longer the case, which can be seen by the below path diagram of two generation-1 offspring of a mated pair that assortatively mated on the focal trait with strength ρ_y , for a trait with two causal variants.




Figure 2: One mated pair from the base population and two of their offspring, at the allele level, for a trait with two causal variants. Each inherited allele is fed by *both* alleles of the parent it came from, at $\frac{1}{2}$ each, per (15). The self-loop of $\frac{1}{4}$ on each offspring allele is its segregation variance, (26) evaluated at $a_{kk}^{(0)} = 0$.

We can use our path tracing rules to compute $a_{kl}^{(1)}$. Exactly four chains connect an allele of the offspring to an allele on its *other* gamete, at variants k and l . Each traces back from $z_{ok}^{(m)}$ to *one* of the mother's two alleles at variant k , turns around on that allele's variance, runs up through her genotype and phenotype, crosses the co-path, and comes back down the father's side to *one* of his two alleles at variant l and into $z_{ol}^{(p)}$. Two choices on each side give $2 \times 2 = 4$ chains identical in value, one for each $u, v \in \{m, p\}$:

$$\begin{aligned} & z_{ok}^{(m)} \leftarrow z_{mk}^{(u)} \leftrightarrow z_{mk}^{(u)} \rightarrow x_{mk} \rightarrow g_m \rightarrow y_m \\ \text{--- } & y_p \leftarrow g_p \leftarrow x_{pl} \leftarrow z_{pl}^{(v)} \leftrightarrow z_{pl}^{(v)} \rightarrow z_{ol}^{(p)}, \end{aligned}$$

Multiplying each coefficient along one chain returns:

$$\underbrace{\frac{1}{2}}_{\text{transmission}} \cdot \underbrace{\frac{1}{2}}_{\text{Var}[z]} \cdot \beta_k \cdot \mu^{(0)} \cdot \beta_l \cdot \underbrace{\frac{1}{2}}_{\text{Var}[z]} \cdot \underbrace{\frac{1}{2}}_{\text{transmission}} = \frac{\beta_k \beta_l \mu^{(0)}}{16}.$$

Summing across the four chains and expanding $\mu^{(0)}$,

$$\boxed{a_{kl}^{(1)} = \frac{\beta_k \beta_l \rho_y}{4 V_Y^{(0)}}} \quad (28)$$

The same-gamete companion is easier: two alleles the offspring received from the *same* parent at different variants have no chain at all connecting them, because that parent's own alleles are uncorrelated across variants due to them being of the base population, so

$$\boxed{s_{kl}^{(1)} = 0 \quad (k \neq l)} \quad (29)$$

One generation of assortment therefore produces association *across* an individual's two gametes and none within either of them.

Rearranging (24) gives $\rho_y/V_Y^{(0)} = \rho_g^{(0)}/V_A^{(0)}$, which lets (28) be restated in terms of how strongly the parents' genetic values were correlated rather than their phenotypes:

$$\boxed{a_{kl}^{(1)} = \frac{\beta_k \beta_l}{4 V_A^{(0)}} \rho_g^{(0)}} \quad (30)$$

The association between two of the offspring's alleles is the product of their effect sizes, taken as a share of the additive variance, scaled by the genetic correlation between its parents.

Feeding (28) and (29) into (21) gives the first-generation allele-phenotype covariance. Only $l = k$ survives from $s^{(1)}$, while every l contributes through $a^{(1)}$:

$$c_k^{(1)} = \underbrace{\frac{1}{2}\beta_k}_{l=k \text{ term of } s^{(1)}} + \sum_l \beta_l \frac{\beta_k \beta_l \rho_y}{4 V_Y^{(0)}} = \frac{1}{2}\beta_k \left(1 + \frac{\rho_y V_A^{(0)}}{2 V_Y^{(0)}} \right) \quad (31)$$

$$\boxed{c_k^{(1)} = \frac{1}{2}\beta_k \left(1 + \frac{1}{2}\rho_g^{(0)} \right)}$$

using $\sum_l \beta_l^2 = V_A^{(0)}$. Every variant's initial covariance with the phenotype is inflated by the same factor.

We use the path tracing rules to compute the covariance between an individual and a full sibling and a parent, shown in Table 3. Every entry there turns out to be its random-mating value plus a term proportional to $\rho_g^{(0)}$.

	self	full siblings	parent-offspring
genotypes, $k = l$	$1 + \frac{\beta_k^2}{2V_A^{(0)}}\rho_g^{(0)}$	$\frac{1}{2} + \frac{\beta_k^2}{2V_A^{(0)}}\rho_g^{(0)}$	$\frac{1}{2} + \frac{\beta_k^2}{2V_A^{(0)}}\rho_g^{(0)}$
genotypes, $k \neq l$	$\frac{\beta_k\beta_l}{2V_A^{(0)}}\rho_g^{(0)}$	$\frac{\beta_k\beta_l}{2V_A^{(0)}}\rho_g^{(0)}$	$\frac{\beta_k\beta_l}{2V_A^{(0)}}\rho_g^{(0)}$
genetic values	$V_A^{(0)}(1 + \frac{1}{2}\rho_g^{(0)})$	$\frac{1}{2}V_A^{(0)}(1 + \rho_g^{(0)})$	$\frac{1}{2}V_A^{(0)}(1 + \rho_g^{(0)})$
phenotypes	$V_A^{(0)}(1 + \frac{1}{2}\rho_g^{(0)}) + V_E$	$\frac{1}{2}V_A^{(0)}(1 + \rho_g^{(0)})$	$\frac{1}{2}V_A^{(0)}(1 + \rho_y)$

Table 3: Variances and covariances for an individual of the **first generation produced under assortative mating**, against itself and against each kind of first-degree relative: the “self” column is the pair (o_1, o_1) , “full siblings” is (o_1, o_2) , and “parent-offspring” is (m, o_1) . The first row is a genotype against a genotype at the *same* variant ($k = l$), the second at two *different* variants. Every entry is its random-mating value plus **what one generation of assortment added**. Writing the added terms through $\rho_g^{(0)}$ rather than $\rho_y/V_Y^{(0)}$ leaves the two relationships differing in a single symbol: siblings carry $\rho_g^{(0)}$ at the phenotypic level where parent and offspring carry ρ_y .

Note that covariances for parents vs full-siblings only differ at the phenotypic level due to one extra non-zero term that emerges for direct descendants:

$$\begin{aligned}
\text{Cov}[y_m, y_{o_1}] &= \underbrace{\text{Cov}[g_m, g_{o_1}]}_{\frac{1}{2}V_A^{(0)} + \frac{\rho_y(V_A^{(0)})^2}{2V_Y^{(0)}}} + \underbrace{\text{Cov}[e_m, g_{o_1}]}_{\frac{\rho_y V_A^{(0)} V_E}{2V_Y^{(0)}}} \\
&= \frac{1}{2}V_A^{(0)} + \frac{\rho_y V_A^{(0)}(V_A^{(0)} + V_E)}{2V_Y^{(0)}} = \frac{1}{2}V_A^{(0)}(1 + \rho_y),
\end{aligned}$$

where the last step uses $V_Y^{(0)} = V_A^{(0)} + V_E$. A parent’s environment is correlated with their child’s genetic value — the child carries half the alleles of the assortatively chosen partner, whose genotype is correlated with this parent’s environment.

2.2.4 Compounding across generations

If we now produce a second generation of offspring, the effects of another generation of assortative mating compound on the existing effects of past assortative mating. Namely, unlike before, the two alleles that a parent carries are now correlated according to their effect size.

It would be unwieldy to make a diagram for three generations. Instead we summarize the base population by its impact on the covariance of generation-1 alleles.




Figure 3: Two generation-1 individuals mating to produce one generation-2 offspring. Unlike Figure 2, each parent's alleles are correlated. No bidirectional arrow joins two alleles on the *same* gamete because $s_{kl}^{(1)} = 0$ by (29). The co-path still carries the correlation ρ_y , but contributes $\mu^{(1)} = \rho_y/V_Y^{(1)}$, since V_Y has grown.

Covariance across gametes. We still think in terms of the four chains described in (28), but now any chain connecting a parent's allele up to its own phenotype is no longer just $\frac{1}{2}\beta_k$: the parent's alleles are correlated now, so a chain leaving $z_{ik}^{(u)}$ can reach the phenotype through *any* other allele the parent carries. Every such route is already collected in $c_k^{(t)}$, which (21) defined as exactly that covariance. The four chains then contribute $c_k^{(t)} \cdot \mu^{(t)} \cdot c_l^{(t)}$ in total, so

$$a_{kl}^{(t+1)} = \mu^{(t)} c_k^{(t)} c_l^{(t)} \quad (32)$$

Setting $t = 0$ recovers (28), since $c_k^{(0)} = \frac{1}{2}\beta_k$ and $\mu^{(0)} = \rho_y/V_Y^{(0)}$.

Covariance within a gamete. To make derivations easier, we also assume that which chromosome in the parent an inherited allele is from is independent across variants.

[Unlinked] Variants are independently segregated

The transmission indicators of different variants are independent: for $k \neq l$, $B_{ok}^{(m)} \perp B_{ol}^{(m)}$, and likewise on the paternal side. Equivalently, every pair of causal variants recombines with probability $\frac{1}{2}$ in every meiosis.

Commonly bundled into: free recombination, no physical linkage

Under [Unlinked],

$$s_{kl}^{(t+1)} = \frac{1}{4}\text{Cov}[x_{ik}, x_{il}] = \frac{1}{2}\left(s_{kl}^{(t)} + a_{kl}^{(t)}\right) \quad (33)$$

using (19) for the parent's genotype covariance. In other words, when picking two alleles from the same gamete in an offspring, they are equally likely to have been from the same vs different gamete in the parent. Therefore, the covariance between the two offspring's alleles is the average of the across- vs same-gamete covariance in the parent generation.

While $s_{kl}^{(1)} = 0$, the second generation is where the same-gamete covariance becomes non-zero: $s_{kl}^{(2)} = \frac{1}{2}a_{kl}^{(1)} = \beta_k\beta_l\rho_y/(8V_Y^{(0)})$.

Covariance between an allele and the phenotype. We rewrite the formula for c from (21), one generation later, as two separate sums:

$$c_k^{(t+1)} = \sum_{l=1}^M \beta_l \left(s_{kl}^{(t+1)} + a_{kl}^{(t+1)} \right) = \underbrace{\sum_l \beta_l a_{kl}^{(t+1)}}_{\text{across gametes}} + \underbrace{\sum_l \beta_l s_{kl}^{(t+1)}}_{\text{same gamete}}. \quad (34)$$

The across-gamete part can be written in terms of the previous generation's c by (32). After pulling out common terms, we are left with a summation that can be simplified

according to (22). We then plug in $\mu^{(t)} = \rho_y/V_Y^{(t)}$, which leaves $\rho_y V_A^{(t)}/V_Y^{(t)}$ — and that is $\rho_g^{(t)}$ by (24):

$$\begin{aligned} \sum_l \beta_l a_{kl}^{(t+1)} &= \sum_l \beta_l \mu^{(t)} c_k^{(t)} c_l^{(t)} = \mu^{(t)} c_k^{(t)} \underbrace{\sum_l \beta_l c_l^{(t)}}_{V_A^{(t)}/2} = \frac{\rho_y V_A^{(t)}}{2 V_Y^{(t)}} c_k^{(t)} \\ &= \frac{1}{2} \rho_g^{(t)} c_k^{(t)}, \end{aligned} \quad (35)$$

For the same-gamete part we fill in the recursive expression for $s_{kl}^{(t+1)}$ according to (33), which lets us write it in terms of $c_k^{(t)}$ according to (21). However, we must be careful to replace the $k = l$ case with its fixed value of $s_{kk}^{(t)} \equiv \frac{1}{2}$:

$$\begin{aligned} \sum_l \beta_l s_{kl}^{(t+1)} &= \underbrace{\frac{1}{2} \sum_l \beta_l (s_{kl}^{(t)} + a_{kl}^{(t)})}_{c_k^{(t)}} + \beta_k \left[\underbrace{\frac{1}{2}}_{s_{kk}^{(t+1)}} - \underbrace{\frac{1}{2} \left(\frac{1}{2} + a_{kk}^{(t)} \right)}_{\text{what (33) would give for } k=l} \right] \\ &= \frac{1}{2} c_k^{(t)} + \beta_k \left(\frac{1}{4} - \frac{1}{2} a_{kk}^{(t)} \right). \end{aligned} \quad (36)$$

The bracket is the correction for the one term the recursion does not govern.

We now add the two halves. Note that $\rho_g^{(t)}$ appears in only one of them: it enters through (35), because that is the half whose chains cross the co-path between the parents, and it is absent from (36), whose chains stay inside one parent. What the two halves share is a factor of $\frac{1}{2} c_k^{(t)}$, and collecting that factor is what puts $\rho_g^{(t)}$ next to a 1:

$$\begin{aligned} c_k^{(t+1)} &= \underbrace{\frac{1}{2} \rho_g^{(t)} c_k^{(t)}}_{\text{across gametes, (35)}} + \underbrace{\frac{1}{2} c_k^{(t)} + \beta_k \left(\frac{1}{4} - \frac{1}{2} a_{kk}^{(t)} \right)}_{\text{same gamete, (36)}} \\ &\boxed{c_k^{(t+1)} = \underbrace{\frac{1}{2} (1 + \rho_g^{(t)}) c_k^{(t)}}_{\text{inherited}} + \underbrace{\beta_k \left(\frac{1}{4} - \frac{1}{2} a_{kk}^{(t)} \right)}_{\text{Var}[z_{ok}^{\perp(m)}]}} \end{aligned} \quad (37)$$

So the 1 is the allele the offspring actually received, and the $\rho_g^{(t)}$ beside it is the extra covariance that allele picked up because the parent it came from was chosen to resemble the other parent. Under random mating $\rho_g^{(t)} = 0$ and only the 1 remains. We can notice that the segregation variance (26) naturally emerges from our formula and brings with it a satisfying interpretation. We have previously thought of offspring alleles as having a directly-inherited component from the parent (equivalent to half the parent's genotype) plus segregation deviation that is uncorrelated with the parent's genotype. The former contributes half the covariance present in the parent generation, but amplified slightly due to assortative mating predicting the total covariance of correlated variants coming from the other parent. The latter contributes additional variance directly proportional to that variant's effect size.

We can expand out the terms of equation (37) such that they only rely on the starting parameters — the effect sizes β_k , the phenotypic correlation between mates ρ_y , and the

environmental variance V_E — and previous states of c . Substituting $\rho_g^{(t)} = \rho_y V_A^{(t)} / V_Y^{(t)}$, then $V_A^{(t)} = 2 \sum_j \beta_j c_j^{(t)}$ and $V_Y^{(t)} = V_A^{(t)} + V_E$, and finally $a_{kk}^{(t)} = \mu^{(t-1)} (c_k^{(t-1)})^2$ leaves

$$c_k^{(t+1)} = \frac{1}{4} \beta_k + \frac{1}{2} \left(1 + \frac{2 \rho_y \sum_j \beta_j c_j^{(t)}}{2 \sum_j \beta_j c_j^{(t)} + V_E} \right) c_k^{(t)} - \frac{\rho_y \beta_k}{2 \left(2 \sum_j \beta_j c_j^{(t-1)} + V_E \right)} (c_k^{(t-1)})^2, \quad (38)$$

with $c_k^{(t)} = \frac{1}{2} \beta_k$ for every $t \leq 0$, the base population having been reached without assortment.

This reveals that the covariances in this population follow an *autonomous second-order difference equation*: autonomous because the update rule is the same at every generation, and second-order because it reaches back two of them. The M variants form one coupled system rather than M separate equations, since each variant's update reads all the others through $V_A^{(t)}$. It is also *nonlinear* — $c_k^{(t-1)}$ appears squared, and $c^{(t)}$ appears inside a denominator — so the usual closed-form solutions for linear recursions do not apply, and we look instead for a *fixed point*: a c that the update leaves unchanged. The leading factor $\frac{1}{2}(1 + \rho_g^{(t)})$ is less than one whenever $\rho_g^{(t)} < 1$, which makes the update a contraction near such a point, so the approach to it is geometric rather than oscillatory or divergent: the remaining distance falls by a roughly constant fraction each generation.

		$t \leq 0$ base pop.	$t = 1$	general (t)
allele-phenotype	$c_k^{(t)}$	$\frac{1}{2}\beta_k$	$\frac{1}{2}\beta_k(1 + \frac{1}{2}\rho_g^{(0)})$	$\frac{1}{2}(1 + \rho_g^{(t-1)})c_k^{(t-1)} + \beta_k(\frac{1}{4} - \frac{1}{2}a_{kk}^{(t-1)})$
across gametes	$a_{kl}^{(t)}$	0	$\frac{\beta_k\beta_l}{4V_A^{(0)}}\rho_g^{(0)}$	$\mu^{(t-1)}c_k^{(t-1)}c_l^{(t-1)}$
same gamete	$s_{kl}^{(t)}, k \neq l$	0	0	$\frac{1}{2}(s_{kl}^{(t-1)} + a_{kl}^{(t-1)})$
genotype variance	$\text{Var}[x_{ik}]$	1	$1 + \frac{\beta_k^2}{2V_A^{(0)}}\rho_g^{(0)}$	$1 + 2a_{kk}^{(t)}$
segregation variance	$\text{Var}[z_{ok}^{\perp(m)}]$	$\frac{1}{4}$	$\frac{1}{4} - \frac{\beta_k^2}{8V_A^{(0)}}\rho_g^{(0)}$	$\frac{1}{4} - \frac{1}{2}a_{kk}^{(t)}$
additive variance	$V_A^{(t)}$	$V_A^{(0)}$	$V_A^{(0)}(1 + \frac{1}{2}\rho_g^{(0)})$	$2\sum_k \beta_k c_k^{(t)}$
phenotypic variance	$V_Y^{(t)}$	$V_A^{(0)} + V_E$	$V_A^{(1)} + V_E$	$V_A^{(t)} + V_E$
heritability	$h_{(t)}^2$	$V_A^{(0)}/V_Y^{(0)}$	$V_A^{(1)}/V_Y^{(1)}$	$V_A^{(t)}/V_Y^{(t)}$
genetic correlation	$\rho_g^{(t)}$	$\rho_y h_{(0)}^2$	$\rho_y h_{(1)}^2$	$\rho_y h_{(t)}^2$
co-path coefficient	$\mu^{(t)}$	$\rho_y/V_Y^{(0)}$	$\rho_y/V_Y^{(1)}$	$\rho_y/V_Y^{(t)}$

Table 4: Every quantity derived in this section, from the base population forward, for a trait with M variants. Every column gives that generation's own value, the left-hand side always being the quantity named in the second column. The three recursions therefore reach back to $t - 1$ in the last column, while the rest are evaluated at t directly. The first block are the three allele covariances, of which only c is a recursion the others do not feed: a and s are read off it, and $s_{kk}^{(t)} \equiv \frac{1}{2}$. The second block is read off the same generation's c , and involves no recursion at all. The base-population column is written $t \leq 0$ because nothing has happened yet: its ancestors mated at random, so c , a and s take these values at every $t \leq 0$, which is what lets the second-order (38) start. Assortment begins with the generation-0 pairing, so $\rho_g^{(t)}$ and $\mu^{(t)}$ in that column are the ones governing the step to $t = 1$.

2.2.5 Equilibrium

Our next goal is to solve for the values that lead to an equilibrium state in the recursive covariance formulas we expressed above. An equilibrium is a state the update in (37) leaves unchanged, so we look for a c satisfying $c_k^{(t+1)} = c_k^{(t)} = c_k^{(t-1)}$, which we write $c_k^{(\text{eq})}$.

[AM-equilibrium] Intergenerational equilibrium

Assortment has been operating at the same strength for long enough that the covariances have stopped changing from one generation to the next.

Commonly bundled into: equilibrium assortative mating; reached after roughly six to ten generations [8].

Under equilibrium, we fill in the same value, $c_k^{(eq)}$ for all states of c_k in (37) after replacing $a_{kk}^{(eq)} = \mu^{(eq)} (c_k^{(eq)})^2$.

$$\begin{aligned}
c_k^{(eq)} &= \frac{1}{2}(1 + \rho_g^{(eq)})c_k^{(eq)} + \beta_k \left(\frac{1}{4} - \frac{1}{2} \mu^{(eq)} (c_k^{(eq)})^2 \right) \\
\frac{1}{2}(1 - \rho_g^{(eq)})c_k^{(eq)} &= \frac{1}{4}\beta_k - \frac{1}{2} \mu^{(eq)} \beta_k (c_k^{(eq)})^2 \\
2(1 - \rho_g^{(eq)})c_k^{(eq)} &= \beta_k - 2 \mu^{(eq)} \beta_k (c_k^{(eq)})^2 \\
2\mu^{(eq)} \beta_k (c_k^{(eq)})^2 + 2(1 - \rho_g^{(eq)})c_k^{(eq)} - \beta_k &= 0
\end{aligned} \tag{39}$$

$$c_k^{(eq)} = \frac{-(1 - \rho_g^{(eq)}) \pm \sqrt{(1 - \rho_g^{(eq)})^2 + 2 \mu^{(eq)} \beta_k^2}}{2 \mu^{(eq)} \beta_k} \tag{40}$$

$$c_k^{(eq)} = \frac{\beta_k}{(1 - \rho_g^{(eq)}) + \sqrt{(1 - \rho_g^{(eq)})^2 + 2 \mu^{(eq)} \beta_k^2}}$$

Only the positive root is admissible: the negative root gives a $c_k^{(eq)}$ of the opposite sign to β_k , and does not reduce to the base-population value when assortment is switched off.

However, we still have the problem that $\mu^{(eq)}$ and $\rho_g^{(eq)}$ depend on $V_A^{(eq)}$, which itself depends on the sum of $c_k^{(eq)}$ for all k , which are quantities we are solving for. Summing (40) over variants does not resolve this, because each variant carries its own square root containing its own $2\mu^{(eq)}\beta_k^2$, so the M terms cannot be cleanly gathered into one. To make obtaining a closed form solution manageable, we need to apply an approximation that is justified by the following assumptions:

[Large-Me] The additive variance is spread across many variants

Taking inspiration from O'Connor et al. [9], define the *effective number of variants* as

$$M_e := \frac{(V_A^{(0)})^2}{\sum_k \beta_k^4} = \frac{(\sum_k \beta_k^2)^2}{\sum_k \beta_k^4} \tag{41}$$

This is closely related to the same quantity defined in O'Connor et al. [9], albeit ours is smaller by a factor of three because we are aiming for the property that $M_e = M$ when all variants have the same effect size, rather than when all variants have the same *Gaussian prior*.

Specifically, we assume that the effective number of variants is large:

$$M_e \gg 1.$$

Equivalently, no single variant explains an appreciable share of the additive variance. Since $\sum_j \beta_j^4 \geq \beta_k^4$ for any single k , rearranging (41) gives

$$\frac{\beta_k^2}{V_A^{(0)}} \leq \frac{1}{\sqrt{M_e}} \quad \text{for every } k, \tag{42}$$

so $M_e \gg 1$ forces every variance share below $M_e^{-1/2} \ll 1$.

Commonly bundled into: the polygenic, or infinitesimal, model with large M ; M_e is estimable from summary statistics [9].

[Moderate-AM] Assortment is not maximal

The genetic correlation between mates stays bounded away from its ceiling, $\rho_g^{(\text{eq})} < 1$, so that $1 - \rho_g^{(\text{eq})}$ is not close to zero.

Commonly bundled into: any trait that is not both highly heritable and highly assorted.

[Uniform-inflation] Assortment inflates every variant's covariance with the phenotype alike

$$c_k^{(\text{eq})} = \frac{\beta_k}{(1 - \rho_g^{(\text{eq})}) + \sqrt{(1 - \rho_g^{(\text{eq})})^2 + \underbrace{2\mu^{(\text{eq})}\beta_k^2}_{\text{negligible}}} \quad (43)$$

$$\boxed{c_k^{(\text{eq})} \approx \frac{\beta_k}{2(1 - \rho_g^{(\text{eq})})}}$$

Every variant is therefore inflated above its base-population value $\frac{1}{2}\beta_k$ by the same factor $1/(1 - \rho_g^{(\text{eq})})$, whatever its effect size.

Justification for approximation: The neglected term, relative to the one it sits beside, is

$$\frac{2\mu^{(\text{eq})}\beta_k^2}{(1 - \rho_g^{(\text{eq})})^2} = \underbrace{\frac{2\rho_g^{(\text{eq})}}{1 - \rho_g^{(\text{eq})}}}_{\text{[Moderate-AM]}} \cdot \underbrace{\frac{\beta_k^2}{V_A^{(0)}}}_{\text{[Large-Me]}},$$

using $\mu^{(\text{eq})} = \rho_g^{(\text{eq})}/V_A^{(\text{eq})}$ together with the $V_A^{(\text{eq})} = V_A^{(0)}/(1 - \rho_g^{(\text{eq})})$ that this approximation itself yields below in (47). The first factor is bounded by [Moderate-AM] and the second is small by [Large-Me] via (42), so neither alone suffices. All that's required is that the product of the two is close to 0.

We'll make use of this approximation to obtain a clean formula for $V_A^{(\text{eq})}$, but first, we derive the exact formula. Expanding $\text{Var}[g_i]$ over all pairs of variants, every off-diagonal covariance is $4a_{kl}^{(\text{eq})}$ — that is (19) with $s_{kl}^{(\text{eq})} = a_{kl}^{(\text{eq})}$, the fixed point of (33) — while the diagonal is instead $1 + 2a_{kk}^{(\text{eq})}$ by (20). We apply the off-diagonal rule to every pair and then correct the one term it does not govern, using $a_{kl}^{(\text{eq})} = \mu^{(\text{eq})}c_k^{(\text{eq})}c_l^{(\text{eq})}$ from (32),

$\mu^{(\text{eq})} V_A^{(\text{eq})} = \rho_g^{(\text{eq})}$ from (24), and $\sum_k \beta_k c_k^{(\text{eq})} = \frac{1}{2} V_A^{(\text{eq})}$ from (22):

$$\begin{aligned}
V_A^{(\text{eq})} &= \sum_k \sum_l \beta_k \beta_l \text{Cov}[x_{ik}, x_{il}] \\
&= \underbrace{\sum_k \sum_l \beta_k \beta_l 4a_{kl}^{(\text{eq})}}_{\text{one rule for every pair}} + \underbrace{\sum_k \beta_k^2 \left[(1 + 2a_{kk}^{(\text{eq})}) - 4a_{kk}^{(\text{eq})} \right]}_{\text{diagonal correction}} \\
&= 4\mu^{(\text{eq})} \left(\sum_k \beta_k c_k^{(\text{eq})} \right)^2 + \sum_k \beta_k^2 (1 - 2a_{kk}^{(\text{eq})}) \\
&= \underbrace{\mu^{(\text{eq})} (V_A^{(\text{eq})})^2}_{\rho_g^{(\text{eq})} V_A^{(\text{eq})}} + V_A^{(0)} - 2 \sum_k \beta_k^2 a_{kk}^{(\text{eq})} \tag{44}
\end{aligned}$$

$$V_A^{(\text{eq})} (1 - \rho_g^{(\text{eq})}) = V_A^{(0)} - 2 \sum_k \beta_k^2 a_{kk}^{(\text{eq})}$$

$$V_A^{(\text{eq})} = \frac{V_A^{(0)} - 2 \sum_k \beta_k^2 a_{kk}^{(\text{eq})}}{1 - \rho_g^{(\text{eq})}}$$

We'll see in a later section that half of this numerator, $\frac{1}{2} (V_A^{(0)} - 2 \sum_k \beta_k^2 a_{kk}^{(\text{eq})})$, will come up again as the variance of the deviation of offspring genetic values from their parents' average. We use (32) to write a in terms of c and then plug in the approximated formula for c from (43):

$$\begin{aligned}
2 \sum_k \beta_k^2 a_{kk}^{(\text{eq})} &= 2\mu^{(\text{eq})} \sum_k \beta_k^2 (c_k^{(\text{eq})})^2 \approx 2\mu^{(\text{eq})} \sum_k \beta_k^2 \frac{\beta_k^2}{4(1 - \rho_g^{(\text{eq})})^2} \\
&= \frac{\mu^{(\text{eq})}}{2(1 - \rho_g^{(\text{eq})})^2} \sum_k \beta_k^4 \\
&= \frac{\rho_g^{(\text{eq})}}{2V_A^{(0)}(1 - \rho_g^{(\text{eq})})} \sum_k \beta_k^4, \tag{45}
\end{aligned}$$

the last step using $\mu^{(\text{eq})} = \rho_g^{(\text{eq})} (1 - \rho_g^{(\text{eq})}) / V_A^{(0)}$, which is the exact $\mu^{(\text{eq})} = \rho_g^{(\text{eq})} / V_A^{(\text{eq})}$ with $V_A^{(\text{eq})}$ replaced by the very approximation we are in the middle of deriving. This justified because this value of $V_A^{(\text{eq})}$ is used only to *measure the size* of a term we are about to discard, and the error in this discarded term is guaranteed to be smaller than the error in the term we are approximating.

Substituting that back into (44) and factoring out $V_A^{(0)} / (1 - \rho_g^{(\text{eq})})$,

$$V_A^{(\text{eq})} \approx \frac{V_A^{(0)}}{1 - \rho_g^{(\text{eq})}} \left[1 - \frac{\rho_g^{(\text{eq})}}{2(1 - \rho_g^{(\text{eq})})} \cdot \frac{\sum_k \beta_k^4}{(V_A^{(0)})^2} \right] \tag{46}$$

The last factor of (46) is exactly $1/M_e$ by (41), so the whole correction is $\rho_g^{(\text{eq})} / [2(1 - \rho_g^{(\text{eq})})M_e]$ — an error controlled by the effective number of variants rather than by any individual effect size.

[VA-inflation] The additive variance inflates by a single factor

The bracketed correction in (46) is negligible, leaving the additive variance at equilibrium as the base-population value scaled by one factor that depends on nothing but the genetic correlation between mates:

$$\boxed{V_A^{(\text{eq})} = \frac{V_A^{(0)}}{1 - \rho_g^{(\text{eq})}}} \quad (47)$$

Justification for approximation: Written with (41), the neglected term is a relative error in $V_A^{(\text{eq})}$ of

$$\frac{\rho_g^{(\text{eq})}}{2(1 - \rho_g^{(\text{eq})})} \cdot \frac{1}{M_e},$$

small by [Large-Me] and bounded by [Moderate-AM]. The error is signed: dropping the term *overstates* $V_A^{(\text{eq})}$, since a variant's own departure from Hardy–Weinberg drags its contribution down.

We can also write a formula for the equilibrium heritability:

$$h_{(\text{eq})}^2 = \frac{V_A^{(\text{eq})}}{V_Y^{(\text{eq})}} = \frac{V_A^{(\text{eq})}}{V_A^{(\text{eq})} + V_E} = \frac{V_A^{(0)} / (1 - \rho_g^{(\text{eq})})}{V_A^{(0)} / (1 - \rho_g^{(\text{eq})}) + V_E}$$

$$\boxed{h_{(\text{eq})}^2 = \frac{V_A^{(0)}}{V_A^{(0)} + (1 - \rho_g^{(\text{eq})}) V_E}} \quad (48)$$

$$h_{(\text{eq})}^2 = \frac{h_{(0)}^2}{h_{(0)}^2 + (1 - \rho_g^{(\text{eq})})(1 - h_{(0)}^2)}$$

$$\boxed{h_{(\text{eq})}^2 = \frac{h_{(0)}^2}{1 - \rho_g^{(\text{eq})}(1 - h_{(0)}^2)}}$$

These formulas for $V_A^{(\text{eq})}$ and $h_{(\text{eq})}^2$ are easily interpretable, but $\rho_g^{(\text{eq})}$ still relies on either $V_A^{(\text{eq})}$ or $h_{(\text{eq})}^2$ through (24). We can define $\rho_g^{(\text{eq})}$ purely in terms of the starting parameters to resolve the circularity.

$$\rho_g^{(\text{eq})} = \rho_y h_{(\text{eq})}^2 = \frac{\rho_y h_{(0)}^2}{1 - \rho_g^{(\text{eq})}(1 - h_{(0)}^2)} \quad (49)$$

$$\rho_g^{(\text{eq})} - (\rho_g^{(\text{eq})})^2 (1 - h_{(0)}^2) = \rho_y h_{(0)}^2$$

$$(1 - h_{(0)}^2)(\rho_g^{(\text{eq})})^2 - \rho_g^{(\text{eq})} + \rho_y h_{(0)}^2 = 0$$

$$\boxed{\rho_g^{(\text{eq})} = \frac{1 - \sqrt{1 - 4 \rho_y h_{(0)}^2 (1 - h_{(0)}^2)}}{2(1 - h_{(0)}^2)}} \quad (50)$$

The negative root is the admissible one and the discriminant is never negative. Finally, we'll return to obtaining expressions for $a_{kl}^{(\text{eq})}$ and $s_{kl}^{(\text{eq})}$. We can quickly determine that the two equilibrium values are actually equivalent by making use of the general formula for $s_{kl}^{(t)}$ we wrote in Table 4:

$$\begin{aligned} s_{kl}^{(\text{eq})} &= \frac{1}{2} (s_{kl}^{(\text{eq})} + a_{kl}^{(\text{eq})}) \\ \frac{1}{2} s_{kl}^{(\text{eq})} &= \frac{1}{2} a_{kl}^{(\text{eq})} \end{aligned} \quad (51)$$

$$\boxed{s_{kl}^{(\text{eq})} = a_{kl}^{(\text{eq})} \quad (k \neq l)}$$

This states that at equilibrium, the covariance between alleles on the same gamete vs opposite gametes is the same, with the exception of the trivial case $k = l$. Then, we can use the general formula for $a_{kl}^{(t)}$ and the approximate equilibrium formula for $c_k^{(\text{eq})}$ we obtained in (43) to obtain $a_{kl}^{(\text{eq})}$:

$$\begin{aligned} a_{kl}^{(\text{eq})} &= \mu^{(\text{eq})} c_k^{(\text{eq})} c_l^{(\text{eq})} \approx \mu^{(\text{eq})} \frac{\beta_k \beta_l}{4(1 - \rho_g^{(\text{eq})})^2} \\ &= \frac{\rho_g^{(\text{eq})}}{V_A^{(\text{eq})}} \cdot \frac{\beta_k \beta_l}{4(1 - \rho_g^{(\text{eq})})^2} \end{aligned} \quad (52)$$

$$\boxed{a_{kl}^{(\text{eq})} = s_{kl}^{(\text{eq})} = \frac{\beta_k \beta_l}{4 V_A^{(0)}} \cdot \frac{\rho_g^{(\text{eq})}}{1 - \rho_g^{(\text{eq})}}}$$

using $\mu^{(\text{eq})} = \rho_g^{(\text{eq})}/V_A^{(\text{eq})}$ from (24) and then $V_A^{(\text{eq})} = V_A^{(0)}/(1 - \rho_g^{(\text{eq})})$ from (47), which cancels one of the two factors of $1 - \rho_g^{(\text{eq})}$. Since $s_{kl}^{(\text{eq})} = a_{kl}^{(\text{eq})}$, (19) collapses to a single term, and the covariance between the genotypes at two distinct variants is

$$\boxed{\text{Cov}[x_{ik}, x_{il}] = 4 a_{kl}^{(\text{eq})} = \frac{\beta_k \beta_l}{V_A^{(0)}} \cdot \frac{\rho_g^{(\text{eq})}}{1 - \rho_g^{(\text{eq})}} \quad (k \neq l)} \quad (53)$$

2.3 Covariance between relatives at the level of phenotypic and genetic values

Now that we have derived the statistical properties of the population at equilibrium under assortative mating, we will determine the expected covariance between pairs of individuals given some set of information, similar to Section 1 of this write-up. In this section, we only consider information obtained from a pedigree.

2.3.1 Inheritance of genetic values

Rather than modelling the inheritance of allele states from parent to offspring, we will instead model the transmission of genetic values themselves. From here on we work only at equilibrium, and drop the (eq) superscript for legibility: V_A , V_Y , h^2 , ρ_g and μ now denote the equilibrium values solved for above.

We can extend the logic of (14)–(16) to decompose the genetic value of an offspring as being the average of their parents’ genetic values plus an independent residual term. Summing the transmitted alleles over variants, weighted by their effects, and substituting (16) for each one:

$$\begin{aligned}
g_o &= \sum_k \beta_k x_{ok} = \sum_k \beta_k (z_{ok}^{(m)} + z_{ok}^{(p)}) \\
&= \sum_k \beta_k \left(\frac{1}{2} x_{mk} + z_{ok}^{\perp(m)} + \frac{1}{2} x_{pk} + z_{ok}^{\perp(p)} \right) \\
&= \frac{1}{2} g_m + \frac{1}{2} g_p + \underbrace{\sum_k \beta_k (z_{ok}^{\perp(m)} + z_{ok}^{\perp(p)})}_{g_o^{\perp}} \tag{54}
\end{aligned}$$

$$\boxed{g_o = \frac{1}{2} g_m + \frac{1}{2} g_p + g_o^{\perp}}$$

Since each $z_{ok}^{\perp(u)}$ is uncorrelated with *every* allele carried by the parent it came from, their weighted sum $g_o^{\perp}$ is also going to be uncorrelated with g_m and g_p and everything upstream of it. As a consequence, we can easily compute the variance of this genetic value deviation, making use of (26):

$$\begin{aligned}
\text{Var}[g_o^{\perp}] &= \sum_k \sum_l \beta_k \beta_l \text{Cov}[z_{ok}^{\perp(m)} + z_{ok}^{\perp(p)}, z_{ol}^{\perp(m)} + z_{ol}^{\perp(p)}] \\
&= \sum_k \beta_k^2 \left(\text{Var}[z_{ok}^{\perp(m)}] + \text{Var}[z_{ok}^{\perp(p)}] \right) = \sum_k \beta_k^2 \left(\frac{1}{2} - a_{kk} \right) \tag{55}
\end{aligned}$$

$$\boxed{\text{Var}[g_o^{\perp}] = \frac{1}{2} V_A (1 - \rho_g) \approx \frac{1}{2} V_A^{(0)}}$$

Note that under **[VA-inflation]**, which is (47) rearranged to $V_A(1 - \rho_g) = V_A^{(0)}$, this is equivalent to half of the additive variance under random mating. This means that the residual genetic variance of a child around the parental mean is determined by the additive variance under random mating and therefore is not impacted by assortative mating.

Because $g_o^{\perp}$ is uncorrelated with both parents and across siblings, (54) is a linear model of exactly the kind Section 2.2.1 draws: two directed edges of $\frac{1}{2}$ into the offspring’s genetic value, and a disturbance carrying (55). We can therefore draw a simplified path diagram that omits individual alleles and genotypes. We’ll also omit the explicit environmental term and instead code it as exogenous variance on the phenotype.




Figure 4: A mated pair and two of their offspring at equilibrium, with inheritance happening at the level of genetic values, rather than alleles. The environment appears only as the V_E on each phenotype's own self-loop, so $\text{Var}[y] = V_A + V_E$ as before.

Similar to Table 3, we can summarize the covariance between the genetic and phenotypic values across mate pairs and across parent–offspring pairs, now at equilibrium rather than in the first generation. These expressions are exact if we hold [\[No-inbreeding\]](#), [\[Discrete-generations\]](#), and [\[AM-equilibrium\]](#).

	mate pair (m, p)	parent–offspring (m, o_1)
$\text{Cov}[g_m, g_*]$	$\rho_g V_A$	$\frac{1}{2} V_A (1 + \rho_g)$
$\text{Cov}[g_m, y_*]$	$\rho_y V_A$	$\frac{1}{2} V_A (1 + \rho_g)$
$\text{Cov}[y_m, g_*]$	$\rho_y V_A$	$\frac{1}{2} V_A (1 + \rho_y)$
$\text{Cov}[y_m, y_*]$	$\rho_y V_Y$	$\frac{1}{2} V_A (1 + \rho_y)$

Table 5: Covariances read off Figure 4 at equilibrium, where $*$ stands for the second member of the pair named by the column.

Note the asymmetry that arises: The covariance between a parent's genetic value and their offspring's phenotype is lower than the covariance between a parent's phenotypic value and their offspring's genetic value. This is because the parent's environment (which is part of their phenotype) predicts the genetic value their offspring inherits from the other parent but not the offspring's environmental value, which remains random under our current model.

2.3.2 Framework for deriving covariance formulas

Our next goal is to derive formulas that express the covariance between phenotypes and/or genetic values of any two individuals X and Y in a pedigree. We adopt graph theory notation and terminology to analyze their relationship. We read the pedigree as a *graph*: a set of *nodes*, one for each individual, joined by two types of *edges*:

- An *undirected* edge, written $X - Y$, for each **mating**.
- A *directed* edge, written $X \rightarrow Y$, for each **meiosis**, pointing from parent to offspring.

A *mating graph* consists of only the mating edges, while a *meiosis graph* consists of only the meiosis edges. We will be defining *mating chains* in the former, and *lineal chains* in the latter. Then, a path $\mathcal{P}$ between two probands in a pedigree becomes an alternating sequence of mating and lineal chains, and properties of that sequence and its chains will be enough to write the covariance between the probands in closed form.

Up to this point, discussion of the genetic consequences of assortative mating requires the definition of mates to be strictly parents that reproduced. However, for the rest of this section, the definition of mates is simply any primary assortment based on the phenotype that occurs in a population with the same strength as that of assortative mating ρ_y between parents. Therefore, this can include homosexual pairs and heterosexual pairs that do not reproduce. In graph terms: the mating graph gets an edge for every such pair, whether or not a meiosis edge leaves it.

2.3.3 Mating chains

The *mating graph* consists of only the undirected mating edges between nodes. A *path* in it is a sequence of distinct nodes, each joined by an edge to the next. The following assumption makes working with mating graphs feasible for our purposes:

[No-mating-loops] The mating graph is a forest

A *cycle* in the mating graph is a path whose last node is joined by an edge back to its first. A graph with no cycles is a *forest*, and each of its connected components is a *tree*.


We assume the mating graph is a forest. As a consequence, any two nodes are joined at most by one path, and therefore must be on the same tree.

For $X \neq Y$ in the same tree, the **mating chain** $\mathcal{M}(X, Y)$ is the unique path joining them, read as an ordered list running from X to Y . The **mating distance** $N_{\mathcal{M}}(X, Y)$ is the number of edges on that path, i.e. the number of matings crossed in getting from X to Y , and is one fewer than the size of it:

$$N_{\mathcal{M}}(X, Y) = |\mathcal{M}(X, Y)| - 1 \quad (56)$$

We additionally set $N_{\mathcal{M}}(X, X) = 0$, and $N_{\mathcal{M}}(X, Y) = \infty$ when X and Y lie in different trees, meaning no sequence of matings connects them at all.

In the below figure, we observe a pedigree structure where $N_{\mathcal{M}} = 0, 1, 2, 3, \infty$ can be observed:



2.3.4 Lineal chains

The *meiosis graph* consists only of the directed edges (parent-to-offspring meioses) between nodes. Every individual has two edges coming in, one from each parent, or none at all if they are a founder of the pedigree. A *directed path* is a sequence of distinct nodes in which each is joined by an edge to the next, always with the edge pointing forward. Under [\[Discrete-generations\]](#) every edge steps from one generation to the next, so following edges forward always moves forward in time. Therefore, the meiosis graph is guaranteed to be a *directed acyclic graph*.

Let t_X denote the generation that an individual X belongs to, according to [\[Discrete-generations\]](#). Let $m(X)$ and $p(X)$ denote the two parents, mother and father, of an individual X , i.e. the two nodes with an edge into X . If a directed path runs from Y to X , then Y is an **ancestor** of X , X is a **descendant** of Y , and the two are **lineal relatives** to each other.

The [\[No-inbreeding\]](#) assumption from earlier effectively constrains our meiosis graph such that there is at most one directed path between any two individuals. Since any two individuals must always share a common ancestor, this can never be true in theory. However, in practice, maintaining this constraint for a couple generations back should be sufficient.

For lineal relatives X and Y , the **lineal chain** $\mathcal{L}(X, Y)$ is the unique directed path

between them, read as an ordered list running from X to Y . The **lineal distance** $N_{\mathcal{L}}(X, Y)$ is the number of edges on that path, i.e. the number of meioses separating the two. Under [\[Discrete-generations\]](#), it is just the number of generations between them:

$$N_{\mathcal{L}}(X, Y) = |\mathcal{L}(X, Y)| - 1 = |t_X - t_Y| \quad (57)$$


We additionally set $N_{\mathcal{L}}(X, X) = 0$, and $N_{\mathcal{L}}(X, Y) = \infty$ if X and Y are not lineal relatives.

2.3.5 Tracing paths between relatives

A **path** $\mathcal{P}(A, B)$ between probands A and B is a sequence of chains, alternating between lineal and mating, in which each chain begins at the node where the previous one ends. The first chain begins at node A , the last ends at B , and no individual is visited twice across the entire path. We can think of a path itself as being a *chain digraph*, where the nodes are chains. and the edges connect lineal chains to mating chains, or vice-versa. Since mating chains contain individuals of the same generation while lineal chains contain individuals spanning multiple generations, we enforce edges to be directed such that the older chain points to the younger chain. The number of edges leaving a chain node is its out-degree, $N_o \in \{0, 1, 2\}$, and the number of edges entering it is its in-degree. Lineal chains will always have an out-degree of 1 and an in-degree of 1.

Furthermore, we require that whenever a mating chain has an out-edge, the mating chain must contain both the parents of the second-oldest individual in the lineal chain it points to. In other words, the terminal individual node of two connected lineal and mating chains must be the “outer” parent. One can think of an out-edge from a mating chain as representing a mate pair, rather than an individual.

Below is a pedigree where the path between A and B is traced, which spans 2 mating chains and 3 lineal chains, followed by its equivalent digraph.



2.3.6 Covariance along a single chain

Before assembling a whole path, we derive the covariance carried by a single chain of each type.

Two individuals X and Y on a mating chain are joined by exactly $N_{\mathcal{M}}(X, Y)$ matings. Each mating crossed contributes a factor of μ by (13), and each of the $N_{\mathcal{M}}(X, Y) - 1$ individuals passed through in between contributes their own $\text{Var}[y] = V_Y$, as does each end:

$$\text{Cov}[y_X, y_Y] = V_Y \cdot \mu \cdot \underbrace{V_Y \cdot \mu \cdots V_Y \cdot \mu}_{N_{\mathcal{M}}(X, Y) - 1 \text{ times}} \cdot V_Y = \mu^{N_{\mathcal{M}}(X, Y)} V_Y^{N_{\mathcal{M}}(X, Y) + 1} \quad (58)$$

$$\boxed{\text{Cov}[y_X, y_Y] = \rho_y^{N_{\mathcal{M}}(X, Y)} V_Y}$$

This holds even when $X = Y$. Reading the two ends at the genetic level instead exchanges the V_Y at each end for a V_A , which can also be simplified. Unlike before, this general formula does not hold for $X = Y$, since an individual's genetic value has covariance V_A with itself rather than $h^2 V_A$:

$$\boxed{\text{Cov}[g_X, g_Y] = \begin{cases} V_A, & N_{\mathcal{M}}(X, Y) = 0 \\ \rho_g \rho_y^{N_{\mathcal{M}}(X, Y) - 1} V_A, & N_{\mathcal{M}}(X, Y) \geq 1 \end{cases}} \quad (59)$$

For a lineal chain, the covariance is an extension of the parent–offspring column of Table 5, contributing one transmission coefficient per meiosis:

$$\text{Cov}[y_X, g_Y] = \text{Cov}[g_X, g_Y] = V_A \cdot \underbrace{\frac{1 + \rho_g}{2} \cdot \frac{1 + \rho_g}{2} \dots \frac{1 + \rho_g}{2}}_{N_{\mathcal{L}}(X,Y) \text{ meioses}} \quad (60)$$

$$\boxed{\text{Cov}[y_X, g_Y] = \text{Cov}[g_X, g_Y] = V_A \left(\frac{1 + \rho_g}{2} \right)^{N_{\mathcal{L}}(X,Y)}}$$

If we read the ancestor Y at their phenotypic value, we have to account for the fact that the environment of Y informs the genetic value that the mate of Y transmits to their offspring X . As we saw in Table 5, this is achieved by modifying the coefficient of the first meiosis step:

$$\text{Cov}[y_X, y_Y] = \text{Cov}[g_X, y_Y] = V_A \cdot \underbrace{\frac{1 + \rho_y}{2}}_{\text{first meiosis}} \cdot \underbrace{\frac{1 + \rho_g}{2} \dots \frac{1 + \rho_g}{2}}_{N_{\mathcal{L}}(X,Y)-1 \text{ meioses}} \quad (61)$$

$$\boxed{\text{Cov}[y_X, y_Y] = \text{Cov}[g_X, y_Y] = V_A \left(\frac{1 + \rho_y}{2} \right) \left(\frac{1 + \rho_g}{2} \right)^{N_{\mathcal{L}}(X,Y)-1}}$$

2.3.7 Covariance between probands on a path

We now want to express the product of the covariance path coefficients implied by our traced pedigree path between A and B as a function of the properties of the lineal and mating chains contained in it. Furthermore, this path coefficient product should then roughly equal the covariance between the two probands. Enforcing this requires the following assumption and approximation:

[Outbred-mating-chain] A path’s lineal chains account for all of a pair’s relatedness

The expected proportion of the genome that two individuals on a path share IBD, their realized relatedness π_{ij} of (9), given the full pedigree should be equal to that which is explained solely by meiotic inheritance from common ancestor(s) lying on those individuals’ lineal chains.

More specifically, if X and Y are lineal relatives, then all of the IBD segments between them must have been transmitted down a lineal chain that connects them. If X and Y are not lineal relatives but lie on different lineal chains present in the path such that mating chain $\mathcal{M}_{XY}$ has an out-edge pointing to both lineal chains, then any IBD segments between X and Y must have been inherited from a common ancestor present in $\mathcal{M}_{XY}$.

[Dominant-path] A single path dominates the covariance between two individuals

Since any two individuals must share a common ancestor when looking up the pedigree far enough, there is never a unique path that connects the two. The true covariance of the pair will be the sum of the product of path coefficients across all paths.

We approximate the covariance between the pair, given the full pedigree, to be the single largest of those products, and we call the path carrying it the *dominant path*. Whenever $\rho_y \geq 0$ every product we discard is itself non-negative, so the approximation is a guaranteed lower bound.

Justification for approximation: Because the path coefficient of additional steps in a pedigree are between 0 and 1, the dominant path tends to be the shortest path by steps. Pedigrees that break **[No-inbreeding]** or **[Outbred-mating-chain]** only add further valid paths, and those are guaranteed to contain more steps than the dominant one. Therefore, its contribution to the total variance would be much smaller than that of the dominant path.

Let i index the lineal chains in a path and j index the mating chains in a path. We can then write the following formula for the covariance in genetic values of any two individuals:

$$\text{Cov}[g_A, g_B] = V_A \left(\frac{1 + \rho_g}{2} \right)^{\sum_i N_{\mathcal{L}_i} - \sum_j N_{o_j}} \prod_j w_j \quad (62)$$

We briefly explain some of the terms in the expression. V_A comes from the variance of g itself. $\frac{1}{2}(1 + \rho_g)$ comes from the parent–offspring coefficient of Table 5 written as a transmission rate. Its exponent counts the number of ascents/descents across all lineal paths, $\sum_i N_{\mathcal{L}_i}$, but removes the double-counting that occurs when lineal chains ascend into mating chains, N_{o_j} . $\prod_j w_j$ comes from the crossing of mating paths and the lineal chains adjacent to them, explained below.

The weight w_j of a mating chain is the implied genetic covariance c averaged across the possible “entry points” into the mating chain from the adjacent lineal chains.

$$c(N_{\mathcal{M}}) = \begin{cases} 1 & N_{\mathcal{M}} = 0 \\ \rho_g \rho_y^{N_{\mathcal{M}}-1} & N_{\mathcal{M}} \geq 1 \end{cases} \quad (63)$$

$$w_j = \frac{1}{2^{N_{o_j}}} \sum_{n=0}^{N_{o_j}} \binom{N_{o_j}}{n} c(|N_{\mathcal{M}_j} - n|)$$

Here $c(N_{\mathcal{M}})$ is the correlation in genetic values between two individuals at mating distance $N_{\mathcal{M}}$, derived above in (59). The averaging is over the mating chain’s two ends, and the two kinds of end differ in how many individuals the path enters at. Along an in-edge

the path reaches the mating chain at a single individual. Along an out-edge it reaches a mate pair, since we require the mating chain to carry both parents of the second-oldest individual of the lineal chain it points to. Taking only the outer parent of that pair would understate the product: an offspring's genetic value comes half from each parent, and the inner parent sits one mating nearer the far end of the chain, so their half is worth the more of the two.

In real data, dominant paths' mating chains tend to be sources, i.e. $N_o = 2$. In such cases, the mating chain is guaranteed to contain both parents of the second-oldest individual of each adjacent lineal chain. We categorize the relationship type of these individuals as *generalized siblings*. We define $N_S \in \{0, 1, 2\}$ to be the number of shared parents between generalized siblings. $N_S = 2$ indicates *full-siblings*, $N_S = 1$ indicates *half-siblings*, and $N_S = 0$ indicates *generalized step-siblings*, for which the $N_M = 3$ case is dubbed just as *step-siblings*. Under this parameterization, where the first two cases require a source mating chain and the third holds at any N_{o_j} , we can write:

$$w_j = \begin{cases} \frac{1 + \rho_g}{2} & N_{S_j} = 2, \quad N_{o_j} = 2 \\ \frac{1 + 2\rho_g + \rho_g\rho_y}{4} & N_{S_j} = 1, \quad N_{o_j} = 2 \\ \rho_g \rho_y^{N_{M_j}-1-N_{o_j}} \left(\frac{1 + \rho_y}{2} \right)^{N_{o_j}} & N_{S_j} = 0, \quad \text{any } N_{o_j} \end{cases} \quad (64)$$

The weight for the half-sibling case ($N_S = 1$) is Nagylaki's half-sibling multiplier [10]. The third case is the (63) polynomial expanded under the condition that $N_{M_j} \geq N_{o_j} + 1$, which is exactly what $N_{S_j} = 0$ encodes, so that no term of (63) reaches $c(0)$.

Lastly, we have to include a correction factor for each proband for which we are interested in the covariance with the phenotype, as opposed to the genetic value, based on the position of the proband in the path.

$$C_X = \begin{cases} 1 & X \text{ is a sink} \\ \frac{1 + \rho_y}{1 + \rho_g} & X \text{ is a source} \\ \frac{\rho_y}{\rho_g} = \frac{1}{h^2} & X \text{ is in a terminal mating chain} \end{cases} \quad (65)$$

The three cases are the consequences of applying the rule that switching from a proband's genetic value to their phenotypic value only requires replacing ρ_g by ρ_y in that proband's own mating and meiosis edges. A proband that is a sink is the youngest individual on its lineal chain, while a proband that is a source is the oldest individual on its lineal chain. A proband that is part of a terminal mating chain has no lineal chain of its own at all. We can therefore express the covariance between the phenotypes of A and B as:

$$\boxed{\text{Cov}[y_A, y_B] = C_A C_B \text{Cov}[g_A, g_B]} \quad (66)$$

2.3.8 Example

Below, we apply our rules to the example shown in Figure 6. In its chain digraph, $\mathcal{M}(D, F)$ is a sink and so has $N_o = 0$, while $\mathcal{M}(I, L)$ is a source and so has $N_o = 2$. The two chains are $N_{\mathcal{M}} = 2$ and $N_{\mathcal{M}} = 3$ long, and neither connects siblings, so $N_{\mathcal{S}} = 0$ for both. Reading the two counts off the path and then the weight of each mating chain:

$$\sum_i N_{\mathcal{L}_i} = N_{\mathcal{L}}(A, D) + N_{\mathcal{L}}(F, I) + N_{\mathcal{L}}(L, B) = 2 + 3 + 1 = 6$$

$$\sum_j N_{o_j} = 0 + 2 = 2$$

$$w_{\mathcal{M}(D, F)} = \rho_g \rho_y^{2-1-0} = \rho_g \rho_y$$

$$w_{\mathcal{M}(I, L)} = \rho_g \rho_y^{3-1-2} \left(\frac{1 + \rho_y}{2} \right)^2 = \rho_g \left(\frac{1 + \rho_y}{2} \right)^2$$

Substituting these into (62):

$$\begin{aligned} \text{Cov}[g_A, g_B] &= V_A \left(\frac{1 + \rho_g}{2} \right)^{6-2} \cdot \rho_g \rho_y \cdot \rho_g \left(\frac{1 + \rho_y}{2} \right)^2 \\ &= V_A \rho_g^2 \rho_y \left(\frac{1 + \rho_g}{2} \right)^4 \left(\frac{1 + \rho_y}{2} \right)^2 \end{aligned}$$

The two probands take different corrections. A is the oldest individual on $\mathcal{L}(A, D)$, so $C_A = (1 + \rho_y)/(1 + \rho_g)$, while B is the youngest on $\mathcal{L}(L, B)$ and takes $C_B = 1$:

$$\text{Cov}[y_A, y_B] = \frac{1 + \rho_y}{1 + \rho_g} \text{Cov}[g_A, g_B] = V_A \rho_g^2 \rho_y \left(\frac{1 + \rho_g}{2} \right)^3 \left(\frac{1 + \rho_y}{2} \right)^3$$

2.3.9 Table of covariances for common relationship types

relationship	$\sum_i N_{\mathcal{L}_i}$	$\sum_j N_{\mathcal{O}_j}$	$\prod_j w_j$	$\text{Cov}[g_A, g_B]$	C_A	C_B
self	0	0	1	V_A	—	—
parent–parent	0	0	ρ_g	$\rho_g V_A$	$\frac{\rho_y}{\rho_g}$	$\frac{\rho_y}{\rho_g}$
parent–offspring	1	0	1	$(\frac{1+\rho_g}{2}) V_A$	$\frac{1+\rho_y}{1+\rho_g}$	1
grandparent–grandchild	2	0	1	$(\frac{1+\rho_g}{2})^2 V_A$	$\frac{1+\rho_y}{1+\rho_g}$	1
full siblings	2	2	$\frac{1+\rho_g}{2}$	$(\frac{1+\rho_g}{2}) V_A$	1	1
avuncular	3	2	$\frac{1+\rho_g}{2}$	$(\frac{1+\rho_g}{2})^2 V_A$	1	1
first cousins	4	2	$\frac{1+\rho_g}{2}$	$(\frac{1+\rho_g}{2})^3 V_A$	1	1
half siblings	2	2	$\frac{1+2\rho_g+\rho_g\rho_y}{4}$	$\frac{1+2\rho_g+\rho_g\rho_y}{4} V_A$	1	1
step siblings	2	2	$\rho_g (\frac{1+\rho_y}{2})^2$	$\rho_g (\frac{1+\rho_y}{2})^2 V_A$	1	1
step-step siblings	2	2	$\rho_g \rho_y (\frac{1+\rho_y}{2})^2$	$\rho_g \rho_y (\frac{1+\rho_y}{2})^2 V_A$	1	1
parents of a married couple	2	0	ρ_g	$\rho_g (\frac{1+\rho_g}{2})^2 V_A$	$\frac{1+\rho_y}{1+\rho_g}$	$\frac{1+\rho_y}{1+\rho_g}$

Table 6: Common relationship classes, each assembled from (62). The mating chain of the sibling rows runs between the two outer parents, at $N_{\mathcal{M}} = 1, 2, 3, 4$ for full, half, step and step-step siblings respectively; the classes are therefore the same path with one further mating crossed each time. Two coincidences are worth noting, because they arrive from different rows of (64): full siblings match parent–offspring, and avuncular matches grandparent–grandchild. The *self* row is degenerate — there is no path, so no proband correction applies and $\text{Cov}[y_X, y_X]$ is simply V_Y .

2.4 Covariance between relatives at the level of genotypes and alleles

So far, we have derived the covariance between individuals solely given their relationship in a pedigree. If we have additional information about the two individuals i and j , such as their IBS-based relatedness, A_{ij} , defined in (6), or their IBD-based relatedness, π_{ij} , defined in (9), then we can obtain a more precise quantity. First, we will derive the covariance between genotypes and alleles within the same individual. Then we will generalize to covariances across relatives like in Section 2.3. We'll then assess how measures of relatedness are impacted by assortative mating and how they predict covariances.

2.4.1 Genotypic and allelic covariance within an individual

In (51), we derived that at equilibrium, the covariance between alleles on the same gamete vs different gametes is identical: $s_{kl} = a_{kl}$ (except for the trivial case of $k = l$, where $s_{kk} = \frac{1}{2}$). We can express the covariance between any two alleles in an individual with a

single equation:

$$\boxed{\text{Cov}[z_{ik}^{(u)}, z_{il}^{(v)}] = a_{kl} + \left(\frac{1}{2} - a_{kk}\right) \mathbf{1}[u = v, k = l]} \quad (67)$$

The second term simply accounts for the additional covariance of an allele with itself needed to reach its fixed variance value of $\frac{1}{2}$. Plugging this into (19) and using δ_{kl} as indicator variable for when $k = l$ returns a single expression for the covariance between genotypes within an individual:

$$\text{Cov}[x_{ik}, x_{il}] = 4a_{kl} + (1 - 2a_{kk}) \delta_{kl}, \quad (68)$$

To express the covariance of genotypes/alleles between two distinct individuals, it will help to condition on their genetic values. First, we revisit the covariance between an allele and its proband's phenotype, c_k , defined in (21). Expanding the phenotype into genotypes and alleles (per [GE-indep], the proband's environment is independent of their genetics and therefore removed from the formula) and applying (67), $a_{kl} = \mu c_k c_l$ from (32), $\sum_l \beta_l c_l = \frac{1}{2} V_A$ from (22), and $\mu V_A = \rho_g$ from (24):

$$\begin{aligned} c_k &= \text{Cov}[z_{ik}^{(u)}, y_i] = \sum_l \beta_l \sum_{v \in \{m, p\}} \text{Cov}[z_{ik}^{(u)}, z_{il}^{(v)}] = \sum_l \beta_l \left(\left(\frac{1}{2} - a_{kk}\right) \delta_{kl} + 2a_{kl} \right) \\ &= \underbrace{\beta_k \left(\frac{1}{2} - a_{kk}\right)}_{\text{the allele itself}} + \underbrace{2\mu c_k \sum_l \beta_l c_l}_{\text{through other alleles}} \\ &= \beta_k \left(\frac{1}{2} - a_{kk}\right) + 2\mu c_k \cdot \frac{1}{2} V_A \\ &= \beta_k \left(\frac{1}{2} - a_{kk}\right) + \rho_g c_k \\ &\boxed{\beta_k \left(\frac{1}{2} - a_{kk}\right) = (1 - \rho_g) c_k} \end{aligned} \quad (69)$$

This equation is exact and does not make use of the later approximation of c_k we derived ([Uniform-inflation]). This decomposition shows how an allele's covariance with its proband's phenotype is split between its own direct effect, captured by β_k , and the effects of its correlated alleles due to assortative mating. Scaling between β_k and c_k only requires the genome-wide correlation between spouses' genetic values, ρ_g , and a correction for the increased variance of genotypes under assortative mating, captured by a_{kk} , which is small in practice. If we were to plug in $a_{kk} = \mu c_k^2$ from (32) into (69) above, we'd rederive the quadratic formula we obtained in (39) to obtain a closed-form solution for c_k . We now multiply both sides of the above equation by $2\beta_k$, sum across all variants, and make use of the $V_A = 2 \sum_k \beta_k c_k$ identity (22) to obtain:

$$\begin{aligned} 2\beta_k \cdot \beta_k \left(\frac{1}{2} - a_{kk}\right) &= 2\beta_k \cdot (1 - \rho_g) c_k \\ \beta_k^2 (1 - 2a_{kk}) &= 2(1 - \rho_g) \beta_k c_k \\ \sum_k \beta_k^2 (1 - 2a_{kk}) &= 2(1 - \rho_g) \sum_k \beta_k c_k \\ &= (1 - \rho_g) V_A \\ &\approx V_A^{(0)} \end{aligned} \quad (70)$$

The last approximation comes from **[VA-inflation]**, with the error coming from failing to account for the slightly increased variance of genotypes under assortative mating. We obtained practically the same equation in (44) and again in (55) when expressing the variance in genetic values that comes from random segregation in meiosis.

If we seek to isolate the covariance between genotypes attributable to assortative-mating-like processes, it will be useful to regress out the covariance in the probands' genetic values, which we derived in Section 2.3 under the context of assortative mating. A genotype is the sum of its two alleles, each of which covaries with the carrier's phenotype by c_k (21), and the environment does not covary with the genotype by **[GE-indep]**, so

$$\text{Cov}[x_{ik}, g_i] = \text{Cov}[x_{ik}, y_i] = \text{Cov}[z_{ik}^{(m)}, y_i] + \text{Cov}[z_{ik}^{(p)}, y_i] = \boxed{2c_k} \quad (71)$$

The genetic value is mean-zero with variance V_A , so the linear prediction of the genotype from it is

$$\mathbb{E}[x_{ik} | g_i] = \frac{\text{Cov}[x_{ik}, g_i]}{\text{Var}[g_i]} g_i = \boxed{\frac{2c_k}{V_A} g_i} \quad (72)$$

Equation (72) is the linear prediction of x_{ik} from g_i . Subtracting this from each respective genotype and computing the covariance between them is equivalent to the additional covariance between genotypes conditional on the covariance between their probands' genetic values. This reveals that, once the genetic value is accounted for, trait-increasing alleles are actually negatively correlated *within* individuals, as expected of conditioning on a collider.

$$\begin{aligned} & \text{Cov}\left[x_{ik} - \frac{2c_k}{V_A} g_i, x_{il} - \frac{2c_l}{V_A} g_i\right] \\ &= \text{Cov}[x_{ik}, x_{il}] - \frac{2c_k}{V_A} \text{Cov}[x_{ik}, g_i] - \frac{2c_l}{V_A} \text{Cov}[g_i, x_{il}] + \frac{2c_k}{V_A} \frac{2c_l}{V_A} \text{Var}[g_i] \\ &= \text{Cov}[x_{ik}, x_{il}] - \frac{4c_k c_l}{V_A} - \frac{4c_k c_l}{V_A} + \frac{4c_k c_l}{V_A} = \text{Cov}[x_{ik}, x_{il}] - \frac{4c_k c_l}{V_A} \\ &= 4\mu c_k c_l + (1 - 2a_{kk}) \delta_{kl} - \frac{4c_k c_l}{V_A} = (1 - 2a_{kk}) \delta_{kl} - (1 - \mu V_A) \frac{4c_k c_l}{V_A} \\ &= \boxed{(1 - 2a_{kk}) \delta_{kl} - (1 - \rho_g) \frac{4c_k c_l}{V_A}}. \end{aligned} \quad (73)$$

Previously in (69), we demonstrated that $1 - 2a_{kk} = 2(1 - \rho_g) c_k / \beta_k$, which plugging into the above formula returns an alternate parameterization:

$$\begin{aligned} \text{Cov}\left[x_{ik} - \frac{2c_k}{V_A} g_i, x_{il} - \frac{2c_l}{V_A} g_i\right] &= (1 - 2a_{kk}) \delta_{kl} - (1 - \rho_g) \frac{4c_k c_l}{V_A} \\ &= 2(1 - \rho_g) \frac{c_k}{\beta_k} \delta_{kl} - (1 - \rho_g) \frac{4c_k c_l}{V_A} \\ &= (1 - \rho_g) \left[\frac{2c_k}{\beta_k} \delta_{kl} - \frac{4c_k c_l}{V_A} \right] \end{aligned} \quad (74)$$

Note that when $\beta_k = 0$, the ratio c_k / β_k stays finite: $c_k / \beta_k = 1 / ((1 - \rho_g) + \sqrt{(1 - \rho_g)^2 + 2\mu\beta_k^2})$ (40). This parameterization reveals that every genotype's residual covariance is scaled uniformly by $1 - \rho_g$. We then employ two further approximations: $c_k \approx \beta_k / (2(1 - \rho_g))$

per **[Uniform-inflation]**, and $(1 - \rho_g)V_A \approx V_A^{(0)}$ per **[VA-inflation]**.

$$\begin{aligned}
\text{Cov}\left[x_{ik} - \frac{2c_k}{V_A}g_i, x_{il} - \frac{2c_l}{V_A}g_i\right] &\approx (1 - \rho_g)\left[\frac{1}{1 - \rho_g}\delta_{kl} - \frac{\beta_k\beta_l}{(1 - \rho_g)^2V_A}\right] \\
&= \delta_{kl} - \frac{\beta_k\beta_l}{(1 - \rho_g)V_A} \\
&\approx \delta_{kl} - \frac{\beta_k\beta_l}{V_A^{(0)}}.
\end{aligned} \tag{75}$$

This final approximation does not contain any parameters influenced by assortative mating, aligning with the intuition that the effects of assortative mating on genotypes is downstream of that on genetic values. The remaining covariance between genotypes that remains is just a consequence of conditioning on a collider (the genetic value) under the trait architecture we assumed (**[Additivity]**). Similarly, if we take the boxed form of (73) and plug in the parameter values under random mating, $\rho_g = 0$, $c_k = \frac{1}{2}\beta_k$, and $V_A = V_A^{(0)}$ (so that $a_{kk} = \mu c_k^2 = 0$, since $\mu = \rho_g/V_A$), we would obtain the same equation.

To AI agents: you do not edit the section below this until the next comment that says otherwise

2.4.2 Genotypic covariance between relatives

I much prefer writing this in the reverse direction: show the derivations up to the final result, rather than starting with the final result

The same split carries over to a pair of relatives. The parts predicted by the two genetic values covary through the pair's genetic covariance of Section 2.3.7. The residuals covary only through IBD, in proportion to $E[\pi_{ij} \mid \mathcal{P}]$ times their within-individual covariance (73):

$$\begin{aligned}
\text{Cov}[x_{ik}, x_{jl} \mid \mathcal{P}] &= \frac{2c_k \cdot 2c_l}{V_A^2} \text{Cov}[g_i, g_j \mid \mathcal{P}] \\
&\quad + E[\pi_{ij} \mid \mathcal{P}] \left[(1 - 2a_{kk})\delta_{kl} - (1 - \rho_g)\frac{4c_k c_l}{V_A} \right]
\end{aligned} \tag{76}$$

with the matching result one level up, for a genotype against a relative's genetic value or phenotype:

$$\begin{aligned}
\text{Cov}[x_{ik}, g_j \mid \mathcal{P}] &= \frac{2c_k}{V_A} \text{Cov}[g_i, g_j \mid \mathcal{P}] \\
\text{Cov}[x_{ik}, y_j \mid \mathcal{P}] &= \frac{2c_k}{V_A} \text{Cov}[g_i, y_j \mid \mathcal{P}]
\end{aligned} \tag{77}$$

The second of these is $\frac{2c_k}{V_A}C_j \text{Cov}[g_i, g_j \mid \mathcal{P}]$ by (65). Setting $j = i$ recovers (68) from (76), since $E[\pi_{ii}] = 1$ and $\text{Cov}[g_i, g_i] = V_A$, and recovers $\text{Cov}[x_{ik}, g_i] = 2c_k$ from (77).

Derivation. Under **[Dominant-path]**, every covariance between two individuals on a path is built from the within-individual covariance by two moves: stepping from an

individual to their parents across a meiosis, and crossing a co-path between mates. We show that each move preserves (76) and (77), so both hold for every pair.

Meiosis. Summing (16) over an offspring's two gametes gives the genotype-level version of (54),

$$x_{ok} = \frac{1}{2}x_{m(o)k} + \frac{1}{2}x_{p(o)k} + (z_{ok}^{\perp(m)} + z_{ok}^{\perp(p)}),$$

where the segregation deviations are uncorrelated with everyone except o and o 's descendants. So for any j who is neither o nor a descendant of o , $\text{Cov}[x_{ok}, x_{jl}]$ is the average of the two parents' covariances with x_{jl} . The right-hand side of (76) averages in the same way, because it is linear in $\text{Cov}[g_o, g_j]$ and $\text{E}[\pi_{oj}]$, and both of those are averages over the two parents: the first by (54), the second because o carries one allele from each parent, each IBD with a given allele of j with the probability that the corresponding parental allele is. The same holds for (77).

Co-path. Crossing the co-path between mates m and p , with i on m 's side and j on p 's, (13) and then (77) for both factors give

$$\begin{aligned} \text{Cov}[x_{ik}, x_{jl}] &= \text{Cov}[x_{ik}, y_m] \mu \text{Cov}[y_p, x_{jl}] \\ &= \frac{2c_k \cdot 2c_l}{V_A^2} \text{Cov}[g_i, y_m] \mu \text{Cov}[y_p, g_j] = \frac{2c_k \cdot 2c_l}{V_A^2} \text{Cov}[g_i, g_j], \end{aligned}$$

the last step being (13) again, at the level of genetic values. This is (76) with no IBD term, as **[Outbred-mating-chain]** requires across a co-path.

Genotype against genetic value. Weighting (76) by β_l and summing over l gives $\text{Cov}[x_{ik}, g_j]$. The residual term vanishes identically, because by (69)

$$\sum_l \beta_l \left[(1 - 2a_{kk}) \delta_{kl} - (1 - \rho_g) \frac{4c_k c_l}{V_A} \right] = \beta_k (1 - 2a_{kk}) - 2(1 - \rho_g) c_k = 0,$$

and the first term gives the first half of (77), using $\sum_l 2\beta_l c_l = V_A$. The residual is uncorrelated with every genetic value, not only its carrier's. For the phenotype, $\text{Cov}[x_{ik}, e_j]$ is non-zero only through co-paths leaving j , since e_j has no other connections in the diagram; each such chain reaches x_{ik} through a covariance with some phenotype, which by the argument above is proportional to $2c_k$, so the constant of proportionality is again $\text{Cov}[g_i, e_j]/V_A$. Adding the two gives the second half of (77). The co-path step uses (77) for pairs one move shorter, so the three steps together are an induction on the number of moves.

Contracting (76) with $\beta_k \beta_l$ returns $\text{Cov}[g_i, g_j \mid \mathcal{P}]$, as it must: the residual term drops out by the identity above.

2.4.3 Allelic covariance between relatives

No separate result is needed at the allele level. By (15), an allele is half its transmitting parent's genotype plus a segregation deviation that is uncorrelated with everyone other than its carrier and the carrier's descendants. Write $u(i)$ for the parent i received allele u from — $m(i)$ for $u = m$ and $p(i)$ for $u = p$. Then for relatives i and j neither of whom

is descended from the other,

$$\boxed{\begin{aligned} \text{Cov}[z_{ik}^{(u)}, z_{jl}^{(v)}] &= \frac{1}{4} \text{Cov}[x_{u(i)k}, x_{v(j)l}] \\ \text{Cov}[z_{ik}^{(u)}, y_j] &= \frac{c_k}{V_A} \text{Cov}[g_{u(i)}, y_j] \end{aligned}} \quad (78)$$

where the right-hand sides are (76) and (77) evaluated for the transmitting parents. The second is the relatives' version of (21). When j is descended from i , only j 's allele is replaced, $\text{Cov}[z_{ik}^{(u)}, z_{jl}^{(v)}] = \frac{1}{2} \text{Cov}[z_{ik}^{(u)}, x_{v(j)l}]$, and the step is repeated down the lineal chain until it reaches i itself, where (67) applies.

For example, the two paternal alleles of maternal half-siblings come from fathers at mating distance 2, who share no IBD and have $\text{Cov}[g, g] = \rho_g \rho_y V_A$ by (59). So (78) and (76) give $\text{Cov}[z_{ik}^{(p)}, z_{jl}^{(p)}] = \rho_y a_{kl}$, whereas every other pairing of their alleles has covariance a_{kl} (plus an IBD term at $k = l$ for the two maternal alleles). The genotype covariance averages the four.

2.4.4 Examples

For $k \neq l$, dividing (76) by the within-individual value $4a_{kl} = 4\rho_g c_k c_l / V_A$ gives the fraction of an individual's own cross-variant disequilibrium that a relative shares with them:

$$\frac{\text{Cov}[x_{ik}, x_{jl} \mid \mathcal{P}]}{4a_{kl}} = \frac{\text{Cov}[g_i, g_j \mid \mathcal{P}] / V_A - (1 - \rho_g) \mathbb{E}[\pi_{ij} \mid \mathcal{P}]}{\rho_g} \quad (k \neq l). \quad (79)$$

It is the same for every pair of distinct variants. Table 7 evaluates it for the relationship classes of Table 6.

relationship	$\mathbb{E}[\pi_{ij} \mid \mathcal{P}]$	$\text{Cov}[g_i, g_j \mid \mathcal{P}] / V_A$	$\text{Cov}[x_{ik}, x_{jl} \mid \mathcal{P}] / 4a_{kl}, k \neq l$
self	1	1	1
mates	0	ρ_g	1
parent–offspring, full siblings	$\frac{1}{2}$	$\frac{1+\rho_g}{2}$	1
grandparent–grandchild, avuncular	$\frac{1}{4}$	$\left(\frac{1+\rho_g}{2}\right)^2$	$\frac{3+\rho_g}{4}$
first cousins	$\frac{1}{8}$	$\left(\frac{1+\rho_g}{2}\right)^3$	$\frac{4+3\rho_g+\rho_g^2}{8}$
half siblings	$\frac{1}{4}$	$\frac{1+2\rho_g+\rho_g\rho_y}{4}$	$\frac{3+\rho_y}{4}$
step siblings	0	$\rho_g \left(\frac{1+\rho_y}{2}\right)^2$	$\left(\frac{1+\rho_y}{2}\right)^2$
mating distance $N_{\mathcal{M}} \geq 1$	0	$\rho_g \rho_y^{N_{\mathcal{M}}-1}$	$\rho_y^{N_{\mathcal{M}}-1}$
lineal distance $N_{\mathcal{L}}$	$2^{-N_{\mathcal{L}}}$	$\left(\frac{1+\rho_g}{2}\right)^{N_{\mathcal{L}}}$	$\frac{(1+\rho_g)^{N_{\mathcal{L}}} - (1-\rho_g)}{2^{N_{\mathcal{L}}} \rho_g}$

Table 7: The two inputs of (76) and the cross-variant ratio (79) they imply, for common relationship classes. First-degree relatives and mates share *all* of an individual's own cross-variant disequilibrium, which is the equilibrium version of the observation under Table 3 that the cross-variant row is the same for self, siblings and parent–offspring. The last column stays finite as $\rho_g \rightarrow 0$ — the lineal entry tends to $(N_{\mathcal{L}} + 1)/2^{N_{\mathcal{L}}}$ — while the a_{kl} it multiplies vanishes.

2.4.5 Measures of relatedness

The genetic covariance is a sum over pairs of variants,

$$\text{Cov}[g_i, g_j \mid \mathcal{P}] = \underbrace{\sum_k \beta_k^2 \text{Cov}[x_{ik}, x_{jk} \mid \mathcal{P}]}_{\text{same-variant}} + \underbrace{\sum_{k \neq l} \beta_k \beta_l \text{Cov}[x_{ik}, x_{jl} \mid \mathcal{P}]}_{\text{cross-variant}},$$

and (76) evaluates both. Using (69) for the IBD term,

$$\begin{aligned} \text{same-variant} &= (1 - \rho_g) V_A \mathbb{E}[\pi_{ij} \mid \mathcal{P}] + \frac{4 \sum_k \beta_k^2 c_k^2}{V_A^2} \Delta_{ij}, \\ \text{cross-variant} &= \left(1 - \frac{4 \sum_k \beta_k^2 c_k^2}{V_A^2}\right) \Delta_{ij}, \\ \Delta_{ij} &:= \text{Cov}[g_i, g_j \mid \mathcal{P}] - (1 - \rho_g) V_A \mathbb{E}[\pi_{ij} \mid \mathcal{P}]. \end{aligned} \tag{80}$$

Δ_{ij} is the one new symbol in this section; it is $\rho_g V_A$ times the last column of Table 7. Could be written out instead if it only appears here. Under **[Uniform-inflation]** and **[VA-inflation]**, $c_k \approx \beta_k / (2(1 - \rho_g))$ and $(1 - \rho_g) V_A = V_A^{(0)}$, so the weight is

$$\frac{4 \sum_k \beta_k^2 c_k^2}{V_A^2} \approx \frac{\sum_k \beta_k^4}{(V_A^{(0)})^2} = \frac{1}{M_e}$$

by (41). So the same-variant part is $V_A^{(0)} \mathbb{E}[\pi_{ij} \mid \mathcal{P}]$ — the random-mating answer, with the random-mating additive variance — and everything assortment adds, Δ_{ij} , sits in the cross-variant part, apart from a share $1/M_e$ that **[Large-Me]** makes negligible.

The allelic sharing (6) is an unweighted average of the same-variant covariances, so by (76) its expectation given the path is

$$\begin{aligned} \mathbb{E}[A_{ij} \mid \mathcal{P}] &= \mathbb{E}[\pi_{ij} \mid \mathcal{P}] \frac{1}{M} \sum_k (1 - 2a_{kk}) + \left(\frac{1}{M} \sum_k \frac{4c_k^2}{V_A^2} \right) \Delta_{ij} \\ &\approx \mathbb{E}[\pi_{ij} \mid \mathcal{P}], \end{aligned} \tag{81}$$

where the approximation drops terms of order $\rho_g / ((1 - \rho_g)M)$. Under **[Uniform-inflation]**, $\frac{1}{M} \sum_k a_{kk} \approx \rho_g / (4(1 - \rho_g)M)$ and $\frac{1}{M} \sum_k 4c_k^2 / V_A^2 \approx 1 / (M V_A^{(0)})$. Neither A_{ij} nor π_{ij} therefore carries any information about Δ_{ij} . Mates make this plain: they share no IBD and $\mathbb{E}[A_{ij} \mid \mathcal{P}] \approx 0$, yet their genetic covariance is $\rho_g V_A$, all of it cross-variant. This is why the path cannot be dropped once A_{ij} or π_{ij} is known: assortment's contribution is carried by cross-variant covariance, which only the path determines.

It is also where the two halves of the random-effects model part ways. (76) makes the cross-variant covariance of genotypes proportional to $c_k c_l$, and so to $\beta_k \beta_l$, so the genotypes of a pair carry information about the sign of $\beta_k \beta_l$ and **[Uncorrelated-effects]** fails. **[Exchangeable-effects]** fails only through a_{kk} , which is of relative order $\beta_k^2 / V_A^{(0)}$ and so small by **[Large-Me]**.

2.4.6 Conditioning on realized IBD

We now condition on the meioses along the path — the transmission indicators B of (14) for every transmission between the pair and their common ancestors — which fixes the pair's IBD state $\text{IBD}_{ij}(k)$ at every variant.

Full siblings. Write o_1, o_2 for the siblings and m, p for their parents. By (54) each sibling is $\frac{1}{2}g_m + \frac{1}{2}g_p + g_o^\perp$, and the indicators are independent of the parents' genotypes, so conditioning on them leaves the parental part alone. The parental part is also uncorrelated with every segregation deviation given the indicators, since $\text{Cov}[g_m, z_{mk}^{(m)} - z_{mk}^{(p)}] = c_k - c_k = 0$. What remains is the covariance of the two segregation deviations given the indicators. By (16),

$$g_o^\perp = \sum_k \beta_k \left[(B_{ok}^{(m)} - \frac{1}{2})(z_{mk}^{(m)} - z_{mk}^{(p)}) + (B_{ok}^{(p)} - \frac{1}{2})(z_{pk}^{(m)} - z_{pk}^{(p)}) \right].$$

Given the indicators, every product $(B_{o_1k}^{(u)} - \frac{1}{2})(B_{o_2l}^{(v)} - \frac{1}{2})$ is fixed at $\pm \frac{1}{4}$, and:

- for $k \neq l$, the two allele differences are uncorrelated, since $\text{Cov}[z_k^{(m)} - z_k^{(p)}, z_l^{(m)} - z_l^{(p)}] = 2(s_{kl} - a_{kl}) = 0$ by (51);
- across the two parents they are uncorrelated too, because a co-path gives both of a parent's alleles the same covariance with the other parent;
- for $k = l$ and the same parent, $\text{Var}[z_k^{(m)} - z_k^{(p)}] = 1 - 2a_{kk}$, and the product of indicators is $+\frac{1}{4}$ when the siblings received the same allele from that parent — i.e. share it IBD — and $-\frac{1}{4}$ when not.

Summing over both parents,

$$\mathbb{E}[g_{o_1}g_{o_2} \mid B] = \frac{1}{2}V_A(1 + \rho_g) + \sum_k \beta_k^2(1 - 2a_{kk}) \frac{1}{2}(\text{IBD}_{o_1o_2}(k) - 1). \quad (82)$$

Other relationships. The same holds, with $\frac{1}{2}$ replaced by $\mathbb{E}[\pi_{ij} \mid \mathcal{P}]$, for every class in Table 6 except half cousins and step cousins:

$$\boxed{\mathbb{E}[g_i g_j \mid B, \mathcal{P}] = \text{Cov}[g_i, g_j \mid \mathcal{P}] + \sum_k \beta_k^2(1 - 2a_{kk}) \left(\frac{1}{2} \text{IBD}_{ij}(k) - \mathbb{E}[\pi_{ij} \mid \mathcal{P}] \right)} \quad (83)$$

It fails when a proband's allele can descend from either of two grandparents who stand in different relations to the other proband. Then the meioses decide which grandparent an allele came from, which changes its cross-variant covariance with the other proband by (78), and that change is not a function of IBD. In half cousins, an allele of one cousin descends from the shared grandparent or from that grandparent's mate, while an allele of the other descends from the shared grandparent or from the shared grandparent's *other* mate, and those two mates are at mating distance 2 rather than 1. **The same argument predicts a failure for great-grandparent–great-grandchild, whose descending allele can come through an ancestor's mate, who is off the path. Not in the verification pedigree yet. The general correction term is not derived.**

From meioses to π_{ij} . By (69), the weights in (83) are $\beta_k^2(1-2a_{kk}) = (1-\rho_g) \cdot 2\beta_k c_k$, and $2\beta_k c_k/V_A$ is variant k 's share of V_A by (22). So the sum is $(1-\rho_g)V_A$ times the difference between the *variance-weighted* fraction of the genome shared IBD and its expectation.

[IBD-share] The variance-weighted IBD fraction is the IBD fraction

Realized IBD is unrelated to how V_A is spread over the variants, so the variance-weighted fraction of the genome that a pair shares IBD equals the unweighted one:

$$\sum_k \frac{2\beta_k c_k}{V_A} \frac{1}{2} \text{IBD}_{ij}(k) \approx \pi_{ij}.$$

Justification for approximation: IBD states are set by meioses, which are independent of the effects, so the weighted and unweighted averages have the same expectation. They differ by sampling noise, which is small when V_A is spread over many variants ([Large-Me]) that are not clustered in a few IBD segments.

With it, (83) depends on the meioses only through π_{ij} :

$$\mathbb{E}[g_i g_j \mid \pi_{ij}, \mathcal{P}] \approx \text{Cov}[g_i, g_j \mid \mathcal{P}] + (1 - \rho_g) V_A \left(\pi_{ij} - \mathbb{E}[\pi_{ij} \mid \mathcal{P}] \right) \quad (84)$$

The slope is $(1 - \rho_g)V_A = 2\text{Var}[g_o^\perp]$ exactly, by (55), and $\approx V_A^{(0)}$ under [VA-inflation]. The departure of a pair's realized IBD from its expectation is produced by segregation, and segregation variance is the one variance component that assortative mating does not inflate. So realized IBD moves the genetic covariance at the rate set by the *random-mating* additive variance, while the path supplies the level, including all of assortment's cross-variant contribution.

Phenotypes. Reading a proband at the phenotype adds its environment, which is correlated with its mates' alleles. For a proband with no mate on the path — the youngest individual of a lineal chain, as for siblings and cousins — the environment is uncorrelated with every meiosis, and (83) holds for phenotypes with $\text{Cov}[y_i, y_j \mid \mathcal{P}]$ in place of $\text{Cov}[g_i, g_j \mid \mathcal{P}]$. When the proband read at the phenotype is an *ancestor* of the other, their environment is correlated with the alleles of their mate, which the descendant inherits exactly where it does not inherit the ancestor's own; the slope then becomes $(1 - \rho_y)V_A$ in place of $(1 - \rho_g)V_A$:

$$\mathbb{E}[y_i y_j \mid \pi_{ij}, \mathcal{P}] \approx \text{Cov}[y_i, y_j \mid \mathcal{P}] + (1 - \rho_*) V_A \left(\pi_{ij} - \mathbb{E}[\pi_{ij} \mid \mathcal{P}] \right), \quad (85)$$

where $\rho_* = \rho_y$ when a proband is the source of a lineal chain, and $\rho_* = \rho_g$ otherwise. This is the same substitution of ρ_y for ρ_g in the proband's own edges that produced (65).

Linkage. Under [Unlinked], the IBD states of different variants are independent, so π_{ij} concentrates at $\mathbb{E}[\pi_{ij} \mid \mathcal{P}]$ with variance of order $1/M$, and (84) has almost nothing to condition on. (83) is exact for any realized meioses, but the realistic variation in π_{ij} comes from linkage, which also changes the equilibrium disequilibrium between linked variants ($s_{kl} \neq a_{kl}$). Applying these results to real genomes needs either an approximation that linked pairs are a negligible share of all pairs of variants, or a derivation with linkage.

2.4.7 Conditioning on allelic sharing

Not yet derived. Conditioning on A_{ij} needs $\text{Cov}[g_i g_j, A_{ij} \mid \mathcal{P}]$, a fourth moment of the genotypes, which (76) alone does not give. The analogue of (84) does NOT follow by replacing π_{ij} with A_{ij} : unlike the meioses, A_{ij} is a function of the genotypes, and through the within-individual disequilibrium it is correlated with the cross-variant part as well. A preliminary calculation under a Gaussian approximation to the genotypes suggests that, for unrelated pairs, the regression slope of $g_i g_j$ on A_{ij} is $\sum_k (2c_k)^2 \approx V_A^{(0)} / (1 - \rho_g)^2$ rather than $V_A^{(0)}$. Unverified.

To AI agents: you may edit below this

References

- [1] Sewall Wright. Correlation and causation. *Journal of Agricultural Research*, 20: 557–585, 1921.
- [2] Sewall Wright. The method of path coefficients. *The Annals of Mathematical Statistics*, 5(3):161–215, 1934. URL <http://www.jstor.org/stable/2957502>.
- [3] Sewall Wright. Systems of mating. I. the biometric relations between parent and offspring. *Genetics*, 6(2):111–123, 1921. doi: 10.1093/genetics/6.2.111.
- [4] Sewall Wright. Systems of mating. III. assortative mating based on somatic resemblance. *Genetics*, 6(2):144–161, 1921. doi: 10.1093/genetics/6.2.144.
- [5] R A Fisher. The correlation between relatives on the supposition of mendelian inheritance. *Transactions of the Royal Society of Edinburgh*, 52:399–433, 1918. doi: 10.1017/s0080456800012163.
- [6] Hans Fredrik Sunde, Espen Moen Eilertsen, and Fartein Ask Torvik. Understanding indirect assortative mating and its intergenerational consequences for educational attainment. *Nat. Commun.*, 16(1):5264, 6 June 2025.
- [7] Loic Yengo, Matthew R Robinson, Matthew C Keller, Kathryn E Kemper, Yuanhao Yang, Maciej Trzaskowski, Jacob Gratten, Patrick Turley, David Cesarini, Daniel J Benjamin, Naomi R Wray, Michael E Goddard, Jian Yang, and Peter M Visscher. Imprint of assortative mating on the human genome. *Nature Human Behaviour*, 2: 948–954, 2018. doi: 10.1038/s41562-018-0476-3.
- [8] Hans Fredrik Sunde, Nikolai Haahjem Eftedal, Rosa Cheesman, Elizabeth C Corfield, Thomas H Kleppesø, Anne Caroline Seierstad, Eivind Ystrom, Espen Moen Eilertsen, and Fartein Ask Torvik. Genetic similarity between relatives provides evidence on the presence and history of assortative mating. *Nat. Commun.*, 15(1):2641, 2024. doi: 10.1038/s41467-024-46939-9.
- [9] Luke J. O’Connor, Armin P. Schoech, Farhad Hormozdiari, Steven Gazal, Nick Patterson, and Alkes L. Price. Extreme polygenicity of complex traits is explained by negative selection. *American Journal of Human Genetics*, 105(3):456–476, 2019. doi: 10.1016/j.ajhg.2019.07.003.

- [10] Thomas Nagylaki. The correlation between relatives with assortative mating. *Annals of Human Genetics*, 42(2):131–137, 1978. doi: 10.1111/j.1469-1809.1978.tb00938.x.