= Eclampsia

Assessment step 1:

=

= Cell your helps

✓ Assessment

- Airway ~~[illegible]~~
- Breakin

Circulation

✓ ~~IV canol [illegible]~~

Assessment

BP: 160/110 ·

RR: 20 ✓

Pulse 80

Reflexes 1+ ✓

IV line open

urine protein 2+

urine catheter 40

FHR 144

~~maintain dose~~ maintenance dose ①

· 3 hour ago

MgSO4 - 4 g IV and 10 g IM [5g per buttock]

assessment

✓ RR

✓ Reflexes

✓ urine catheter (output and input)

BP 158/106mmHg = Emergency

= ~~[illegible]~~ Nifedpine 10mg oral = Lebetalol iv

Check BP every 10 - 20mint

= check urine catheter

✓ out put in put

every 1hour = 30ML

= maintenan dose - P

8 ~~calle~~ Give 5g IM every 4 hour alternatatin

- Since The loading doso ~~gi~~ was give 3 hour ago, next maintenane dose 1 hour safety are

maintenan dose check

✓ = Before very maintenan dose check

✓ RR 16m/h

✓ reflexes

✓ urine ~~out~~ out put ~~30~~ Least 30ML hour

IV fluid

= continue Ringers. carefully controlled

Avoid IV fluid becouces othe risk edema

calcium gluconat 10% keep availabe as an ~~at~~ antidote

magnesium toxicity