case study

Vital smgns
- BP 188/ 106mmHg
Urine proteauria 2+
RR 20 ~~lungs~~
Pulse 80 reflect +1
If cateter placed 140ml urine collect
FHR 144
GA = 32 weeks
NO smgns of labour

diagnosis
eclampsia with sere hypertension

Management step by step
1= call for student nursing.
~~2~~ = call the emmergency team
= do not leave the patient alone
= prepare emmergency equipment
step(2) check Airway and breathing.
= patient place Inthe left leteral position
= maintain clear Airway
= give oxygen if ~~needed~~
= patient protect fram injury
= never not anything Inter mouth
Step 3 =
✓ Check Vital smgns
✓ Such as BP PR, RR and oxygen
step(4)
✓ control convulsion
✓ give magnesium supplate
step(5) control Blood pressure
- give labetatol or Hydalasine ~~as~~

Step (6)

- ✓ monitor mother and baby
- ✓ monitor urine output
- ✓ monitor FHR

Step (7)

- ✓ ~~pay~~ prepare delivary.
- ✓ First mother than Plan delivary.

Step (8)

- ✓ continue magnesium ~~for~~ 24hours after delivary
- ✓ loading 4g 10+ 10gml (5g in each ~~buttonia~~
- ✓ Medication:
- ✓ magnesium sulphate
- (2) maintenance 5gml every 4hours.

~~analog~~

- ✓ ~~p~~ porpose: control and prevent convulsion