


# VECTOR BOSON FUSION AND VECTOR BOSON SCATTERING

Dieter Zeppenfeld  
Karlsruhe Institute of Technology




Bundesministerium  
für Bildung  
und Forschung

- Higgs coupling measurement and VBF
- Effective Lagrangians and anomalous couplings
- VVVV vertices and VBS

# Tensor structure of the $HVV$ coupling

Most general  $HVV$  vertex  $T^{\mu\nu}(q_1, q_2)$



Physical interpretation of terms:

**SM Higgs**  $\mathcal{L}_I \sim HV_\mu V^\mu \longrightarrow a_1$

loop induced couplings for neutral scalar

**CP even**  $\mathcal{L}_{eff} \sim HV_{\mu\nu} V^{\mu\nu} \longrightarrow a_2$

**CP odd**  $\mathcal{L}_{eff} \sim HV_{\mu\nu} \tilde{V}^{\mu\nu} \longrightarrow a_3$

Must distinguish  $a_1, a_2, a_3$  experimentally

$$T^{\mu\nu} = a_1 g^{\mu\nu} + a_2 (q_1 \cdot q_2 g^{\mu\nu} - q_1^\nu q_2^\mu) + a_3 \varepsilon^{\mu\nu\rho\sigma} q_{1\rho} q_{2\sigma}$$

The  $a_i = a_i(q_1, q_2)$  are scalar form factors

## Connection to effective Lagrangian

We need model of the underlying UV physics to determine the form factors  $a_i(q_1, q_2)$

Approximate its low-energy effects by an effective Lagrangian

$$\mathcal{L}_{\text{eff}} = \frac{f_{WW}}{\Lambda^2} \phi^\dagger \hat{W}_{\mu\nu} \hat{W}^{\mu\nu} \phi + \frac{f_\phi}{\Lambda^2} \left( \phi^\dagger \phi - \frac{v^2}{2} \right) (D_\mu \phi)^\dagger D^\mu \phi + \dots + \sum_i \frac{f_i^{(8)}}{\Lambda^4} \mathcal{O}_i^{(8)} + \dots$$

Gives leading terms for form factors, e.g. for hWW coupling

$$a_1 = \frac{2m_W^2}{v} \left( 1 + \frac{f_\phi}{\Lambda^2} \frac{v^2}{2} \right) + \sum_i c_i^{(1)} \frac{f_i^{(8)}}{\Lambda^4} v^2 q^2 + \dots$$

$$a_2 = c^{(2)} \frac{f_{WW}}{\Lambda^2} v + \sum_i c_i^{(2)} \frac{f_i^{(8)}}{\Lambda^4} v q^2 + \dots$$

$$a_3 = c^{(3)} \frac{\tilde{f}_{WW}}{\Lambda^2} v + \sum_i c_i^{(3)} \frac{\tilde{f}_i^{(8)}}{\Lambda^4} v q^2 + \dots$$

Describe same physics (for a particular vertex) by taking some minimal set of effective Lagrangian coefficients  $f_i$  as form factors

## Implementation in VBFNLO

Start from effective Lagrangians (set PARAMETR1=.true. in anom\_HVV.dat )

$$\begin{aligned} \mathcal{L} = & \frac{g_{5e}^{HZZ}}{2\Lambda_5} H Z_{\mu\nu} Z^{\mu\nu} + \frac{g_{5o}^{HZZ}}{2\Lambda_5} H \tilde{Z}_{\mu\nu} Z^{\mu\nu} + \frac{g_{5e}^{HWW}}{\Lambda_5} H W_{\mu\nu}^+ W_-^{\mu\nu} + \frac{g_{5o}^{HWW}}{\Lambda_5} H \tilde{W}_{\mu\nu}^+ W_-^{\mu\nu} + \\ & \frac{g_{5e}^{HZ\gamma}}{\Lambda_5} H Z_{\mu\nu} A^{\mu\nu} + \frac{g_{5o}^{HZ\gamma}}{\Lambda_5} H \tilde{Z}_{\mu\nu} A^{\mu\nu} + \frac{g_{5e}^{H\gamma\gamma}}{2\Lambda_5} H A_{\mu\nu} A^{\mu\nu} + \frac{g_{5o}^{H\gamma\gamma}}{2\Lambda_5} H \tilde{A}_{\mu\nu} A^{\mu\nu} \end{aligned}$$

or , alternatively, (set PARAMETR3=.true. in anom\_HVV.dat )

$$\mathcal{L}_{\text{eff}} = \frac{f_{WW}}{\Lambda_6^2} \phi^\dagger \hat{W}_{\mu\nu} \hat{W}^{\mu\nu} \phi + \frac{f_{BB}}{\Lambda_6^2} \phi^\dagger \hat{B}_{\mu\nu} \hat{B}^{\mu\nu} \phi + \text{CP-odd part} + \dots$$

see VBFNLO manual for details on how to set the anomalous coupling choices


Remember to choose form factors in anom\_HVV.dat

$$F_1 = \frac{M^2}{q_1^2 - M^2} \frac{M^2}{q_2^2 - M^2} \quad \text{or} \quad F_2 = -2 M^2 C_0(q_1^2, q_2^2, (q_1 + q_2)^2, M^2)$$

## Jet transverse momentum

Form factors affect momentum transfer and thus jet transverse momenta (Here:  $a_2$  only)


Figy, DZ hep-ph/0403297



- Change in tagging jet  $p_T$  distributions is sensitive indicator of anomalous couplings
- Can choose form-factor such as to approximate SM  $p_T$  distributions of the two tagging jets

## Azimuthal angle correlations

Tell-tale signal for non-SM coupling is azimuthal angle between tagging jets



Dip structure at  $90^\circ$  (CP even) or  $0/180^\circ$  (CP odd) only depends on tensor structure of  $hVV$  vertex. Very little dependence on form factor, LO vs. NLO, Higgs mass etc.

Same physics in decay plane correlations for  $h \rightarrow ZZ^* \rightarrow 4 \text{ leptons}$

## Vector boson scattering

The  $m_h = 126$  GeV Higgs will unitarize  $VV \rightarrow VV$  scattering **provided** it has SM  $hVV$  couplings

⇒ Check this by either

- precise measurements of the  $hVV$  couplings at the light Higgs resonance
- measurement of  $VV \rightarrow VV$  differential cross sections at high  $p_T$  and invariant mass

Full  $qq \rightarrow qqVV$  with  $VV$  leptonic and semileptonic decay is implemented in VBFNLO with NLO QCD corrections and large set of dimension 6 and 8 terms in the effective Lagrangian

## Going beyond dimension 6

Reason for dimension 8 operators like

$$\begin{aligned}\mathcal{L}_{S,0} &= \left[ (D_\mu \Phi)^\dagger D_\nu \Phi \right] \times \left[ (D^\mu \Phi)^\dagger D^\nu \Phi \right] \\ \mathcal{L}_{M,1} &= \text{Tr} \left[ \hat{W}_{\mu\nu} \hat{W}^{\nu\beta} \right] \times \left[ (D_\beta \Phi)^\dagger D^\mu \Phi \right] \\ \mathcal{L}_{T,1} &= \text{Tr} \left[ \hat{W}_{\alpha\nu} \hat{W}^{\mu\beta} \right] \times \text{Tr} \left[ \hat{W}_{\mu\beta} \hat{W}^{\alpha\nu} \right]\end{aligned}$$

- Dimension 6 operators only do not allow to parameterize  $VVVV$  vertex with arbitrary helicities of the four gauge bosons

For example:  $\mathcal{L}_{S,0}$  is needed to describe  $V_L V_L \rightarrow V_L V_L$  scattering


- New physics may appear at 1-loop level for dimension 6 operators but at tree level for some dimension 8 operators



## $VV \rightarrow W^+ W^-$ with dimension 8 operators

Effect of  $\mathcal{L}_{eff} = \frac{f_{M,1}}{\Lambda^4} \text{Tr} [\hat{W}_{\alpha\gamma} \hat{W}^{\mu\beta}] \times \text{Tr} [\hat{W}_{\mu\beta} \hat{W}^{\alpha\gamma}]$


with  $T_1 = \frac{f_{M,1}}{\Lambda^4}$  constant on  $pp \rightarrow W^+ W^- jj \rightarrow e^+ \nu_e \mu^- \bar{\nu}_\mu jj$



- Small increase in cross section at high  $WW$  invariant mass??

## $VV \rightarrow W^+ W^-$ with dimension 8 operators

Effect of constant  $T_1 = \frac{f_{M,1}}{\Lambda^4}$  on  $pp \rightarrow W^+ W^- jj \rightarrow e^+ \nu_e \mu^- \bar{\nu}_\mu jj$



- Huge increase in cross section at high  $m_{WW}$  is completely unphysical
- Need form factor for analysis or some other unitarization procedure

## Conclusions

- VBF production of a light Higgs provides for important input to the coupling measurements
- VBS at high  $VV$  invariant mass and high  $p_T$  of the weak bosons complements these measurements
- Model independent parameterizations of deviations from the SM are provided by a variety of programs
- Form factors or some other unitarization procedure cannot be avoided when using effective Lagrangians for  $VV$  scattering at the LHC (for quite some time)