

Fast Matrix Exponentiation

With Applications

Nurseiit Abdimomyn

UNIST

October 26, 2018

1 Introduction

2 Matrix exponentiation

3 Applications

Matrix Multiplication

Consider two matrices:

Matrix Multiplication

Consider two matrices:

- Matrix A is $n * k$ dimensional.

Matrix Multiplication

Consider two matrices:

- Matrix A is $n * k$ dimensional.
- Matrix B is $k * m$ dimensional.

Matrix Multiplication

Consider two matrices:

- Matrix A is $n * k$ dimensional.
- Matrix B is $k * m$ dimensional.

Notice that A 's columns and B 's rows number are identical!

Matrix Multiplication

Consider two matrices:

- Matrix A is $n * k$ dimensional.
- Matrix B is $k * m$ dimensional.

Notice that A 's columns and B 's rows number are identical!

Then we define matrix $C = A * B$ as:

$$\begin{bmatrix} c_{11} & c_{12} & \dots & c_{1m} \\ c_{21} & c_{22} & \dots & c_{2m} \\ \dots & \dots & \dots & \dots \\ c_{n1} & c_{n2} & \dots & c_{nm} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1k} \\ a_{21} & a_{22} & \dots & a_{2k} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nk} \end{bmatrix} * \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2m} \\ \dots & \dots & \dots & \dots \\ b_{k1} & b_{k2} & \dots & b_{km} \end{bmatrix}$$

Matrix Multiplication

Consider two matrices:

- Matrix A is $n * k$ dimensional.
- Matrix B is $k * m$ dimensional.

Notice that A 's columns and B 's rows number are identical!

Then we define matrix $C = A * B$ as:

$$\begin{bmatrix} c_{11} & c_{12} & \dots & c_{1m} \\ c_{21} & c_{22} & \dots & c_{2m} \\ \dots & \dots & \dots & \dots \\ c_{n1} & c_{n2} & \dots & c_{nm} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1k} \\ a_{21} & a_{22} & \dots & a_{2k} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nk} \end{bmatrix} * \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2m} \\ \dots & \dots & \dots & \dots \\ b_{k1} & b_{k2} & \dots & b_{km} \end{bmatrix}$$

Thus, C is an $n * m$ dimensional matrix.

Which is calculated as: $c_{ij} = \sum_{r=1}^k a_{ir} * b_{rj}$

Couple of things to notice

- Matrix C has $n * m$ elements and each element is computed in k steps with given formula.

Couple of things to notice

- Matrix C has $n * m$ elements and each element is computed in k steps with given formula.
Thus, we can obtain C in $O(n * m * k)$, given A and B .

Couple of things to notice

- Matrix C has $n * m$ elements and each element is computed in k steps with given formula.
Thus, we can obtain C in $O(n * m * k)$, given A and B .
- If $n = m = k$ (i.e. both A and B have n rows and n columns), then C has n rows and n columns, and can be computed in $O(n^3)$.

Some useful properties of matrix multiplication

- It is *not commutative*: $A * B \neq B * A$ in general case.

Some useful properties of matrix multiplication

- It is *not commutative*: $A * B \neq B * A$ in general case.
- It is *associative*:
 $A * B * C = (A * B) * C = A * (B * C)$ in case $A * B * C$ exists;

Some useful properties of matrix multiplication

- It is *not commutative*: $A * B \neq B * A$ in general case.
- It is *associative*:
 $A * B * C = (A * B) * C = A * (B * C)$ in case $A * B * C$ exists;
- If you have a matrix with n rows and n columns, then multiplying it by I_n gives the same matrix.
i. e. $I_n * A = A * I_n = A$.

Some useful properties of matrix multiplication

- It is *not commutative*: $A * B \neq B * A$ in general case.
- It is *associative*:
 $A * B * C = (A * B) * C = A * (B * C)$ in case $A * B * C$ exists;
- If you have a matrix with n rows and n columns, then multiplying it by I_n gives the same matrix.
 - i. e. $I_n * A = A * I_n = A$.Where I_n is a matrix with n rows and n columns of such form:

$$I_n = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

1 Introduction

2 Matrix exponentiation

3 Applications

Matrix exponentiation

Suppose we need to find some $n * n$ dimensional square matrix A to the power p or A^p

Matrix exponentiation

Suppose we need to find some $n * n$ dimensional square matrix A to the power p or A^p

We can do so via:

```
function matpow_naive(A, p):  
    result = I_n  
    for i = 1..p:  
        result = result * A  
    return result
```

Which will run in $O(n^3 * p)$

Fast Matrix exponentiation

Can we do it any faster?

Fast Matrix exponentiation

Can we do it any faster?

Yes, we can apply the *BinPower* algorithm here:

```
function matBinPow(A, p):  
    result = I_n  
    while p > 0:  
        if p % 2 == 1:  
            result = result * A  
        A = A * A  
        p = p / 2  
    return result
```

Which will run in $O(n^3 * \log p)$

1 Introduction

2 Matrix exponentiation

3 Applications

Finding Nth Fibonacci number

Fibonacci numbers, F_n are defined as:

- $F_0 = F_1 = 1$
- $F_i = F_{i-1} + F_{i-2}$ for $i > 1$.

Finding Nth Fibonacci number

Fibonacci numbers, F_n are defined as:

- $F_0 = F_1 = 1$
- $F_i = F_{i-1} + F_{i-2}$ for $i > 1$.

We want to calculate $F_n \bmod M$, where $n < 10^{18}$ and $M = 10^9 + 7$.

Finding Nth Fibonacci number

Suppose we have a vector (matrix with one row and several columns) of (F_{i-2}, F_{i-1}) and we want to multiply it by some matrix, so that we get (F_{i-1}, F_i) as a result.

Finding Nth Fibonacci number

Suppose we have a vector (matrix with one row and several columns) of (F_{i-2}, F_{i-1}) and we want to multiply it by some matrix, so that we get (F_{i-1}, F_i) as a result.

Let's call this matrix M :

$$(F_{i-2}, F_{i-1}) * M = (F_{i-1}, F_i)$$

Two questions we should answer arise immediately:

- 1 What are the dimensions of M ?
- 2 What are the exact values in M ?

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

- The *size*.

We multiply (F_{i-2}, F_{i-1}) , which has 1 row and 2 columns, by M .
The result is (F_{i-1}, F_i) , which has 1 row and 2 columns.

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

- The *size*.

We multiply (F_{i-2}, F_{i-1}) , which has 1 row and 2 columns, by M .
The result is (F_{i-1}, F_i) , which has 1 row and 2 columns.

By definition, if we multiply a matrix with N rows and K columns by a matrix with K rows and L columns, we get a matrix with N rows and L columns.

Therefore, matrix M has $K = 2$ rows and $L = 2$ columns.

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

- The *values*.

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

- The *values*.

We now know that M has 2 rows and 2 columns, 4 values overall.

Lets denote them by letters, as we usually do with unknown variables:

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Finding Nth Fibonacci number

We can answer them using the definition of matrix multiplication:

- The *values*.

We now know that M has 2 rows and 2 columns, 4 values overall.

Lets denote them by letters, as we usually do with unknown variables:

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

We want to find a, b, c and d .

Finding Nth Fibonacci number

We have,

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Finding Nth Fibonacci number

We have,

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

We want to find a, b, c and d .

Let's see what we get if we multiply (F_{i-2}, F_{i-1}) by M by definition:

Finding Nth Fibonacci number

We have,

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

We want to find a, b, c and d .

Let's see what we get if we multiply (F_{i-2}, F_{i-1}) by M by definition:

$$(F_{i-2}, F_{i-1}) * \begin{bmatrix} a & b \\ c & d \end{bmatrix} = (a * F_{i-2} + c * F_{i-1}, b * F_{i-2} + d * F_{i-1})$$

Finding Nth Fibonacci number

We have,

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

We want to find a, b, c and d .

Let's see what we get if we multiply (F_{i-2}, F_{i-1}) by M by definition:

$$(F_{i-2}, F_{i-1}) * \begin{bmatrix} a & b \\ c & d \end{bmatrix} = (a * F_{i-2} + c * F_{i-1}, b * F_{i-2} + d * F_{i-1})$$

Moreover, we know that the result of this multiplication must be (F_{i-1}, F_i) :
 $(a * F_{i-2} + c * F_{i-1}, b * F_{i-2} + d * F_{i-1}) = (F_{i-1}, F_i)$

Finding Nth Fibonacci number

Now we can write the system of equations:

- $a * F_{i-2} + c * F_{i-1} = F_{i-1}$
- $b * F_{i-2} + d * F_{i-1} = F_i$

Finding Nth Fibonacci number

Now we can write the system of equations:

- $a * F_{i-2} + c * F_{i-1} = F_{i-1}$
- $b * F_{i-2} + d * F_{i-1} = F_i$

The easiest solution is to set:

$$a = 0, b = 1, c = 1, d = 1$$

Finding Nth Fibonacci number

Now we can write the system of equations:

- $a * F_{i-2} + c * F_{i-1} = F_{i-1}$
- $b * F_{i-2} + d * F_{i-1} = F_i$

The easiest solution is to set:

$$a = 0, b = 1, c = 1, d = 1$$

Thus, we obtain:

$$(F_{i-2}, F_{i-1}) * \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = (F_{i-1}, F_i)$$

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?
Hang on, you'll see in a moment.

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?

Hang on, you'll see in a moment.

Initially, we have F_0 and F_1 .

$$(F_0, F_1) = (1, 1)$$

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?

Hang on, you'll see in a moment.

Initially, we have F_0 and F_1 .

$$(F_0, F_1) = (1, 1)$$

By multiplying this vector with the matrix M we get:

$$(F_1, F_2) = (1, 2) \text{ or } (1, 1) * M = (1, 2).$$

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?

Hang on, you'll see in a moment.

Initially, we have F_0 and F_1 .

$$(F_0, F_1) = (1, 1)$$

By multiplying this vector with the matrix M we get:

$$(F_1, F_2) = (1, 2) \text{ or } (1, 1) * M = (1, 2).$$

If we multiply $(1, 2)$ by M :

$$(1, 2) * M = (2, 3)$$

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?

Hang on, you'll see in a moment.

Initially, we have F_0 and F_1 .

$$(F_0, F_1) = (1, 1)$$

By multiplying this vector with the matrix M we get:

$$(F_1, F_2) = (1, 2) \text{ or } (1, 1) * M = (1, 2).$$

If we multiply $(1, 2)$ by M :

$$(1, 2) * M = (2, 3)$$

But we could get the same result by multiplying $(1, 1)$ by M^2 .

$$(1, 1) * M^2 = (2, 3).$$

Finding Nth Fibonacci number

You may wonder why on Earth would we need to overcomplicate the problem, introduce matrices or else?

Hang on, you'll see in a moment.

Initially, we have F_0 and F_1 .

$$(F_0, F_1) = (1, 1)$$

By multiplying this vector with the matrix M we get:

$$(F_1, F_2) = (1, 2) \text{ or } (1, 1) * M = (1, 2).$$

If we multiply $(1, 2)$ by M :

$$(1, 2) * M = (2, 3)$$

But we could get the same result by multiplying $(1, 1)$ by M^2 .

$$(1, 1) * M^2 = (2, 3).$$

Now, can you see the pattern?

Finding Nth Fibonacci number

In general, we get:

$$(1, 1) * M^k = (F_k, F_{k+1})$$

Finding Nth Fibonacci number

In general, we get:

$$(1, 1) * M^k = (F_k, F_{k+1})$$

Computing M^k takes $O(2^3 * \log(k))$ time.

Thus, we can now find N -th Fibonacci number in $O(\log(N))$ time.

General Solution for Recurrence Relations

Thank you!