AI Assignment 1

$Nurseiit\ Abdimomyn-20172001$

26/09/2021

1.1 We have $\theta = [\theta_0, \theta_1, \theta_2]^T$ and $x_i = [1, x_{i0}, x_{i1}]^T$ for $h_{\theta}(x) = \theta^T x$.

So we derive for the gradient:

$$\frac{\partial L(\theta)}{\partial \theta_0} = \frac{\partial}{\partial \theta_0} \sum_{i} (\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i)^2$$
$$= \sum_{i} \frac{\partial}{\partial \theta_0} (\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i)^2$$

$$= \sum_{i} \frac{\partial}{\partial \theta_0} (\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i)^2$$

$$= \sum_{i} 2(\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i)$$

Doing the same for θ_1 and θ_2 we get:

$$\frac{\partial L(\theta)}{\partial \theta_1} = \sum_{i} 2(\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i) * x_{i0}$$

$$\frac{\partial L(\theta)}{\partial \theta_2} = \sum_{i} 2(\theta_0 + \theta_1 x_{i0} + \theta_2 x_{i1} - y_i) * x_{i1}$$

To find a minimum by gradient descent we use:

 $\theta^{k+1} \leftarrow \theta^k - \eta_k \frac{\partial L(\theta)}{\partial \theta}$ where we have $\eta = 0.01$ for all k.

Thus,
$$\theta^{k+1} \leftarrow \theta^k - 0.01 * \frac{\partial L(\theta)}{\partial \theta}$$
.

Initial parameter $\theta^0 = [0, 0, 0]^T$, so compute for θ^1 simultaneously:

$$\theta_0^1 = \theta_0^0 - 0.01 * \sum_{i} 2(\theta_0^0 + \theta_1^0 x_{i0} + \theta_2^0 x_{i1} - y_i)$$

$$= 0 - 0.01 * \sum_{i} 2(0 + 0 * x_{i0} + 0 * x_{i1} - y_{i})$$

$$= 0.02 * \sum_{i} y_i = 0.02 * 37 = 0.74.$$

$$\theta_1^1 = \theta_1^0 - 0.01 * \sum_i 2(\theta_0^0 + \theta_1^0 x_{i0} + \theta_2^0 x_{i1} - y_i) * x_{i0}$$

$$= 0 - 0.01 * \sum_i 2(0 + 0 * x_{i0} + 0 * x_{i1} - y_i) * x_{i0}$$

$$= 0.02 * \sum_i (y_i * x_{i0})$$

$$= 0.02 * (6 * 1 + 4 * 2 + 7 * 3 + 10 * 3 + 10 * 4) = 0.02 * 105 = 2.1.$$

$$\begin{split} \theta_2^1 &= \theta_2^0 - 0.01 * \sum_i 2(\theta_0^0 + \theta_1^0 x_{i0} + \theta_2^0 x_{i1} - y_i) * x_{i1} \\ &= 0 - 0.01 * \sum_i 2(0 + 0 * x_{i0} + 0 * x_{i1} - y_i) * x_{i1} \\ &= 0.02 * \sum_i (y_i * x_{i1}) \\ &= 0.02 * (6 * 2 + 4 * 1 + 7 * 2 + 10 * 3 + 10 * 5) = 0.02 * 110 = 2.2. \\ \text{So for the first iteration we have } \theta^1 = [0.74, 2.1, 2.2]^T. \end{split}$$

Second iteration:

$$\theta_0^2 = \theta_0^1 - 0.01 * \sum_i 2(\theta_0^1 + \theta_1^1 x_{i0} + \theta_2^1 x_{i1} - y_i)$$

$$= 0.74 - 0.01 * \sum_i 2(0.74 + 2.1x_{i0} + 2.2x_{i1} - y_i)$$

$$= 0.74 - 0.02 * 22.6 = 0.288.$$

$$\theta_1^2 = \theta_1^1 - 0.01 * \sum_i 2(\theta_0^1 + \theta_1^1 x_{i0} + \theta_2^1 x_{i1} - y_i) * x_{i0}$$

$$= 2.1 - 0.01 * \sum_i 2(0.74 + 2.1 x_{i0} + 2.2 x_{i1} - y_i) * x_{i0}$$

$$= 2.1 - 0.02 * \sum_i (0.74 + 2.1 x_{i0} + 2.2 x_{i1} - y_i) * x_{i0} = 0.6536$$

$$\theta_2^2 = \theta_2^1 - 0.01 * \sum_i 2(\theta_0^1 + \theta_1^1 x_{i0} + \theta_2^1 x_{i1} - y_i) * x_{i1}$$
$$= \theta_2^1 - 0.02 * \sum_i (\theta_0^1 + \theta_1^1 x_{i0} + \theta_2^1 x_{i1} - y_i) * x_{i1}$$

$$= 2.2 - 0.02 * \sum_{i} (0.74 + 2.1x_{i0} + 2.2x_{i1} - y_i) * x_{i1} = 0.6776.$$

Thus, $\theta^2 = [0.288, 0.6536, 0.6776]^T$.

$$1.2 \ \theta^* = (X^T X)^{-1} X^T y$$

$$= \left(\begin{bmatrix} 2 & 1 & 2 & 3 & 5 \\ 1 & 2 & 3 & 3 & 4 \end{bmatrix} * \begin{bmatrix} 1 & 2 \\ 2 & 1 \\ 3 & 2 \\ 3 & 3 & 4 \end{bmatrix} \right)^{-1} * \begin{bmatrix} 2 & 1 & 2 & 3 & 5 \\ 1 & 2 & 3 & 3 & 4 \end{bmatrix} * \begin{bmatrix} 6 \\ 4 \\ 7 \\ 10 \\ 10 \end{bmatrix}$$

$$= \left(\begin{bmatrix} 39 & 43 \\ 39 & 39 \end{bmatrix} \right)^{-1} * \begin{bmatrix} 2 & 1 & 2 & 3 & 5 \\ 1 & 2 & 3 & 3 & 4 \end{bmatrix} * \begin{bmatrix} 6 \\ 4 \\ 7 \\ 10 \\ 10 \end{bmatrix}$$

$$= \begin{bmatrix} -0.25 & 0.27564 \\ 0.25 & -0.25 \end{bmatrix} * \begin{bmatrix} 2 & 1 & 2 & 3 & 5 \\ 1 & 2 & 3 & 3 & 4 \end{bmatrix} * \begin{bmatrix} 6 \\ 4 \\ 7 \\ 10 \\ 10 \end{bmatrix}$$

$$= \begin{bmatrix} -0.22436 & 0.30128 & 0.32692 & 0.07692 & -0.14744 \\ 0.25 & -0.25 & -0.25 & 0 & 0.25 \end{bmatrix} * \begin{bmatrix} 6 \\ 4 \\ 7 \\ 10 \\ 10 \end{bmatrix}$$

$$= \begin{bmatrix} 1.4422 \\ 1.25 \end{bmatrix}$$