

Table 1. Primer sequences used for amplification of TCF7L2 rs12255372

Primer	Sequence (5'→3')	Product size
Forward	CTGGAAACTAAGGCGTGAGG	346 bp
Reverse	GGGTCGATGTTGTTGAGCTT	346 bp

Table 2. Characteristics of Study Participants

Variable	T2DM (n=111)	Non-DM (n=60)	T1DM (n=10)	p-value
Age (years)	40.3 ± 6.73	31.5 ± 7.18	24.8 ± 6.05	<0.001
Sex (M/F)	23/88	4/56	2/8	0.001
BMI (kg/m²)	23.2 ± 1.40	23.6 ± 1.69	22.7 ± 1.83	0.008
Fasting glucose (mg/dL)	159 (142–193)	87 (82–91)	199 (190–210)	<0.001
HbA1c (%)	8.83 ± 1.68	5.1 ± 0.17	8.4 ± 1.14	<0.001
Insulin	9.8 ± 10.40	7.36 ± 1.38	2.0 ± 0.27	<0.001
C-peptide (ng/mL)	2.18 (1.59–2.53)	2.33 (1.67–2.55)	0.30 (0.20–0.30)	<0.001
HOMA-IR	3.5 (2.8–3.9)	1.5 (1.3–1.7)	1.0 (0.9–1.1)	<0.001

Table 3. Distribution of rs12255372 Genotypes and Alleles

Genotype/Allele	T2DM (n=111)	Non-DM (n=60)	p-value
GG	107 (96.4%)	58 (96.7%)	0.93
GT	4 (3.6%)	2 (3.3%)	
G allele	218 (98.2%)	118 (98.3%)	0.92
T allele	4 (1.8%)	2 (1.7%)	

Table 4. Association Between rs12255372 and T2DM

Genotype/Allele	OR (95% CI)	p-value
GG	Reference	0.93
GT	1.08 (0.19–6.16)	

Table 5. Comparison of C-Peptide Levels according to genotype

Genotype	Median C-peptide(IQR) ng/mL	p-value
GG (n=175)	2.1 (1.5–2.5)	0.157
GT (n=6)	2.4 (2.1–2.6)	

Table 6. Comparison of HOMA-IR according to genotype

Genotype	Median HOMA-IR (IQR)	p-value
GG (n=175)	2.7 (1.6–3.6)	0.609
GT (n=6)	3.0 (1.9–3.7)	

Table 7. Multivariate Logistic Regression Analysis for T2DM

Variable	OR	95% CI	p-value
rs12255372 genotype	0.646	0.070–5.969	0.700
Age	1.323	1.199–1.459	<0.001
BMI	0.752	0.677–0.835	<0.001
Sex	3.077	1.020–9.283	0.046