

Notes on Statistical Theory

Benjamin Nutter

Contents

I	Probability Distributions	7
	The Bernoulli Distribution	9
1	Cumulative Distribution Function	9
2	Expected Values	10
3	Moment Generating Function	10
4	Theorems for the Bernoulli Distribution	11
4.1	Validity of the Distribution	11
4.2	Sum of Bernoulli Random Variables	11
	The Binomial Distribution	13
5	Cumulative Distribution Function	13
6	Expected Values	13
7	Moment Generating Function	16
8	Maximum Likelihood Estimator	17
8.1	Likelihood Function	17
8.2	Log-likelihood Function	18
8.3	MLE for p	18
9	Theorems for the Binomial Distribution	18
9.1	Validity of the Distribution	18
9.2	Sum of Binomial Random Variables	19
9.3	Sum of Bernoulli Random Variables	19
	The Chi-square Distribution	21
	The Exponential Distribution	23
	The Gamma Distribution	25
	The Geometric Distribution	27
	The Hypergeometric Distribution	29

The Multinomial Distribution	31
The Normal Distribution	33
The Poisson Distribution	35
The Skew-Normal Distribution	37
The Uniform Distribution	39
The Weibull Distribution	41
II Frequentist Hypothesis Testing	43
Mantel-Haenszel Test	45
III Supplemental Subjects	47
Chebychev's Theorem	49
Combinations	51
The Correlation Coefficient	53
Covariance	55
Experimental Designs	57
Moments and Moment Generating Functions	59
Summation	61
The Method of Transformations	63
Variance Paramter	65
IV Non-Statistical Proofs	67
The Binomial Theorem	69
Functions	71
The Geometric Series	73

<i>CONTENTS</i>	5
Integraion: Techniques and Theorems	75
Logarithmic and Exponential Functions	77
The Real Number System	79

Part I

Probability Distributions

The Bernoulli Distribution

A random variable is said to have a Bernoulli Distribution with parameter p if its probability mass function is:

$$p(x) = \begin{cases} p^x(1-p)^{1-x}, & x = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

Where p is the probability of a success.

1 Cumulative Distribution Function

$$P(x) = \begin{cases} 0 & x < 0 \\ 1-p & x = 0 \\ 1 & 1 \leq x \end{cases}$$




Figure .1: The graphs on the left and right show a Binomial Probability Distribution and Cumulative Distribution Function, respectively, with $p = .4$. Note that this is identical to a Binomial Distribution with parameters $n = 1$ and $p = .4$.

2 Expected Values

$$\begin{aligned}
 E(X) &= \sum_{i=0}^1 x \cdot p(x) \\
 &= \sum_{i=0}^1 x \cdot p^x (1-p)^{1-x} \\
 &= 0 \cdot p^0 (1-p)^{1-0} + 1 \cdot p^1 (1-p)^{1-1} \\
 &= 0 + p(1-p)^0 \\
 &= p
 \end{aligned}$$

$$\begin{aligned}
 E(X^2) &= \sum_{i=0}^1 x^2 \cdot p(x) \\
 &= \sum_{i=0}^1 x^2 \cdot p^x (1-p)^{1-x} \\
 &= \sum_{i=0}^1 0^2 \cdot p^0 (1-p)^{1-0} + 1^2 \cdot p^1 (1-p)^{1-1} \\
 &= 0 \cdot 1 \cdot 1 + 1 \cdot p \cdot 1 \\
 &= 0 + p \\
 &= p
 \end{aligned}$$

$$\mu = E(X) = p$$

$$\sigma^2 = E(X^2) - E(X)^2 = p - p^2 = p(1-p)$$

3 Moment Generating Function

$$\begin{aligned}
 M_X(t) &= E(e^{tX}) = \sum_{i=0}^1 e^{tx} p(x) = \sum_{i=0}^1 e^{tx} p^x (1-p)^{1-x} \\
 &= e^{t0} p^0 (1-p)^{1-0} + e^{t1} p^1 (1-p)^{1-1} = (1-p) + e^t p = pe^t + (1-p)
 \end{aligned}$$

$$M_X^{(1)}(t) = pe^t$$

$$M_X^{(2)}(t) = pe^t$$

$$E(X) = M_X^{(1)}(0) = pe^0 = p \cdot 1 = p$$

$$E(X^2) = M_X^{(2)}(0) = pe^0 = p$$

$$\mu = E(X) = p$$

$$\sigma^2 = E(X^2) - E(X)^2 = p - p^2 = p(1 - p)$$

4 Theorems for the Bernoulli Distribution

4.1 Validity of the Distribution

$$\sum_{x=0}^1 p^x (1-p)^{1-x} = 1$$

Proof:

$$\sum_{x=0}^1 p^x (1-p)^{1-x} = p^0 (1-p)^1 + p^1 (1-p)^0 = (1-p) + p = 1$$

■

4.2 Sum of Bernoulli Random Variables

Let X_1, X_2, \dots, X_n be independent and identically distributed random variables from a Bernoulli distribution with parameter p . Let $Y = \sum_{i=1}^n X_i$.

Then $Y \sim \text{Binomial}(n, p)$

Proof:

$$\begin{aligned} M_Y(t) &= E(e^{tY}) = E(e^{tX_1} e^{tX_2} \dots e^{tX_n}) = E(e^{tX_1}) E(e^{tX_2}) \dots E(e^{tX_n}) \\ &= (pe^t + (1-p))(pe^t + (1-p)) \dots (pe^t + (1-p)) = (pe^t + (1-p))^n \end{aligned}$$

Which is the mgf of a Binomial random variable with parameters n and p .

Thus, $Y \sim \text{Binomial}(n, p)$. ■

The Binomial Distribution

A random variable is said to follow a Binomial distribution with parameters n and p if its probability mass function is:

$$p(x) = \begin{cases} \binom{n}{x} p^x (1-p)^{n-x}, & x = 0, 1, 2, \dots, n \\ 0 & \text{otherwise} \end{cases}$$

Where n is the number of trials performed and p is the probability of a success on each individual trial.

5 Cumulative Distribution Function

$$P(x) = \begin{cases} 0 & x < 0 \\ \sum_{i=0}^x \binom{n}{i} p^i (1-p)^{n-i} & 0 \leq x = 0, 1, 2, \dots, n \\ 1 & n \leq x \end{cases}$$

A recursive form of the cdf can be derived and has some usefulness in computer applications. With it, one need only initiate the first value and additional cumulative probabilities can be calculated. It is derived as follows:

$$\begin{aligned} F(x+1) &= \binom{n}{x+1} p^{x+1} (1-p)^{n-(x+1)} \\ &= \frac{n!}{(x+1)!(n-(x+1))!} p^{x+1} (1-p)^{n-(x+1)} \\ &= \frac{n!}{(x+1)!(n-x-1)!} p^{x+1} (1-p)^{n-x-1} \\ &= \frac{(n-x)n!}{(x+1)x!(n-x)(n-x-1)!} p \cdot p^x \frac{(1-p)^{n-x}}{(1-p)} \\ &= \frac{(n-x)n!}{(x+1)x!(n-x)!} \cdot \frac{p}{1-p} p^x (1-p)^{n-x} \\ &= \frac{p}{1-p} \cdot \frac{n-x}{x+1} \cdot \frac{n!}{x!(n-x)!} p^x (1-p)^{n-x} \\ &= \frac{p}{1-p} \cdot \frac{n-x}{x+1} \cdot \binom{n}{x} p^x (1-p)^{n-x} \\ &= \frac{p}{1-p} \cdot \frac{n-x}{x+1} \cdot F(x) \end{aligned}$$

6 Expected Values

Let X be a binomial random variable with parameters n and p . The expected value of X is:




Figure .1: plot of chunk unnamed-chunk-7

$$\begin{aligned}
 E(X) &= \sum_{x=0}^n x \cdot p(x) \\
 &= \sum_{x=0}^n x \binom{n}{x} p^x (1-p)^{n-x}
 \end{aligned}$$

For convenience, let $q = (1 - p)$

$$\begin{aligned}
 &= \sum_{x=0}^n x \binom{n}{x} p^x q^{n-x} \\
 &= 0 \cdot \binom{n}{0} p^0 q^n + 1 \cdot \binom{n}{1} p^1 q^{n-1} + \dots + n \binom{n}{n} p^n q^{n-n} \\
 &= 0 + 1 \binom{n}{1} p^1 q^{n-1} + 2 \binom{n}{2} p^2 q^{n-2} + \dots + n \binom{n}{n} p^n q^{n-n} \\
 &= np^1 q^{n-1} + n(n-1)p^2 q^{n-2} + \dots + n(n-1)p^{n-1} q^{n-(n-1)} + np^n \\
 &= np[q^{n-1} + (n-1)pq^{n-2} + \dots + p^{n-1}] \\
 &= np \left[\binom{n-1}{0} p^0 q^{n-1} + \binom{n-1}{1} p^1 q^{(n-1)-1} + \dots + \binom{n-1}{n-1} p^{n-1} q^{n-(n-1)} \right] \\
 &= np \left(\sum_{x=0}^{n-1} \binom{n-1}{x} p^x q^{(n-1)-x} \right)
 \end{aligned}$$

By the Binomial Theorem, $\sum_{x=0}^n \binom{n}{x} a^x b^{n-x} = (a+b)^n$

$$= np(p+q)^{n-1}$$

Resubstituting $(1 - p)$ for q gives us

$$\begin{aligned}
 &= np(p + (1 - p))^{n-1} \\
 &= np(p + 1 - p)^{n-1} \\
 &= np(1)^{n-1} \\
 &= np(1) \\
 &= np
 \end{aligned}$$

The Expected Value of X^2 is:

$$\begin{aligned} E(X^2) &= \sum_{x=0}^n x^2 p(x) \\ &= \sum_{x=0}^n x^2 \binom{n}{x} p^x (1-p)^{n-x} \end{aligned}$$

For convenience, let $q = (1-p)$

$$\begin{aligned} &= \sum_{x=0}^n x^2 \binom{n}{x} p^x q^{n-x} \\ &= 0^2 \frac{n!}{0!(n-0)!} p^0 q^n + 1^2 \frac{n!}{1!(n-1)!} p^1 q^{n-1} + \cdots + n^2 \frac{n!}{n!(n-n)!} p^n q^{n-n} \\ &= 0 + 1 \frac{n!}{(n-1)!} p q^{n-1} + 2 \frac{n!}{1 \cdot (n-2)!} p^2 q^{n-2} + \cdots + n \frac{n!}{(n-1)!(n-n)!} p^n \\ &= np \left[1 \frac{(n-1)!}{(n-1)!} p^0 q^{n-1} + 2 \frac{(n-1)!}{1(n-2)!} p^2 q^{n-2} + \cdots + n \frac{(n-1)!}{(n-1)!(n-n)!} p^{n-1} \right] \\ &= np \left[1 \frac{(n-1)!}{(1-1)!((n-1)-(-1-1))!} p^{1-1} q^{n-1} + \cdots + n \frac{(n-1)!}{(n-1)!((n-1)-(n-1))!} p^{n-1} \right] \\ &= np \sum_{x=1}^n x \binom{n-1}{x-1} p^{x-1} q^{(n-1)-(x-1)} \end{aligned}$$

Let $y = x - 1$ and $n = m + 1$

$\Rightarrow x = y + 1$ and $m = n - 1$

$$\begin{aligned} &= \sum_{y=0}^m (y+1) \binom{m}{y} p^y q^{m-y} \\ &= np \left[\sum_{y=0}^m y \binom{m}{y} p^y q^{m-y} + \binom{m}{y} p^y q^{m-y} \right] \\ &= np \left[\sum_{y=0}^m y \binom{m}{y} p^y q^{m-y} + \sum_{y=0}^m \binom{m}{y} p^y q^{m-y} \right] \end{aligned}$$

$\sum_{y=0}^m y \binom{m}{y} p^y q^{m-y}$ is of the form

of the expected value of Y ,

and $E(Y) = mp = (n-1)p$

$\sum_{y=0}^m \binom{m}{y} p^y q^{m-y}$ is the sum of all

probabilities over the domain of Y ,

which is 1.

$$\begin{aligned} &= np(mp + 1) \\ &= np[(n-1)p + 1] \\ &= np(np - p + 1) \\ &= n^2 p^2 - np^2 + np \end{aligned}$$

The mean of X can be calculated as

$$\mu = E(X) = np$$

And the variance of X can be calculated by

$$\begin{aligned}\sigma^2 &= E(X^2) - E(X)^2 \\ &= n^2 p^2 - np^2 + np - n^2 p^2 \\ &= -np^2 + np \\ &= np(-p + 1) \\ &= np(1 - p)\end{aligned}$$

7 Moment Generating Function

$$\begin{aligned}M_X(t) &= E(e^{tX}) = \sum_{x=0}^n e^{tx} p(x) \\ &= \sum_{x=0}^n e^{tx} \binom{n}{x} p^x (1-p)^{n-x} \\ &= \sum_{x=0}^n \binom{n}{x} e^{tx} p^x (1-p)^{n-x} \\ &= \sum_{x=0}^n \binom{n}{x} (pe^{tx})^x (1-p)^{n-x}\end{aligned}$$

By Binomial Theorem REF

$$\begin{aligned}\sum_{x=0}^n \binom{n}{x} b^x a^{n-x} &= (a + b)^n \\ &= [(1-p) + pe^t]^n\end{aligned}$$

$$M_X^{(1)}(t) = n[(1-p) + pe^t]^{n-1} pe^t$$

$$\begin{aligned}M_X^{(2)}(t) &= n[(1-p) + pe^t]^{n-1} pe^t + n(n-1)[(1-p) + pe^t]^{n-2} (pe^t)^2 \\ &= npe^t[(1-p) + pe^t]^{n-1} + n(n-1)pe^{2t}[(1-p) + pe^t]^{n-2}\end{aligned}$$

$$\begin{aligned}
E(X) &= M_X^{(1)}(0) \\
&= n[(1-p) + pe^0]^{n-1} pe^0 \\
&= n[1-p + p^{n-1}p] \\
&= n(1)^{n-1}p = np
\end{aligned}$$

$$\begin{aligned}
E(X^2) &= M_X^{(2)}(0) = npe^0[(1-p) + pe^0]^{n-1} + n(n-2)pe^{2 \cdot 0}[(1-p) + pe^0]^{n-2} \\
&= np(1-p + p)^{n-2} + n(n-1)p^2(1-p + p^{n-2}) \\
&= np(1)^{n-1} + n(n-1)p^2(1)^{n-2} = np + n(n-1)p^2 = np + (n^2 - n)p^2 \\
&= np + n^2 + n^2p^2 - np^2
\end{aligned}$$

$$\mu = E(X) = np$$

$$\begin{aligned}
\sigma^2 &= E(X^2) - E(X)^2 \\
&= np + n^2p^2 - np^2 - n^2p^2 \\
&= np - np^2 \\
&= np(1-p)
\end{aligned}$$

8 Maximum Likelihood Estimator

Since n is fixed in each Binomial experiment, and must therefore be given, it is unnecessary to develop an estimator for n . The mean and variance can both be estimated from the single parameter p .

Let X be a Binomial random variable with parameter p and n outcomes (x_1, x_2, \dots, x_n) . Let $x_i = 0$ for a failure and $x_i = 1$ for a success. In other words, X is the sum of n Bernoulli trials with equal probability of success and $X = \sum_{i=1}^n x_i$.

8.1 Likelihood Function

$$\begin{aligned}
L(\theta) &= L(x_1, x_2, \dots, x_n | \theta) \\
&= P(x_1 | \theta) P(x_2 | \theta) \cdots P(x_n | \theta) \\
&= [\theta^{x_1} (1-\theta)^{1-x_1}] [\theta^{x_2} (1-\theta)^{1-x_2}] \cdots [\theta^{x_n} (1-\theta)^{1-x_n}] \\
&= \exp_{\theta} \left\{ \sum_{i=1}^n x_i \right\} \exp_{(1-\theta)} \left\{ n - \sum_{i=1}^n x_i \right\} \\
&= \theta^X (1-\theta)^{n-X}
\end{aligned}$$

8.2 Log-likelihood Function

$$\begin{aligned}
 \ell(\theta) &= \ln L(\theta) \\
 &= \ln (\theta^X (1 - \theta)^{n-X}) \\
 &= X \ln(\theta) + (n - X) \ln(1 - \theta)
 \end{aligned}$$

8.3 MLE for p

$$\begin{aligned}
 \frac{d\ell(p)}{dp} &= \frac{X}{p} - \frac{n - X}{1 - p} \\
 0 &= \frac{X}{p} - \frac{n - X}{1 - p} \\
 \frac{X}{p} &= \frac{n - X}{1 - p} \\
 (1 - p)X &= p(n - X) \\
 X - pX &= np - pX \\
 X &= np \\
 \frac{X}{n} &= p
 \end{aligned}$$

So $\hat{p} = \frac{X}{n} = \frac{1}{n} \sum_{i=1}^n x_i$ is the maximum likelihood estimator for p .

9 Theorems for the Binomial Distribution

9.1 Validity of the Distribution

$$\sum_{x=0}^n \binom{n}{x} p^x (1 - p)^{n-x} = 1$$

Proof:

$$\begin{aligned}
 \sum_{x=0}^n \binom{n}{x} p^x (1 - p)^{n-x} &= \\
 \sum_{x=0}^n \binom{n}{x} a^x b^{n-x} (a + b)^n &.
 \end{aligned}$$

See Binomial Theorem REF.

$$\begin{aligned}
 &= \big(p + (1 - p)\big)^n \\
 &= (1)^n \\
 &= 1
 \end{aligned}$$

■

9.2 Sum of Binomial Random Variables

Let X_1, X_2, \dots, X_k be independent random variables where X_i comes from a Binomial distribution with parameters n_i and p . That is $X_i \sim (n_i, p)$. Let $Y = \sum_{i=1}^k X_i$. Then $Y \sim \text{Binomial}(\sum_{i=1}^k n_i, p)$.

Proof:

$$\begin{aligned}
 M_Y(t) &= E(e^{tY}) \\
 &= E(e^{t(X_1 + X_2 + \dots + X_k)}) \\
 &= E(e^{tX_1} e^{tX_2} \dots e^{tX_k}) \\
 &= E(e^{tX_1}) E(e^{tX_2}) \dots E(e^{tX_k}) \\
 &= \prod_{i=1}^k [(1-p) + pe^t]^{n_i} \\
 &= [(1-p) + pe^t]^{\sum_{i=1}^k n_i}
 \end{aligned}$$

\ \ Which is the mgf of a Binomial random variable with parameters $\sum_{i=1}^k n_i$ and p .

Thus $Y \sim \text{Binomial}(\sum_{i=1}^k n_i, p)$. ■

9.3 Sum of Bernoulli Random Variables

Let X_1, X_2, \dots, X_n be independent and identically distributed random variables from a Bernoulli distribution with parameter p . Let $Y = \sum_{i=1}^n X_i$.

Then $Y \sim \text{Binomial}(n, p)$

Proof:

$$\begin{aligned}
 M_Y(t) &= E(e^{tY}) \\
 &= E(e^{tX_1} e^{tX_2} \dots e^{tX_n}) \\
 &= E(e^{tX_1}) E(e^{tX_2}) \dots E(e^{tX_n}) \\
 &= (pe^t + (1-p))(pe^t + (1-p)) \dots (pe^t + (1-p)) \\
 &= (pe^t + (1-p))^n
 \end{aligned}$$

Which is the mgf of a Binomial random variable with parameters n and p . Thus, $Y \sim \text{Binomial}(n, p)$. ■

The Chi-square Distribution

The Exponential Distribution

The Gamma Distribution

The Geometric Distribution

The Hypergeometric Distribution

The Multinomial Distribution

The Normal Distribution

The Poisson Distribution

The Skew-Normal Distribution

The Uniform Distribution

The Weibull Distribution

Part II

Frequentist Hypothesis Testing

Mantel-Haenszel Test

Part III

Supplemental Subjects

Chebychev's Theorem

Combinations

The Correlation Coefficient

Covariance

Experimental Designs

Moments and Moment Generating Functions

Summation

The Method of Transformations

Variance Paramter

Part IV

Non-Statistical Proofs

The Binomial Theorem

Functions

The Geometric Series

Integraion: Techniques and Theorems

Logarithmic and Exponential Functions

The Real Number System