

Table 6.3.2 : Prevalence Rates of the Population with at Least One Non-Communicable Disease by Age Group and Sex, 2024

Age Group (Years)	Total Population			Population who has at least one NCD			Prevalence Rate		
	Total	Male	Female	Total	Male	Female	Total	Male	Female
Sri Lanka	21,779,483	10,510,498	11,268,985	4,185,749	1,725,859	2,459,890	19.2	16.4	21.8
00-04	1,215,095	611,067	604,028	13,673	7,507	6,166	1.1	1.2	1
05-09	1,556,507	788,536	767,971	29,688	16,709	12,979	1.9	2.1	1.7
10-14	1,735,181	880,895	854,286	35,588	20,225	15,363	2.1	2.3	1.8
15-19	1,795,024	907,730	887,294	31,370	16,270	15,100	1.7	1.8	1.7
20-24	1,608,577	790,203	818,374	29,256	13,480	15,776	1.8	1.7	1.9
25-29	1,372,421	662,725	709,696	36,874	16,189	20,685	2.7	2.4	2.9
30-34	1,414,013	682,539	731,474	67,032	28,862	38,170	4.7	4.2	5.2
35-39	1,452,617	705,065	747,552	127,106	56,111	70,995	8.8	8	9.5
40-44	1,602,207	785,827	816,380	239,883	107,296	132,587	15	13.7	16.2
45-49	1,490,573	730,288	760,285	351,226	153,081	198,145	23.6	21	26.1
50-54	1,341,549	650,882	690,667	441,704	185,056	256,648	32.9	28.4	37.2
55-59	1,275,751	603,165	672,586	534,828	217,942	316,886	41.9	36.1	47.1
60-64	1,182,973	547,858	635,115	587,563	239,189	348,374	49.7	43.7	54.9
65-69	991,489	446,651	544,838	558,098	226,178	331,920	56.3	50.6	60.9
70-74	795,319	344,465	450,854	488,211	194,928	293,283	61.4	56.6	65.1
75-79	522,832	214,786	308,046	337,335	129,980	207,355	64.5	60.5	67.3
80-84	268,326	102,783	165,543	176,798	63,951	112,847	65.9	62.2	68.2
85-89	111,009	39,731	71,278	71,442	24,326	47,116	64.4	61.2	66.1
90-94	39,232	12,619	26,613	23,568	7,251	16,317	60.1	57.5	61.3
95 & Over	8,788	2,683	6,105	4,506	1,328	3,178	51.3	49.5	52.1

*Excluding the roofless population

Analysis by age group confirms that the prevalence rate of non-communicable diseases (NCDs) rises sharply for both males and females from the age of 35 onwards. According to the data, 94 percent of the population reporting at least one NCD belongs to age groups over 35 years. The prevalence rate in the 20–34 age group remains at a relatively low level, ranging from 1.8 percent to 4.7 percent. However, the prevalence rate for the 40–44 age group stands at 15 percent, and by age 55–59, this figure nearly triples to 41.9 percent. Overall, the rate increases progressively from the 30–34 age group up to the 80–84 age group, after which it shows a slight decline.

When considering each age group individually, it is clear that after the age of 60, more than half of the population in every age category lives with at least one non-communicable disease. By the 80–84 age group, it is reported that two-thirds of the population (65.9%) have at least one NCD.