

Table 6.3.4 : Prevalence Rates of Non-Communicable Diseases by District, 2024

District	Total Population	Diabetes	High Cholesterol	High Blood Pressure	Heart Disease	Kidney Disease	Thalassemia	Cancer	Stroke	Asthma	Epilepsy
Sri Lanka*	21,779,483	8.5	8.2	10.1	2.5	0.8	0.1	0.4	0.6	1.8	0.3
Colombo	2,374,869	10.7	9.1	10.5	2.5	0.5	0.1	0.3	0.4	1.2	0.2
Gampaha	2,435,913	10.5	9.6	10.7	2.5	0.6	0.1	0.4	0.5	1.2	0.2
Kalutara	1,305,708	9.2	8.9	10.2	2.4	0.6	0.1	0.4	0.6	1.2	0.3
Kandy	1,461,755	8.8	8.2	11.6	2.7	0.6	0.1	0.4	0.6	1.9	0.3
Matale	526,849	7.8	7.9	11.3	2.5	1.1	0.1	0.4	0.6	2.0	0.3
Nuwara Eliya	725,251	5.4	4.9	10.2	2.7	0.6	0.1	0.4	0.9	2.2	0.5
Galle	1,097,225	8.0	9.0	9.8	2.7	0.6	0.1	0.4	0.6	1.9	0.3
Matara	837,756	8.4	9.6	10.3	3.0	0.7	0.1	0.4	0.5	2.5	0.4
Hambantota	671,383	8.0	9.5	9.0	2.5	0.7	0.1	0.4	0.6	2.2	0.3
Jaffna	594,699	10.4	9.9	10.5	2.1	0.6	0.1	0.4	0.6	2.4	0.5
Mannar	123,752	7.7	6.6	8.9	2.1	0.8	0.1	0.3	0.5	2.0	0.4
Vavuniya	172,280	7.5	8.2	8.9	2.5	2.4	0.1	0.3	0.7	3.2	0.5
Mullaitivu	122,617	6.9	7.1	8.0	2.0	1.7	0.1	0.4	0.7	4.2	0.7
Kilinochchi	136,710	6.5	7.3	7.4	1.9	1.0	0.1	0.3	0.5	3.7	0.6
Batticaloa	595,889	7.1	6.3	7.9	1.8	0.5	0.1	0.2	0.6	4.4	0.4
Ampara	744,531	8.3	7.8	9.0	2.2	0.8	0.1	0.3	0.6	2.4	0.3
Trincomalee	442,722	6.7	6.1	7.0	1.6	0.7	0.1	0.2	0.5	1.6	0.3
Kurunegala	1,768,023	7.9	7.7	10.2	2.4	1.1	0.1	0.5	0.7	1.3	0.2
Puttalam	818,703	8.2	8.3	9.4	2.3	0.8	0.1	0.4	0.6	1.6	0.2
Anuradhapura	959,901	7.2	7.5	9.7	2.5	1.5	0.2	0.4	0.6	1.5	0.2
Polonnaruwa	447,503	7.8	8.6	10.0	2.8	1.9	0.1	0.4	0.6	2.3	0.2
Badulla	872,239	6.9	6.3	10.3	2.5	1.1	0.1	0.4	0.6	2.5	0.4
Moneragala	527,532	6.6	6.7	8.7	2.4	1.1	0.1	0.5	0.6	2.3	0.3
Ratnapura	1,145,252	7.5	7.5	9.7	2.3	0.8	0.1	0.4	0.7	2.1	0.4
Kegalle	870,421	8.6	8.9	11.7	2.4	0.5	0.1	0.4	0.7	1.7	0.3

*Excluding the roofless population

The highest prevalence rate for heart disease was reported in the Matara district (3%). Regarding the distribution of kidney disease, Vavuniya (2.4%), Polonnaruwa (1.9%), Mullaitivu (1.7%), and Anuradhapura (1.5%) districts report higher values compared to other districts. A prevalence rate of 0.2 percent for thalassemia was reported in the Anuradhapura district, while all other districts reported a rate of 0.1 percent.

Cancer prevalence rates range between 0.2 percent and 0.5 percent; the lowest rates were reported in Batticaloa and Trincomalee districts, while the highest rates were reported in Kurunegala and Moneragala districts. The lowest prevalence rate for stroke was reported in the Colombo district (0.4%), while the highest was reported in the Nuwara Eliya district (0.9%). The highest prevalence rates for asthma were reported in Batticaloa (4.4%), Mullaitivu (4.2%), and Kilinochchi (3.7%) districts. For epilepsy, several districts reported a minimum prevalence rate of 0.2 percent, while the highest rate of 0.7 percent was reported in the Mullaitivu district.