

 style =
width :
75 <
/section ><
!-
- <
section ><
h4 >
Splinesanddata <
/h4 ><
divclass =
row ><
divclass =
"columnnleft" ><
imgsrc =
"../..Figures/terrain2.png" style =
width :
100 <
/div ><
divclass =
"columnnright" ><
imgsrc =
"../..Figures/levels.jpg" style =
width :
55 <
imgsrc =
"../..Figures/mixed2.jpg" style =
width :
45 <
/div ><
/div ><
small ><
ahref =
"https :
//doi.org/10.1016/j.cag.2015.03.006" >
Locallyrefinedsplinesurfacesforrepresentationofterraindata(2015) <
/a ><
/br >
SINTEF, Norway <
/small ><
/section >
-- ><
/section >
 (f, g)
 C_r
 $f - g = h \ell^{r+1} \Leftrightarrow f - g \in \langle \ell^{r+1} \rangle$.
 $\Delta \subseteq$
 R^2
 $\dim S_d^r(\Delta) \leq$
 $d + 22 +$
 $f_1^0 d + 2 - (r + 1)2 -$
 $\sum_{i, \tilde{t}_i=1}^{f_0^0} d + 2 - (r + 1)2$
 $- \sum_{i=1, \tilde{t}_i \geq 2}^{f_0^0} \left[\tilde{t}_i d + 2 - (r + 1)2 - \tilde{b}_i d + 2 - \tilde{\Omega}_i 2 - \tilde{a}_i d + 2 - (\tilde{\Omega}_i + 1)2 \right]$
 $\tilde{t}_i =$
 $\tilde{\Omega}_i = \lfloor \frac{\tilde{t}_i r}{\tilde{t}_i - 1} \rfloor + 1, \tilde{a}_i = \tilde{t}_i(r + 1) + (1 - \tilde{t}_i)\tilde{\Omega}_i,$
 $\tilde{b}_i = \tilde{t}_i - 1 - \tilde{a}_i.$
 $10^{>}$
Splinespacesontriangulations <
/a >
, 2014. <
/br >
D.Toshniwal and N.Villamizar <
/br >
128209; nbsp; <
ahref =
"https :
//doi.org/10.1016/j.aam.2022.102412" >
Algebraic methods for supersmooth splinespaces <
/a >
, Advances in applied mathematics, 142, 2023. <
/p ><
divclass =
fixed ><
imgsrc =
"../..Figures/Sc3.png" style =
width :
55 <
/div ><
/section ><
section ><
h4 >
Hierarchical box splines <
/h4 >