

# Chapter 12: Variance of Discrete RVs

Meike Niederhausen and Nicky Wakim

2023-10-23

# Table of contents

- Learning Objectives
- Let's start building the variance through expected values of functions
- What is the expected value of a function?
- Let's revisit the card example (1/2)
- Let's revisit the card example (2/2)
- Variance of a RV
- Let's calculate the variance and prove it!
- (break) Some Important Variance and Expected Values Results
  - Variance of a function with a single RV
  - Important results for *independent* RVs
  - Variance of sum of independent discrete RVs
  - Corollaries
  - Let's revisit our ghost problems *without replacement*
  - Let's revisit our ghost problems *with replacement*
  - Back to our hotel example from Chapter 11

# Learning Objectives

1. Calculate the variance and standard deviation of discrete random variables
2. Calculate the variance of sums of discrete random variables
3. Calculate the variance of functions of discrete random variables

# Let's start building the variance through expected values of functions

## Example 1

Let  $g$  be a function and let  $g(x) = ax + b$ , for real-valued constants  $a$  and  $b$ . What is  $\mathbb{E}[g(X)]$ ?

# What is the expected value of a function?

Definition: Expected value of function of RV

For any function  $g$  and discrete r.v.  $X$ , the expected value of  $g(X)$  is

$$\mathbb{E}[g(X)] = \sum_{\{\text{all } x\}} g(x)p_X(x).$$

# Let's revisit the card example (1/2)

## Example 2

Suppose you draw 2 cards from a standard deck of cards *with* replacement. Let  $X$  be the number of hearts you draw.

1. Find  $\mathbb{E}[X^2]$ .

Recall Binomial RV with  $n = 2$ :

$$p_X(x) = \binom{2}{x} p^x (1-p)^{2-x} \text{ for } x = 0, 1, 2$$

# Let's revisit the card example (2/2)

## Example 2

Suppose you draw 2 cards from a standard deck of cards *with* replacement. Let  $X$  be the number of hearts you draw.

2. Find  $\mathbb{E}\left[\left(X - \frac{1}{2}\right)^2\right]$ .

Recall Binomial RV with  $n = 2$ :

$$p_X(x) = \binom{2}{x} p^x (1-p)^{2-x} \text{ for } x = 0, 1, 2$$

# Variance of a RV

## Definition: Variance of RV

The variance of a r.v.  $X$ , with (finite) expected value  $\mu_X = \mathbb{E}[X]$  is

$$\sigma_X^2 = \text{Var}(X) = \mathbb{E}[(X - \mu_X)^2] = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

## Definition: Standard deviation of RV

The standard deviation of a r.v.  $X$  is

$$\sigma_X = \text{SD}(X) = \sqrt{\sigma_X^2} = \sqrt{\text{Var}(X)}.$$



# Let's calculate the variance and prove it!

## Lemma 6: "Computation formula" for Variance

The variance of a r.v.  $X$ , can be computed as

$$\begin{aligned}\sigma_X^2 &= \text{Var}(X) \\ &= \mathbb{E}[X^2] - \mu_X^2 \\ &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2\end{aligned}$$

**(break) Some Important Variance and  
Expected Values Results**

# Variance of a function with a single RV

## Lemma 7

For a r.v.  $X$  and constants  $a$  and  $b$ ,

$$\text{Var}(aX + b) = a^2 \text{Var}(X).$$

Proof will be exercise in homework. It's fun! In a mathy kinda way.

# Important results for *independent* RVs

## Theorem 8

For independent r.v.'s  $X$  and  $Y$ , and functions  $g$  and  $h$ ,

$$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)].$$

## Corollary 1

For independent r.v.'s  $X$  and  $Y$ ,

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y].$$

# Variance of sum of independent discrete RVs

## Theorem 9: Variance of sum of independent discrete r.v.'s

For independent discrete r.v.'s  $X_i$  and constants  $a_i, i = 1, 2, \dots, n$ ,

$$\text{Var}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i^2 \text{Var}(X_i).$$

# Corollaries

## Corollary 2

For independent discrete r.v.'s  $X_i, i = 1, 2, \dots, n$ ,

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i).$$

## Corollary 3

For independent identically distributed (i.i.d.) discrete r.v.'s  $X_i, i = 1, 2, \dots, n$ ,

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = n \text{Var}(X_1).$$

# Let's revisit our ghost problems *without replacement*

## Example 3.1

The ghost is trick-or-treating at a different house now. In this case it is known that the bag of candy has 10 chocolates, 20 lollipops, and 30 laffy taffies. The ghost grabs a handful of five pieces of candy. What is the variance for the number of chocolates the ghost takes? Let's solve this for the cases ***without*** replacement.

Recall probability without replacement:

$$p_X(x) = \frac{\binom{K}{x} \binom{N-K}{n-x}}{\binom{N}{n}}$$

# Let's revisit our ghost problems *with replacement*

## Example 3.2

The ghost is trick-or-treating at a different house now. In this case it is known that the bag of candy has 10 chocolates, 20 lollipops, and 30 laffy taffies. The ghost grabs a handful of five pieces of candy. What is the variance for the number of chocolates the ghost takes? Let's solve this for the cases ***with*** replacement.

Recall probability with replacement:

$$p_X(x) = \binom{n}{k} p^k (1 - p)^{n-k}$$



# Back to our hotel example from Chapter 11

## Example 4

A tour group is planning a visit to the city of Minneapolis and needs to book 30 hotel rooms. The average price of a room is \$200 with standard deviation \$10. In addition, there is a 10% tourism tax for each room. What is the **standard deviation** of the cost for the 30 hotel rooms?

Problem to do at home if we don't have enough time.

